

CareSource Ohio  
 HEDIS Annual Rates: 2012-2014

Measure/Data Element	2014	2013	2012	2014 National 75th Percentile
<b>Effectiveness of Care: Prevention and Screening</b>				
<b>Adult BMI Assessment (aba)</b>	71.78%	58.15%	53.77%	85.23%
<b>Weight Assessment and Counseling for Nutrition and Physical Activity for Children/Adolescents (wcc)</b>				
<i>BMI Percentile</i>	34.95%	24.82%	17.76%	73.72%
<i>Counseling for Nutrition</i>	51.46%	37.47%	34.31%	69.21%
<i>Counseling for Physical Activity</i>	46.84%	24.09%	19.95%	60.82%
<b>Childhood Immunization Status (cis)</b>				
<i>DTaP</i>	71.53%	74.94%	74.94%	84.07%
<i>IPV</i>	89.78%	88.56%	88.56%	93.92%
<i>MMR</i>	89.78%	88.81%	88.81%	93.29%
<i>HiB</i>	89.78%	89.05%	89.05%	94.04%
<i>Hepatitis B</i>	85.16%	89.54%	89.54%	93.43%
<i>VZV</i>	89.78%	88.32%	88.32%	93.16%
<i>Pneumococcal Conjugate</i>	73.97%	76.40%	76.40%	84.95%
<i>Hepatitis A</i>	82.00%	64.72%	64.72%	88.54%
<i>Rotavirus</i>	67.64%	60.58%	60.58%	74.31%
<i>Influenza</i>	44.77%	41.36%	41.36%	58.88%
<i>Combination #2</i>	64.72%	70.07%	70.07%	79.72%
<i>Combination #3</i>	60.34%	65.45%	65.45%	77.78%
<i>Combination #4</i>	57.42%	50.61%	50.61%	72.51%
<i>Combination #5</i>	51.82%	48.18%	48.18%	64.48%
<i>Combination #6</i>	33.33%	34.06%	34.06%	50.69%
<i>Combination #7</i>	49.88%	38.44%	38.44%	61.42%
<i>Combination #8</i>	31.87%	27.74%	27.74%	47.73%
<i>Combination #9</i>	29.93%	26.52%	26.52%	43.93%
<i>Combination #10</i>	28.95%	21.41%	21.41%	41.85%
<b>Immunizations for Adolescents (ima)</b>				
<i>Meningococcal</i>	78.10%	60.34%	59.85%	82.39%
<i>Tdap/Td</i>	79.08%	76.64%	78.10%	90.39%
<i>Combination #1</i>	77.86%	59.12%	56.93%	80.90%
<b>Human Papillomavirus Vaccine for Female Adolescents (hpv)</b>	18.49%	18.25%	16.79%	23.62%
<b>Lead Screening in Children (lsc)</b>	67.15%	66.91%	64.36%	80.83%
<b>Breast Cancer Screening (bcs)</b>	53.97%	46.64%	46.03%	65.12%
<b>Cervical Cancer Screening (ccs)</b>	70.80%	72.99%	68.61%	NR
<b>Non-Recommended Cervical Cancer Screening in Adolescent Females (ncs)</b>	3.29%	NA	NA	NR
<b>Chlamydia Screening in Women (chl)</b>				
<i>16-20 Years</i>	51.59%	53.84%	56.38%	58.68%
<i>21-24 Years</i>	63.45%	65.04%	65.00%	69.23%
<i>Total</i>	56.82%	58.67%	60.00%	62.75%

DNE = Did Not exist: Measure did not exist in this year.

NA = Not Applicable

NR = Not Reportable: Not enough data was collected to report this measure.

Notes: Rates were extracted from the IDSS results file for each year listed.

CareSource Ohio  
HEDIS Annual Rates: 2012-2014

Measure/Data Element	2014	2013	2012	2014 National 75th Percentile
<b>Effectiveness of Care: Respiratory Conditions</b>				
<b>Appropriate Testing for Children with Pharyngitis (cwp)</b>	68.72%	71.78%	71.89%	77.96%
<b>Appropriate Treatment for Children With URI (uri)</b>	81.81%	81.11%	80.74%	91.21%
<b>Avoidance of Antibiotic Treatment in Adults with Acute Bronchitis (aab)</b>	16.78%	16.43%	17.15%	30.54%
<b>Use of Spirometry Testing in the Assessment and Diagnosis of COPD (spr)</b>	28.14%	28.50%	31.79%	36.73%
<b>Pharmacotherapy Management of COPD Exacerbation (pce)</b>				
<i>Systemic Corticosteroid</i>	75.08%	73.27%	71.86%	74.94%
<i>Bronchodilator</i>	88.09%	87.05%	84.83%	87.61%
<b>Use of Appropriate Medications for People With Asthma (asm)</b>				
<i>5-11 Years</i>	90.39%	89.00%	86.96%	93.59%
<i>12-18 Years</i>	86.31%	85.38%	82.62%	89.52%
<i>19-50 Years</i>	72.09%	70.77%	63.63%	80.12%
<i>51-64 Years</i>	65.11%	64.72%	58.77%	77.21%
<i>Total</i>	82.96%	58.67%	60.00%	87.26%
<b>Medication Management for People With Asthma (mma)</b>				
<i>5-11 Years - Medication Compliance 50%</i>	53.21%	49.90%	51.00%	NA
<i>5-11 Years - Medication Compliance 75%</i>	29.58%	25.83%	29.21%	32.32%
<i>12-18 Years - Medication Compliance 50%</i>	50.11%	48.60%	45.92%	NA
<i>12-18 Years - Medication Compliance 75%</i>	25.87%	24.10%	23.55%	30.58%
<i>19-50 Years - Medication Compliance 50%</i>	61.48%	56.71%	52.51%	NA
<i>19-50 Years - Medication Compliance 75%</i>	36.89%	34.14%	32.59%	40.36%
<i>51-64 Years - Medication Compliance 50%</i>	79.96%	78.28%	66.07%	NA
<i>51-64 Years - Medication Compliance 75%</i>	57.63%	56.97%	49.74%	56.67%
<i>Total - Medication Compliance 50%</i>	55.40%	52.07%	50.43%	NA
<i>Total - Medication Compliance 75%</i>	31.42%	28.32%	29.06%	34.96%
<b>Asthma Medication Ratio (amr)</b>				
<i>5-11 Years</i>	77.30%	89.00%	86.96%	82.16%
<i>12-18 Years</i>	65.35%	85.38%	82.62%	70.95%
<i>19-50 Years</i>	47.34%	70.77%	63.63%	55.36%
<i>51-64 Years</i>	46.60%	64.72%	58.77%	58.93%
<i>Total</i>	64.22%	58.67%	60.00%	70.88%
<b>Effectiveness of Care: Cardiovascular</b>				
<b>Cholesterol Management for Patients With Cardiovascular Conditions (cmc)</b>				
<i>LDL-C Screening Performed</i>	80.54%	82.48%	82.48%	84.91%
<i>LDL-C Control (&lt;100 mg/dL)</i>	32.60%	32.60%	32.60%	47.64%
<b>Controlling High Blood Pressure (cbp)</b>	52.07%	52.07%	51.09%	63.76%
<b>Persistence of Beta-Blocker Treatment After a Heart Attack (pbh)</b>	93.83%	88.10%	87.27%	91.18%

DNE = Did Not exist: Measure did not exist in this year.

NA = Not Applicable

NR = Not Reportable: Not enough data was collected to report this measure.

Notes: Rates were extracted from the IDSS results file for each year listed.

**CareSource Ohio**  
**HEDIS Annual Rates: 2012-2014**

Measure/Data Element	2014	2013	2012	2014 National 75th Percentile
<b>Effectiveness of Care: Diabetes</b>				
<b>Comprehensive Diabetes Care (cdc)</b>				
<i>Hemoglobin A1c (HbA1c) Testing</i>	80.39%	79.24%	79.13%	87.59%
<i>HbA1c Poor Control (&gt;9.0%)</i>	63.37%	53.63%	56.52%	36.52%
<i>HbA1c Control (&lt;8.0%)</i>	34.68%	37.72%	41.22%	52.89%
<i>HbA1c Control (&lt;7.0%)</i>	22.87%	27.49%	31.39%	40.26%
<i>Eye Exam (Retinal) Performed</i>	52.51%	57.27%	45.22%	63.14%
<i>LDL-C Screening Performed</i>	69.37%	82.48%	69.91%	80.18%
<i>LDL-C Control (&lt;100 mg/dL)</i>	27.23%	32.60%	24.00%	40.44%
<i>Medical Attention for Nephropathy</i>	74.72%	74.05%	73.91%	83.11%
<i>Blood Pressure Control (&lt;140/80 mm Hg)</i>	39.55%	38.26%	38.26%	45.50%
<i>Blood Pressure Control (&lt;140/90 mm Hg)</i>	56.40%	57.22%	57.22%	70.07%
<b>Effectiveness of Care: Musculoskeletal</b>				
<b>Disease Modifying Anti-Rheumatic Drug Therapy in Rheumatoid Arthritis (art)</b>	72.54%	71.89%	72.93%	77.17%
<b>Use of Imaging Studies for Low Back Pain (lbp)</b>	70.92%	73.51%	71.55%	78.53%
<b>Effectiveness of Care: Behavioral Health</b>				
<b>Antidepressant Medication Management (amm)</b>				
<i>Effective Acute Phase Treatment</i>	46.04%	48.93%	47.96%	54.31%
<i>Effective Continuation Phase Treatment</i>	29.82%	32.25%	29.96%	38.23%
<b>Follow-Up Care for Children Prescribed ADHD Medication (add)</b>				
<i>Initiation Phase</i>	56.18%	51.86%	51.54%	46.99%
<i>Continuation and Maintenance (C&amp;M) Phase</i>	66.56%	60.89%	60.37%	57.55%
<b>Follow-Up After Hospitalization for Mental Illness (fuh)</b>				
<i>30-Day Follow-Up</i>	76.97%	70.64%	68.44%	74.09%
<i>7-Day Follow-Up</i>	54.49%	46.01%	40.17%	54.45%
<b>Diabetes Screening for People With Schizophrenia or Bipolar Disorder Who Are Using Antipsychotic Medication (ssd)</b>	82.56%	75.21%	DNE	82.88%
<b>Diabetes Monitoring for People With Diabetes and Schizophrenia (smd)</b>	60.69%	68.11%	DNE	73.76%
<b>Cardiovascular Monitoring for People With Cardiovascular Disease and Schizophrenia (smc)</b>	78.23%	72.99%	DNE	84.21%
<b>Adherence to Antipsychotic Medications for Individuals With Schizophrenia (saa)</b>	57.35%	63.32%	DNE	67.13%
<b>Effectiveness of Care: Medication Management</b>				
<b>Annual Monitoring for Patients on Persistent Medications (mpm)</b>				
<i>ACE Inhibitors or ARBs</i>	87.14%	87.23%	87.52%	89.94%
<i>Digoxin</i>	90.46%	93.19%	92.63%	94.12%
<i>Diuretics</i>	86.92%	87.27%	86.92%	90.57%
<i>Anticonvulsants</i>	61.72%	62.01%	64.67%	69.53%
<i>Total</i>	85.02%	58.67%	60.00%	88.25%

DNE = Did Not exist: Measure did not exist in this year.

NA = Not Applicable

NR = Not Reportable: Not enough data was collected to report this measure.

Notes: Rates were extracted from the IDSS results file for each year listed.

**CareSource Ohio**  
**HEDIS Annual Rates: 2012-2014**

Measure/Data Element	2014	2013	2012	2014 National 75th Percentile
<b>Access/Availability of Care</b>				
<b>Adults' Access to Preventive/Ambulatory Health Services (aap)</b>				
<i>20-44 Years</i>	84.40%	84.91%	85.54%	86.21%
<i>45-64 Years</i>	90.74%	90.72%	91.16%	90.98%
<i>65+ Years</i>	91.59%	90.85%	91.84%	90.70%
<i>Total</i>	85.94%	58.67%	60.00%	88.11%
<b>Children and Adolescents' Access to Primary Care Practitioners (cap)</b>				
<i>12-24 Months</i>	96.32%	96.46%	96.57%	97.86%
<i>25 Months - 6 Years</i>	86.17%	88.28%	88.53%	91.73%
<i>7-11 Years</i>	87.96%	90.39%	90.13%	93.50%
<i>12-19 Years</i>	87.30%	89.88%	89.63%	92.17%
<b>Annual Dental Visit (adv)</b>				
<i>2-3 Years</i>	28.35%	29.70%	30.10%	43.98%
<i>4-6 Years</i>	57.43%	58.13%	59.02%	68.85%
<i>7-10 Years</i>	60.74%	61.07%	62.04%	71.57%
<i>11-14 Years</i>	57.25%	57.35%	57.57%	65.61%
<i>15-18 Years</i>	50.86%	50.46%	50.17%	56.52%
<i>19-21 Years</i>	36.40%	37.37%	37.68%	41.02%
<i>Total</i>	52.44%	58.67%	60.00%	61.13%
<b>Initiation and Engagement of AOD Dependence Treatment (iet)</b>				
<i>Initiation of AOD Treatment: 13-17 Years</i>	36.78%	43.54%	33.50%	47.47%
<i>Engagement of AOD Treatment: 13-17 Years</i>	6.44%	19.70%	10.19%	23.88%
<i>Initiation of AOD Treatment: 18+ Years</i>	29.85%	41.00%	34.83%	43.29%
<i>Engagement of AOD Treatment: 18+ Years</i>	6.64%	11.30%	4.47%	14.17%
<i>Initiation of AOD Treatment: Total</i>	30.26%	41.23%	34.73%	43.48%
<i>Engagement of AOD Treatment: Total</i>	6.63%	12.06%	4.91%	14.97%
<b>Prenatal and Postpartum Care (ppc)</b>				
<i>Timeliness of Prenatal Care</i>	87.35%	87.35%	87.35%	89.62%
<i>Postpartum Care</i>	60.58%	60.58%	63.75%	69.47%
<b>Call Answer Timeliness (cat)</b>				
	92.51%	95.53%	95.07%	91.85%
<b>Utilization</b>				
<b>Frequency of Ongoing Prenatal Care (fpc)</b>				
<i>&lt;21 Percent</i>	4.14%	4.14%	2.92%	15.71%
<i>21-40 Percent</i>	6.33%	6.33%	2.43%	8.83%
<i>41-60 Percent</i>	4.14%	4.14%	4.87%	10.82%
<i>61-80 Percent</i>	12.17%	12.17%	12.90%	17.35%
<i>81+ Percent</i>	73.24%	73.24%	76.89%	71.34%
<b>Well-Child Visits in the First 15 Months of Life (w15)</b>				
<i>0 Visits</i>	3.16%	2.92%	2.43%	2.64%
<i>1 Visit</i>	1.70%	3.16%	0.97%	2.35%
<i>2 Visits</i>	2.19%	1.22%	3.89%	3.70%
<i>3 Visits</i>	5.35%	4.62%	4.14%	6.34%
<i>4 Visits</i>	9.98%	6.33%	10.46%	11.88%
<i>5 Visits</i>	17.27%	9.98%	11.19%	18.73%
<i>6+ Visits</i>	60.34%	71.78%	66.91%	69.75%
<b>Well-Child Visits in the Third, Fourth, Fifth and Sixth Years of Life (w34)</b>				
	71.53%	66.91%	66.67%	77.26%
<b>Adolescent Well-Care Visits (awc)</b>				
	41.12%	39.05%	45.50%	59.21%

DNE = Did Not exist: Measure did not exist in this year.

NA = Not Applicable

NR = Not Reportable: Not enough data was collected to report this measure.

Notes: Rates were extracted from the IDSS results file for each year listed.