

# ACTEMRA SQ

## Products Affected

- ACTEMRA ACTPEN
- ACTEMRA SUBCUTANEOUS
- TYENNE AUTOINJECTOR
- TYENNE SUBCUTANEOUS

PA Criteria	Criteria Details
Exclusion Criteria	PENDING CMS REVIEW
Required Medical Information	PENDING CMS REVIEW
Age Restrictions	PENDING CMS REVIEW
Prescriber Restrictions	PENDING CMS REVIEW
Coverage Duration	PENDING CMS REVIEW
Other Criteria	PENDING CMS REVIEW
Indications	PENDING CMS REVIEW
Off-Label Uses	PENDING CMS REVIEW
Part B Prerequisite	PENDING CMS REVIEW

# ACTIMMUNE

---

## Products Affected

- ACTIMMUNE

PA Criteria	Criteria Details
Exclusion Criteria	PENDING CMS REVIEW
Required Medical Information	PENDING CMS REVIEW
Age Restrictions	PENDING CMS REVIEW
Prescriber Restrictions	PENDING CMS REVIEW
Coverage Duration	PENDING CMS REVIEW
Other Criteria	PENDING CMS REVIEW
Indications	PENDING CMS REVIEW
Off-Label Uses	PENDING CMS REVIEW
Part B Prerequisite	PENDING CMS REVIEW

## ACYCLOVIR (TOPICAL)

---

### Products Affected

- *acyclovir topical ointment*

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	Diagnosis
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	12 months
Other Criteria	N/A
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

# ADALIMUMAB

## Products Affected

- CYLTEZO(CF) PEN
- CYLTEZO(CF) PEN CROHN'S-UC-HS
- CYLTEZO(CF) PEN PSORIASIS-UV
- CYLTEZO(CF) SUBCUTANEOUS SYRINGE KIT 10 MG/0.2 ML, 20 MG/0.4 ML, 40 MG/0.4 ML, 40 MG/0.8 ML
- HUMIRA (PREFERRED NDCS STARTING WITH 00074) SUBCUTANEOUS SYRINGE KIT 40 MG/0.8 ML
- HUMIRA PEN (PREFERRED NDCS STARTING WITH 00074)
- HUMIRA(CF) (PREFERRED NDCS STARTING WITH 00074) SUBCUTANEOUS SYRINGE KIT 10 MG/0.1 ML, 20 MG/0.2 ML, 40 MG/0.4 ML
- HUMIRA(CF) PEN (PREFERRED NDCS NDCS STARTING WITH 00074) SUBCUTANEOUS PEN INJECTOR KIT 40 MG/0.4 ML, 80 MG/0.8 ML
- HUMIRA(CF) PEN CROHNS-UC-HS (PREFERRED NDCS NDCS STARTING WITH 00074)
- HUMIRA(CF) PEN PEDIATRIC UC (ONLY NDCS STARTING WITH 00074)
- HUMIRA(CF) PEN PSOR-UV-ADOL HS (PREFERRED NDCS NDCS STARTING WITH 00074)
- YUFLYMA(CF) AI CROHN'S-UC-HS
- YUFLYMA(CF) AUTOINJECTOR SUBCUTANEOUS AUTO-INJECTOR, KIT 40 MG/0.4 ML, 80 MG/0.8 ML
- YUFLYMA(CF) SUBCUTANEOUS SYRINGE KIT 20 MG/0.2 ML, 40 MG/0.4 ML

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	Concurrent use with another biologic DMARD or targeted synthetic DMARD.
<b>Required Medical Information</b>	Diagnosis, concurrent medications, previous therapies tried
<b>Age Restrictions</b>	Crohn's disease (CD), 6 or older (initial therapy only). Ulcerative colitis (UC), 5 or older (initial therapy only), PP-18 years and older (initial therapy only)
<b>Prescriber Restrictions</b>	Initial therapy only for all dx-RA/JIA/JRA/Ankylosing spondylitis, prescribed by or in consultation with rheumatologist. Psoriatic arthritis (PsA), prescribed by or in consultation with a rheumatologist or dermatologist. Plaque psoriasis (PP), prescribed by or in consultation with a dermatologist. UC/ CD, prescribed by or in consultation with a gastroenterologist. HS - dermatologist.UV-ophthalmologist

PA Criteria	Criteria Details
<b>Coverage Duration</b>	Approve through end of plan year
<b>Other Criteria</b>	<p>INITIAL THERAPY: CROHN'S DISEASE (CD) [one of A, B, C, or D]: A) tried or currently taking corticosteroid (CS) or CS is contraindicated, B) tried one other conventional systemic therapy (e.g., azathioprine, 6-mercaptopurine [6-MP], methotrexate [MTX], certolizumab, infliximab, ustekinumab, vedolizumab), C) had ileocolonic resection, or D) enterocutaneous (perianal or abdominal) or rectovaginal fistulas.</p> <p>JUVENILE IDIOPATHIC ARTHRITIS (JIA)/JRA (one of A, B, C, D, or E): A) tried one other systemic therapy (e.g., MTX, sulfasalazine, leflunomide, NSAID), B) tried a biologic (e.g., etanercept, abatacept, infliximab, anakinra, tocilizumab), C) will be starting on adalimumab concurrently with MTX, sulfasalazine, or leflunomide, D) patient has absolute contraindication to MTX, sulfasalazine, or leflunomide, or E) patient has aggressive disease.</p> <p>HIDRADENITIS SUPPURATIVA (HS): tried one other therapy (e.g., intralesional or oral CS, systemic antibiotics, isotretinoin).</p> <p>PLAQUE PSORIASIS (PP) [A or B]: A) tried at least one traditional systemic agent (e.g., MTX, cyclosporine (CSA), acitretin, PUVA) for at least 3 months, unless intolerant (Note: a trial of a biologic will also count) or B) contraindication to MTX.</p> <p>RHEUMATOID ARTHRITIS (RA): tried one conventional synthetic DMARD for at least 3 months (Note: a 3-month trial of a biologic will also count).</p> <p>ULCERATIVE COLITIS (A or B): A) tried a systemic therapy (e.g., 6-MP, azathioprine, CSA, tacrolimus, infliximab, golimumab SC, or a CS) or B) has pouchitis and tried therapy with an antibiotic, probiotic, CS enema, or mesalamine (Rowasa) enema.</p> <p>CONTINUATION THERAPY: ALL INDICATIONS: patient had a response to therapy.</p>
<b>Indications</b>	All FDA-approved Indications.
<b>Off-Label Uses</b>	N/A
<b>Part B Prerequisite</b>	No

# ADBRY

## Products Affected

- ADBRY SUBCUTANEOUS SYRINGE

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	Concurrent use with another monoclonal antibody therapy (i.e., Dupixent, Cinqair, Fasenra, Nucala, Tezspire, or Xolair). Concurrent use with Janus Kinase Inhibitors (JAKis) [oral or topical].
<b>Required Medical Information</b>	Diagnosis
<b>Age Restrictions</b>	AD-12 years of age and older (initial therapy)
<b>Prescriber Restrictions</b>	Atopic Dermatitis-prescribed by or in consultation with an allergist, immunologist or dermatologist (initial therapy)
<b>Coverage Duration</b>	Initial-Atopic Dermatitis-4 months, Continuation-1 year
<b>Other Criteria</b>	Atopic Dermatitis, initial-patient has atopic dermatitis involvement estimated to be greater than or equal to 10 percent of the body surface area and patient meets a and b: a. Patient has tried at least one medium-, medium-high, high-, and/or super-high-potency prescription topical corticosteroid AND b. Inadequate efficacy was demonstrated with the previously tried topical corticosteroid therapy. Continuation- Approve if the patient has been receiving Adbry for at least 4 months and patient has responded to therapy. Note: A patient who has received less than 4 months of therapy or who is restarting therapy with Adbry should be considered under initial therapy.
<b>Indications</b>	All FDA-approved Indications.
<b>Off-Label Uses</b>	N/A
<b>Part B Prerequisite</b>	No

## ADEMPAS

---

### Products Affected

- ADEMPAS

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	Concurrent Use with Phosphodiesterase Inhibitors Used for Pulmonary Hypertension or Other Soluble Guanylate Cyclase Stimulators.
<b>Required Medical Information</b>	N/A
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	PAH and CTEPH- must be prescribed by or in consultation with a cardiologist or a pulmonologist.
<b>Coverage Duration</b>	1 year
<b>Other Criteria</b>	For PAH - must have PAH (WHO Group 1) and had a right heart catheterization to confirm the diagnosis of PAH (WHO Group 1).
<b>Indications</b>	All FDA-approved Indications.
<b>Off-Label Uses</b>	N/A
<b>Part B Prerequisite</b>	No

# AIMOVIG

## Products Affected

- AIMOVIG AUTOINJECTOR

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	Combination therapy with another cGRP inhibitor for migraine headache prevention
<b>Required Medical Information</b>	Diagnosis, number of migraine headaches per month
<b>Age Restrictions</b>	18 years and older
<b>Prescriber Restrictions</b>	N/A
<b>Coverage Duration</b>	1 year
<b>Other Criteria</b>	Approve if the patient meets the following criteria (A and B): A) Patient has greater than or equal to 4 migraine headache days per month (prior to initiating a migraine-preventative medication), AND B) If pt is currently taking Aimovig, the pt has had significant clinical benefit from the medication.
<b>Indications</b>	All FDA-approved Indications.
<b>Off-Label Uses</b>	N/A
<b>Part B Prerequisite</b>	No



# AKEEGA

## Products Affected

- AKEEGA

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	Diagnosis
<b>Age Restrictions</b>	18 years and older
<b>Prescriber Restrictions</b>	N/A
<b>Coverage Duration</b>	1 year
<b>Other Criteria</b>	Prostate cancer- Approve if the patient meets the following (A, B, C, and D): A)Patient has metastatic castration-resistant prostate cancer, AND B)Patient has a BREast CAncer (BRCA) mutation, AND C)The medication is used in combination with prednisone, AND D)Patient meets one of the following (i or ii): i. The medication is used concurrently with a gonadotropin-releasing hormone (GnRH) analog, Note: Examples are leuprolide acetate, Lupron Depot (leuprolide acetate intramuscular injection), Trelstar (triptorelin pamoate intramuscular injection), Zoladex (goserelin acetate subcutaneous implant), Vantas (histrelin acetate subcutaneous implant), Firmagon (degarelix acetate subcutaneous injection), and Orgovyx (relugolix tablets).OR ii. Patient has had a bilateral orchiectomy.
<b>Indications</b>	All FDA-approved Indications.
<b>Off-Label Uses</b>	N/A
<b>Part B Prerequisite</b>	No

# ALECENSA

## Products Affected

- ALECENSA

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	N/A
<b>Age Restrictions</b>	18 years and older
<b>Prescriber Restrictions</b>	N/A
<b>Coverage Duration</b>	1 year
<b>Other Criteria</b>	Non-small cell lung cancer-approve if the patient has both (A and B): A) either (i or ii): i) medication is used as adjuvant treatment following tumor resection (note: for tumors greater than or equal to 4 cm or node positive) or ii) advanced or metastatic disease and B) anaplastic lymphoma kinase (ALK)-positive disease as detected by an approved test. Anaplastic large cell lymphoma-approve if the patient has anaplastic lymphoma kinase (ALK)-positive disease and (i or ii): (i) the medication is used for palliative-intent therapy, or (ii) pt has relapsed or refractory disease. Erdheim-Chester disease-approve if the patient has anaplastic lymphoma kinase (ALK) rearrangement/fusion-positive disease. Inflammatory Myofibroblastic Tumor- pt has anaplastic lymphoma kinase (ALK)-positive disease AND (i or ii): (i) pt has advanced, recurrent or metastatic disease, or (ii) tumor is inoperable. Large B-Cell Lymphoma- pt has ALK-positive disease AND pt has relapsed or refractory disease.
<b>Indications</b>	All FDA-approved Indications, Some Medically-accepted Indications.
<b>Off-Label Uses</b>	Anaplastic large cell lymphoma, Erdheim Chester disease, Inflammatory Myofibroblastic Tumor, Large B-Cell Lymphoma

PA Criteria	Criteria Details
Part B Prerequisite	No

# ALOSETRON

## Products Affected

- *alosetron*

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	Diagnosis
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	12 months
Other Criteria	N/A
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

# ALPHA 1 PROTEINASE INHIBITORS

## Products Affected

- PROLASTIN-C INTRAVENOUS SOLUTION

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	Diagnosis
Age Restrictions	18 years and older
Prescriber Restrictions	N/A
Coverage Duration	1 year
Other Criteria	Alpha1-Antitrypsin Deficiency with Emphysema (or Chronic Obstructive Pulmonary Disease)-approve if the patient has a baseline (pretreatment) AAT serum concentration of less than 80 mg/dL or 11 micromol/L.
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

# ALUNBRIG

## Products Affected

- ALUNBRIG ORAL TABLET 180 MG, 30 MG, 90 MG
- ALUNBRIG ORAL TABLETS,DOSE PACK

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	ALK status
<b>Age Restrictions</b>	18 years and older
<b>Prescriber Restrictions</b>	N/A
<b>Coverage Duration</b>	1 year
<b>Other Criteria</b>	Erdheim-Chester disease-approve if the patient has anaplastic lymphoma kinase (ALK) rearrangement/fusion-positive disease. Inflammatory myofibroblastic tumor (IMT)-approve if the patient has ALK positive disease and has advanced, recurrent or metastatic disease or the tumor is inoperable. NSCLC, must be ALK-positive, as detected by an approved test, have advanced or metastatic disease and patients new to therapy must have a trial of Alecensa prior to approval of Alunbrig.
<b>Indications</b>	All FDA-approved Indications, Some Medically-accepted Indications.
<b>Off-Label Uses</b>	Erdheim-Chester disease, Inflammatory myofibroblastic tumor (IMT)
<b>Part B Prerequisite</b>	No

## ANTIBIOTICS (IV)

### Products Affected

- *amikacin injection solution 500 mg/2 ml*
- *ampicillin sodium injection recon soln 1 gram, 10 gram, 125 mg*
- *ampicillin-sulbactam injection*
- *azithromycin intravenous*
- *aztreonam*
- **BICILLIN L-A**
- *cefoxitin*
- *ceftazidime*
- *cefuroxime sodium injection recon soln 750 mg*
- *cefuroxime sodium intravenous recon soln 1.5 gram*
- *ciprofloxacin in 5 % dextrose intravenous piggyback 200 mg/100 ml*
- *clindamycin in 5 % dextrose*
- *clindamycin phosphate injection*
- *colistin (colistimethate na)*
- *doxy-100*
- *ertapenem*
- *gentamicin in nacl (iso-osm) intravenous piggyback 100 mg/100 ml, 60 mg/50 ml, 80 mg/100 ml, 80 mg/50 ml*
- *gentamicin injection solution 40 mg/ml*
- *imipenem-cilastatin*
- *levofloxacin in d5w intravenous piggyback 500 mg/100 ml, 750 mg/150 ml*
- *linezolid in dextrose 5%*
- *meropenem intravenous recon soln 1 gram, 500 mg*
- *metronidazole in nacl (iso-os)*
- *moxifloxacin-sod.chloride(iso)*
- *nafcillin injection*
- *oxacillin*
- *oxacillin in dextrose(iso-osm)*
- **PENICILLIN G POT IN DEXTROSE INTRAVENOUS PIGGYBACK 2 MILLION UNIT/50 ML, 3 MILLION UNIT/50 ML**
- *penicillin g potassium injection recon soln 20 million unit*
- *penicillin g sodium*
- **STREPTOMYCIN**
- *tazicef injection*
- **TEFLARO**
- *tigecycline*
- *tobramycin sulfate injection solution*
- *vancomycin intravenous recon soln 1,000 mg, 10 gram, 500 mg, 750 mg*

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	Diagnosis
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	N/A

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Coverage Duration</b>	3 months
<b>Other Criteria</b>	N/A
<b>Indications</b>	All FDA-approved Indications.
<b>Off-Label Uses</b>	N/A
<b>Part B Prerequisite</b>	No



## ANTIFUNGALS (IV)

### Products Affected

- *fluconazole in nacl (iso-osm) intravenous piggyback 200 mg/100 ml, 400 mg/200 ml*
- *voriconazole*

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	Diagnosis
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	3 months
Other Criteria	N/A
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

# ARCALYST

---

## Products Affected

- ARCALYST

PA Criteria	Criteria Details
Exclusion Criteria	PENDING CMS REVIEW
Required Medical Information	PENDING CMS REVIEW
Age Restrictions	PENDING CMS REVIEW
Prescriber Restrictions	PENDING CMS REVIEW
Coverage Duration	PENDING CMS REVIEW
Other Criteria	PENDING CMS REVIEW
Indications	PENDING CMS REVIEW
Off-Label Uses	PENDING CMS REVIEW
Part B Prerequisite	PENDING CMS REVIEW

# ARIKAYCE

---

## Products Affected

- ARIKAYCE

PA Criteria	Criteria Details
Exclusion Criteria	PENDING CMS REVIEW
Required Medical Information	PENDING CMS REVIEW
Age Restrictions	PENDING CMS REVIEW
Prescriber Restrictions	PENDING CMS REVIEW
Coverage Duration	PENDING CMS REVIEW
Other Criteria	PENDING CMS REVIEW
Indications	PENDING CMS REVIEW
Off-Label Uses	PENDING CMS REVIEW
Part B Prerequisite	PENDING CMS REVIEW

# AUBAGIO

## Products Affected

- *teriflunomide*

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	Concurrent use of teriflunomide with other disease-modifying agents used for multiple sclerosis (MS)
<b>Required Medical Information</b>	Relapsing form of MS, to include, clinically-isolated syndrome, relapsing-remitting disease, and active secondary progressive disease.
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	Prescribed by or in consultation with a neurologist or MS specialist.
<b>Coverage Duration</b>	Authorization will be for 1 year.
<b>Other Criteria</b>	N/A
<b>Indications</b>	All FDA-approved Indications.
<b>Off-Label Uses</b>	N/A
<b>Part B Prerequisite</b>	No

# AUGTYRO

---

## Products Affected

- AUGTYRO

PA Criteria	Criteria Details
Exclusion Criteria	PENDING CMS REVIEW
Required Medical Information	PENDING CMS REVIEW
Age Restrictions	PENDING CMS REVIEW
Prescriber Restrictions	PENDING CMS REVIEW
Coverage Duration	PENDING CMS REVIEW
Other Criteria	PENDING CMS REVIEW
Indications	PENDING CMS REVIEW
Off-Label Uses	PENDING CMS REVIEW
Part B Prerequisite	PENDING CMS REVIEW

# AVONEX

## Products Affected

- AVONEX INTRAMUSCULAR PEN INJECTOR KIT
- AVONEX INTRAMUSCULAR SYRINGE KIT

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	Concurrent use of other disease-modifying agent used for multiple sclerosis
<b>Required Medical Information</b>	Relapsing form of Multiple Sclerosis (MS), to include clinically-isolated syndrome, relapsing-remitting disease, and active secondary progressive disease
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	Prescribed by or after consultation with a neurologist or an MS specialist.
<b>Coverage Duration</b>	Authorization will be for 1 year
<b>Other Criteria</b>	N/A
<b>Indications</b>	All FDA-approved Indications.
<b>Off-Label Uses</b>	N/A
<b>Part B Prerequisite</b>	No

# AYVAKIT

## Products Affected

- AYVAKIT

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	Diagnosis
<b>Age Restrictions</b>	18 years and older
<b>Prescriber Restrictions</b>	N/A
<b>Coverage Duration</b>	1 year
<b>Other Criteria</b>	GIST-approve if the tumor is positive for platelet-derived growth factor receptor alpha (PDGFRA) exon 18 mutation or if the patient has tried two of the following: Gleevec (imatinib), Sutent (sunitinib), Sprycel (dasatinib), Stivarga (regorafenib) or Qinlock (ripretinib). Myeloid/Lymphoid Neoplasms with eosinophilia-approve if the tumor is positive for PDGFRA D842V mutation. Systemic mastocytosis-Approve if the patient has a platelet count greater than or equal to 50,000/mcL and patient has either indolent systemic mastocytosis or one of the following subtypes of advanced systemic mastocytosis-aggressive systemic mastocytosis, systemic mastocytosis with an associated hematological neoplasm or mast cell leukemia.
<b>Indications</b>	All FDA-approved Indications, Some Medically-accepted Indications.
<b>Off-Label Uses</b>	Myeloid/Lymphoid neoplasms with Eosinophilia
<b>Part B Prerequisite</b>	No

# BALVERSA

## Products Affected

- BALVERSA

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	Diagnosis, previous therapies, test results
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	1 year
Other Criteria	Urothelial Carcinoma, locally advanced or metastatic-approve if the patient has susceptible fibroblast growth factor receptor 3 genetic alterations AND the patient has progressed during or following prior platinum-containing chemotherapy, other chemotherapy or checkpoint inhibitor therapy.
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Part B Prerequisite	No



# BENLYSTA

---

## Products Affected

- BENLYSTA SUBCUTANEOUS

PA Criteria	Criteria Details
Exclusion Criteria	PENDING CMS REVIEW
Required Medical Information	PENDING CMS REVIEW
Age Restrictions	PENDING CMS REVIEW
Prescriber Restrictions	PENDING CMS REVIEW
Coverage Duration	PENDING CMS REVIEW
Other Criteria	PENDING CMS REVIEW
Indications	PENDING CMS REVIEW
Off-Label Uses	PENDING CMS REVIEW
Part B Prerequisite	PENDING CMS REVIEW

# BESREMI

---

## Products Affected

- BESREMI

PA Criteria	Criteria Details
Exclusion Criteria	Concomitant use with other interferon products
Required Medical Information	Diagnosis
Age Restrictions	18 years and older
Prescriber Restrictions	Prescribed by or in consultation with an oncologist
Coverage Duration	1 year
Other Criteria	N/A
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

## BETASERON/EXTAVIA

### Products Affected

- BETASERON SUBCUTANEOUS KIT

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	Concurrent use with other disease-modifying agent used for multiple sclerosis
<b>Required Medical Information</b>	Relapsing form of Multiple Sclerosis (MS), to include clinically-isolated syndrome, relapsing-remitting disease, and active secondary progressive disease
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	Prescribed by or after consultation with a neurologist or an MS specialist.
<b>Coverage Duration</b>	Authorization will be for 1 year
<b>Other Criteria</b>	N/A
<b>Indications</b>	All FDA-approved Indications.
<b>Off-Label Uses</b>	N/A
<b>Part B Prerequisite</b>	No

## BEXAROTENE (ORAL)

---

### Products Affected

- bexarotene*

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	Diagnosis
Age Restrictions	N/A
Prescriber Restrictions	Prescribed by or in consultation with an oncologist or dermatologist (initial and continuation)
Coverage Duration	1 year
Other Criteria	N/A
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

## BEXAROTENE (TOPICAL)

### Products Affected

- bexarotene*

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	Diagnosis
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	Prescribed by or in consultation with an oncologist or dermatologist (initial and continuation)
<b>Coverage Duration</b>	1 year
<b>Other Criteria</b>	Adult T-Cell Leukemia/Lymphoma- approve if the patient has chronic/smoldering subtype and this medication is used as first-line therapy.
<b>Indications</b>	All FDA-approved Indications, Some Medically-accepted Indications.
<b>Off-Label Uses</b>	Adult T-Cell Leukemia/Lymphoma
<b>Part B Prerequisite</b>	No

## BOSENTAN/AMBRISENTAN

### Products Affected

- *ambrisentan*
- *bosentan*

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	Pulmonary arterial hypertension (PAH) WHO Group 1, results of right heart cath
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	For treatment of pulmonary arterial hypertension, ambrisentan or bosentan must be prescribed by or in consultation with a cardiologist or a pulmonologist. CTEPH - prescribed by or in consultation with a cardiologist or pulmonologist
<b>Coverage Duration</b>	Authorization will be for 1 year.
<b>Other Criteria</b>	CTEPH - pt must have tried Adempas, has a contraindication to Adempas, or is currently receiving bosentan for CTEPH. Pulmonary arterial hypertension (PAH) WHO Group 1, are required to have had a right-heart catheterization to confirm diagnosis of PAH to ensure appropriate medical assessment.
<b>Indications</b>	All FDA-approved Indications, Some Medically-accepted Indications.
<b>Off-Label Uses</b>	Chronic thromboembolic pulmonary hypertension (CTEPH) (bosentan)
<b>Part B Prerequisite</b>	No

# BOSULIF

## Products Affected

- BOSULIF ORAL CAPSULE 100 MG, 50 MG
- BOSULIF ORAL TABLET 100 MG, 400 MG, 500 MG

PA Criteria	Criteria Details
Exclusion Criteria	PENDING CMS REVIEW
Required Medical Information	PENDING CMS REVIEW
Age Restrictions	PENDING CMS REVIEW
Prescriber Restrictions	PENDING CMS REVIEW
Coverage Duration	PENDING CMS REVIEW
Other Criteria	PENDING CMS REVIEW
Indications	PENDING CMS REVIEW
Off-Label Uses	PENDING CMS REVIEW
Part B Prerequisite	PENDING CMS REVIEW

# BRAFTOVI

## Products Affected

- BRAFTOVI

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	Diagnosis, BRAF V600 status
<b>Age Restrictions</b>	18 years and older
<b>Prescriber Restrictions</b>	N/A
<b>Coverage Duration</b>	1 year
<b>Other Criteria</b>	Melanoma - approve if the patient has unresectable, advanced or metastatic melanoma AND has a BRAF V600 mutation. In addition, patients new to therapy must have a trial of Zelboraf or Tafinlar prior to approval of Braftovi. Colon or Rectal cancer-approve if the patient meets the following (A, B, and C): A) The patient has BRAF V600E mutation-positive disease AND B) The patient has previously received a chemotherapy regimen for colon or rectal cancer AND C) The agent is prescribed as part of a combination regimen for colon or rectal cancer. NSCLC- approve if pt has BRAF V600E mutation-positive metastatic disease AND this medication will be taken in combination with Mektovi (binimetinib tablets).
<b>Indications</b>	All FDA-approved Indications.
<b>Off-Label Uses</b>	N/A
<b>Part B Prerequisite</b>	No



# BRUKINSA

## Products Affected

- BRUKINSA

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	Diagnosis, prior therapies
<b>Age Restrictions</b>	18 years and older
<b>Prescriber Restrictions</b>	N/A
<b>Coverage Duration</b>	1 year
<b>Other Criteria</b>	Follicular Lymphoma - approve if pt tried at least two other systemic regimens and will use this in combination with Gazyva (obinutuzumab intravenous infusion). Mantle Cell Lymphoma/CLL/SLL - for patients new to therapy, approve if the patient has tried Calquence. Marginal zone lymphoma - approve if the patient has tried at least one systemic regimen. Waldenstrom macroglobulinemia/lymphoplasmacytic lymphoma - approve. Hairy Cell Leukemia - approve if pt has received therapy for relapsed or refractory disease AND pt has progressive disease.
<b>Indications</b>	All FDA-approved Indications, Some Medically-accepted Indications.
<b>Off-Label Uses</b>	Hairy Cell Leukemia
<b>Part B Prerequisite</b>	No

## C1 ESTERASE INHIBITORS

### Products Affected

- CINRYZE

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	N/A
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	prescribed by or in consultation with an allergist/immunologist or a physician that specializes in the treatment of HAE or related disorders
<b>Coverage Duration</b>	1 year
<b>Other Criteria</b>	Hereditary Angioedema (HAE) Due to C1 Inhibitor (C1-INH) Deficiency [Type I or Type II], Prophylaxis, Initial Therapy: approve if the patient has HAE type I or type II confirmed by low levels of functional C1-INH protein (less than 50 percent of normal) at baseline and lower than normal serum C4 levels at baseline. Patient is currently taking Cinryze for prophylaxis - approve if the patient meets the following criteria (i and ii): i) patient has a diagnosis of HAE type I or II, and ii) according to the prescriber, the patient has had a favorable clinical response since initiating Cinryze as prophylactic therapy compared with baseline.
<b>Indications</b>	All FDA-approved Indications.
<b>Off-Label Uses</b>	N/A
<b>Part B Prerequisite</b>	No

# CABLIVI

## Products Affected

- CABLIVI INJECTION KIT

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	Diagnosis, concurrent medications
Age Restrictions	18 years and older
Prescriber Restrictions	Prescribed by or in consultation with a hematologist
Coverage Duration	Approve for 12 months
Other Criteria	aTTP-approve if the requested medication was initiated in the inpatient setting in combination with plasma exchange therapy AND patient is currently receiving at least one immunosuppressive therapy AND if the patient has previously received Cablivi, he/she has not had more than two recurrences of aTTP while on Cablivi.
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

# CABOMETYX

## Products Affected

- CABOMETYX

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	Diagnosis, histology, RET gene rearrangement status for NSCLC
<b>Age Restrictions</b>	Thyroid carcinoma-12 years and older, other dx (except bone cancer)-18 years and older
<b>Prescriber Restrictions</b>	N/A
<b>Coverage Duration</b>	1 year
<b>Other Criteria</b>	Renal Cell Carcinoma-Approve if the patient has relapsed or stage IV disease. Hepatocellular Carcinoma-approve if the patient has been previously treated with at least one other systemic therapy (e.g., Nexavar, Lenvima). Bone cancer-approve if the patient has Ewing sarcoma or osteosarcoma and has tried at least one previous systemic regimen. Thyroid carcinoma-approve if the patient has differentiated thyroid carcinoma, patient is refractory to radioactive iodine therapy and the patient has tried Lenvima or sorafenib. Endometrial carcinoma-approve if the patient has tried one systemic regimen. GIST-approve if the patient has tried two of the following-imatinib, Ayvakit, sunitinib, dasatinib, Stivarga or Qinlock. NSCLC-approve if the patient has RET rearrangement psotivie tumor.
<b>Indications</b>	All FDA-approved Indications, Some Medically-accepted Indications.
<b>Off-Label Uses</b>	Patients with Non-Small Cell Lung Cancer, Gastrointestinal stromal tumors (GIST), Bone cancer, Endometrial Carcinoma
<b>Part B Prerequisite</b>	No

# CALQUENCE

## Products Affected

- CALQUENCE
- CALQUENCE (ACALABRUTINIB MAL)

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	N/A
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	N/A
<b>Coverage Duration</b>	1 year
<b>Other Criteria</b>	CLL and SLL-approve. Mantle Cell Lymphoma- approve if the patient has tried at least one systemic regimen or is not a candidate for a systemic regimen (e.g., rituximab, dexamethasone, cytarabine, carboplatin, cisplatin, oxaliplatin, cyclophosphamide, doxorubicin, vincristine, prednisone, methotrexate, bendamustine, bortezomib, or lenalidomide). Marginal Zone Lymphoma-approve if patient has tried at least one systemic regimen (e.g., bendamustine, rituximab, cyclophosphamide, doxorubicin, vincristine, prednisone, lenalidomide, or chlorambucil). Waldenstrom Macroglobulinemia/Lymphoplasmacytic Lymphoma-approve if the patient has tried at least one systemic regimen (e.g., Brukinsa [zanubrutinib capsules], Imbruvica [ibrutinib tablets and capsules], rituximab, bendamustine, cyclophosphamide, dexamethasone, bortezomib, fludarabine, or cladribine)
<b>Indications</b>	All FDA-approved Indications, Some Medically-accepted Indications.
<b>Off-Label Uses</b>	Waldenstrom's Macroglobulinemia/Lymphoplasmacytic Lymphoma, Marginal zone lymphoma.

PA Criteria	Criteria Details
Part B Prerequisite	No

# CAPRELSA

## Products Affected

- CAPRELSA ORAL TABLET 100 MG, 300 MG

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	N/A
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	N/A
<b>Coverage Duration</b>	1 year
<b>Other Criteria</b>	MTC - approve. DTC - approve if refractory to radioactive iodine therapy.
<b>Indications</b>	All FDA-approved Indications, Some Medically-accepted Indications.
<b>Off-Label Uses</b>	Differentiated (i.e., papillary, follicular, and oncocytic) Thyroid Carcinoma.
<b>Part B Prerequisite</b>	No

## CARGLUMIC ACID

### Products Affected

- carglumic acid*

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	Diagnosis
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	Prescribed by or in consultation with a metabolic disease specialist or a specialist who focuses in the treatment of metabolic diseases
<b>Coverage Duration</b>	NAGS-Pt meets criteria no genetic test - 3mo. Pt had genetic test - 12mo, other-approve 7 days
<b>Other Criteria</b>	N-Acetylglutamate synthase deficiency with hyperammonemia-Approve if genetic testing confirmed a mutation leading to N-acetylglutamate synthase deficiency or if the patient has hyperammonemia. Propionic Acidemia or Methylmalonic Acidemia with Hyperammonemia, Acute Treatment-approve if the patient's plasma ammonia level is greater then or equal to 50 micromol/L and the requested medication will be used in conjunction with other ammonia-lowering therapies.
<b>Indications</b>	All FDA-approved Indications, Some Medically-accepted Indications.
<b>Off-Label Uses</b>	Acute hyperammonemia due to propionic acidemia (PA) or methylmalonic acidemia (MMA) (generic carglumic acid)
<b>Part B Prerequisite</b>	No



# CAYSTON

---

## Products Affected

- CAYSTON

PA Criteria	Criteria Details
Exclusion Criteria	PENDING CMS REVIEW
Required Medical Information	PENDING CMS REVIEW
Age Restrictions	PENDING CMS REVIEW
Prescriber Restrictions	PENDING CMS REVIEW
Coverage Duration	PENDING CMS REVIEW
Other Criteria	PENDING CMS REVIEW
Indications	PENDING CMS REVIEW
Off-Label Uses	PENDING CMS REVIEW
Part B Prerequisite	PENDING CMS REVIEW

# CHEMET

## Products Affected

- CHEMET

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	Blood lead level
<b>Age Restrictions</b>	Approve in patients between the age of 12 months and 18 years
<b>Prescriber Restrictions</b>	Prescribed by or in consultation with a professional experienced in the use of chelation therapy (eg, a medical toxicologist or a poison control center specialist)
<b>Coverage Duration</b>	Approve for 2 months
<b>Other Criteria</b>	Approve if Chemet is being used to treat acute lead poisoning (not as prophylaxis) and prior to starting Chemet therapy the patient's blood lead level was greater than 45 mcg/dL.
<b>Indications</b>	All FDA-approved Indications.
<b>Off-Label Uses</b>	N/A
<b>Part B Prerequisite</b>	No

# CIBINQO

## Products Affected

- CIBINQO

PA Criteria	Criteria Details
Exclusion Criteria	PENDING CMS REVIEW
Required Medical Information	PENDING CMS REVIEW
Age Restrictions	PENDING CMS REVIEW
Prescriber Restrictions	PENDING CMS REVIEW
Coverage Duration	PENDING CMS REVIEW
Other Criteria	PENDING CMS REVIEW
Indications	PENDING CMS REVIEW
Off-Label Uses	PENDING CMS REVIEW
Part B Prerequisite	PENDING CMS REVIEW

# CIMZIA

## Products Affected

- CIMZIA POWDER FOR RECONST
- CIMZIA SUBCUTANEOUS SYRINGE KIT 400 MG/2 ML (200 MG/ML X 2)

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	Concurrent use with a Biologic DMARD or Targeted Synthetic DMARD
<b>Required Medical Information</b>	Diagnosis, concurrent medications, previous therapies tried
<b>Age Restrictions</b>	18 years and older for CD and PP (initial therapy).
<b>Prescriber Restrictions</b>	All dx initial therapy only. RA/AS, prescribed by or in consultation with a rheumatologist. Crohn's disease, prescribed by or in consultation with a gastroenterologist. PsA prescribed by or in consultation with a rheumatologist or dermatologist. PP, prescribed by or in consultation with a dermatologist. nr-axSpA-prescribed by or in consultation with a rheumatologist
<b>Coverage Duration</b>	Approve through end of plan year
<b>Other Criteria</b>	AS initial tx, approve if the patient has tried TWO of the following drugs in the past: Enbrel, a preferred adalimumab product, Xeljanz/XR, Cosentyx. Note: if the patient does not meet this requirement, a previous trial of another non-preferred adalimumab product will also count. PsA initial tx, approve if the patient has tried TWO of the following drugs in the past: Enbrel, a preferred adalimumab product, Cosentyx, Tremfya, Stelara, Otezla, Orencia, Rinvoq, Skyrizi or Xeljanz/XR. Note: if the patient does not meet this requirement, a previous trial of another non-preferred adalimumab product will also count. RA initial tx, approve if the patient has tried two of the following drugs in the past: Enbrel, a preferred adalimumab product, Orencia, Rinvoq or Xeljanz/XR. Note: if the patient does not meet this requirement, a previous trial of another non-preferred adalimumab product will also count. CD initial tx, approve if patient has previously tried TWO of the following drugs in the past: a preferred adalimumab product, a preferred infliximab product, Stelara, Skyrizi or

PA Criteria	Criteria Details
	<p>Rinvoq. Note: if the patient does not meet this requirement, a previous trial of another non-preferred adalimumab/infliximab product will also count. Plaque Psoriasis (PP), initial tx-approve if the patient has tried TWO of the following drugs in the past: Enbrel, a preferred adalimumab product, Skyrizi, Stelara SC, Otezla, Cosentyx, Tremfya, Sotyktu. A trial of a non-preferred adalimumab also counts. Cont tx, AS/PsA/RA/CD/PP - approve if the pt had a response as determined by the prescriber. Non-radiographic axial spondylitis (nr-axSpA), initial tx-approve if the patient has objective signs of inflammation, defined as at least one of the following: C-reactive protein (CRP) elevated beyond the upper limit of normal for the reporting laboratory OR sacroilitis reported on MRI. nr-axSpA continuation tx-approve if the patient has had a response as determined by the prescriber. Please Note: preferred adalimumab products include Humira (NDCs starting with -00074), Cyltezo, Yuflyma. Preferred infliximab products include Remicade, Zymfentra.</p>
<b>Indications</b>	All FDA-approved Indications.
<b>Off-Label Uses</b>	N/A
<b>Part B Prerequisite</b>	No

# CINACALCET

## Products Affected

- *cinacalcet*

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	Diagnosis
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	Hypercalcemia d/t parathyroid CA-prescr/consult w/onco or endo. Hypercalcemia w/primary hyperparathyroidism-prescr/consult w/nephro or endo. Hyperparathyroidism in post-renal transplant-prescr/consult w/transplant physician/nephro/endo.
<b>Coverage Duration</b>	12 months
<b>Other Criteria</b>	Hypercalcemia due to parathyroid carcinoma-approve. Hypercalcemia in patients with primary hyperparathyroidism-approve if the patient has failed or is unable to undergo a parathyroidectomy due to a contraindication. Secondary Hyperparathyroidism in patients with chronic kidney disease on dialysis - deny under Medicare Part D (claim should be submitted under the ESRD bundles payment benefit). Hyperparathyroidism in Post-Renal Transplant Patients-approve if the baseline (prior to starting cinacalcet therapy) calcium and intact parathyroid hormone (iPTH) levels are above the normal range, as defined by the laboratory reference values.
<b>Indications</b>	All FDA-approved Indications, Some Medically-accepted Indications.
<b>Off-Label Uses</b>	hyperparathyroidism in post-renal transplant patients
<b>Part B Prerequisite</b>	No

# CLOBAZAM

## Products Affected

- *clobazam oral suspension*
- *clobazam oral tablet*
- SYMPAZAN

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	Diagnosis, other medications tried
<b>Age Restrictions</b>	2 years and older (initial therapy)
<b>Prescriber Restrictions</b>	Prescribed by or in consultation with a neurologist (initial therapy)
<b>Coverage Duration</b>	1 year
<b>Other Criteria</b>	Lennox-Gastaut Syndrome, initial therapy-patient has tried and/or is concomitantly receiving one of the following: lamotrigine, topiramate, rufinamide, felbamate, Fintepla, Epidiolex or valproic acid. Treatment refractory seizures/epilepsy, initial therapy-patient has tried and/or is concomitantly receiving at least two other antiepileptic drugs (e.g., valproic acid, lamotrigine, topiramate, clonazepam, levetiracetam, zonisamide, felbamate). Continuation-prescriber confirms patient is responding to therapy.
<b>Indications</b>	All FDA-approved Indications, Some Medically-accepted Indications.
<b>Off-Label Uses</b>	Dravet Syndrome and treatment-refractory seizures/epilepsy
<b>Part B Prerequisite</b>	No

# COMETRIQ

## Products Affected

- COMETRIQ ORAL CAPSULE 100 MG/DAY(80 MG X1-20 MG X3), 60 MG/DAY(80 MG X1-20 MG X1), 140 MG/DAY (20 MG X 3/DAY)

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	Diagnosis.
<b>Age Restrictions</b>	18 years and older
<b>Prescriber Restrictions</b>	N/A
<b>Coverage Duration</b>	1 year
<b>Other Criteria</b>	MTC - approve. Non-Small Cell Lung Cancer with RET Gene Rearrangements - approve. Differentiated (i.e., papillary, follicular, and oncocytic) Thyroid Carcinoma-approve if the patient's carcinoma is refractory to radioactive iodine therapy and patient has tried a Vascular Endothelial Growth Factor Receptor (VEGFR)-targeted therapy.
<b>Indications</b>	All FDA-approved Indications, Some Medically-accepted Indications.
<b>Off-Label Uses</b>	Non-Small Cell Lung Cancer with RET Gene Rearrangements, Differentiated (i.e., papillary, follicular, and oncocytic) Thyroid Carcinoma
<b>Part B Prerequisite</b>	No



# COPIKTRA

## Products Affected

- COPIKTRA

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	Diagnosis, previous therapies
<b>Age Restrictions</b>	18 years and older
<b>Prescriber Restrictions</b>	N/A
<b>Coverage Duration</b>	1 year
<b>Other Criteria</b>	Chronic Lymphocytic Leukemia/ Small Lymphocytic Lymphoma - approve if the patient has tried one systemic regimen (e.g., Imbruvica (ibrutinib capsules, tablets and oral solution), Venclexta (venetoclax tablets), rituximab, Gazyva (obinutuzumab intravenous infusion), chlorambucil, fludarabine, cyclophosphamide, bendamustine, high-dose methylprednisolone, Campath (alemtuzumab intravenous infusion), Calquence (acalabrutinib capsules), Brukinsa (zanubrutinib capsules), or Arzerra (ofatumumab intravenous infusion). T-cell lymphoma- For peripheral T-cell lymphoma, approve. For breast implant-associated anaplastic large cell lymphoma, or hepatosplenic T-cell lymphoma, approve if the patient has relapsed or refractory disease.
<b>Indications</b>	All FDA-approved Indications, Some Medically-accepted Indications.
<b>Off-Label Uses</b>	T-cell Lymphoma
<b>Part B Prerequisite</b>	No

# COSENTYX

## Products Affected

- COSENTYX (2 SYRINGES)
- COSENTYX PEN (2 PENS)
- COSENTYX SUBCUTANEOUS SYRINGE 75 MG/0.5 ML
- COSENTYX UNOREADY PEN

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	Concurrent Use with other Biologics or Targeted Synthetic Disease-Modifying Antirheumatic Drugs (DMARDs)
<b>Required Medical Information</b>	Diagnosis and previous medications use
<b>Age Restrictions</b>	PP-6 yr and older.AS/Spondy/HS initial - 18 years of age and older. PsA-2 years and older. Enthesitis-4 years and older
<b>Prescriber Restrictions</b>	PP initial-presc/consult derm. PsA initial - prescribed by or in consultation with a dermatologist or rheumatologist. AS/spondylo/enthesitis initial- by or in consultation with rheumatologist. HS initial - by or in consult w/ dermatologist
<b>Coverage Duration</b>	Approve through end of plan year
<b>Other Criteria</b>	INITIAL THERAPY: HIDRADENITIS SUPPURATIVA (HS): tried at least one other therapy (e.g. systemic antibiotics, isotretinoin). NON-RADIOGRAPHIC AXIAL SPONDYLOARTHRITIS: objective signs of inflammation and meets a or b: a) C-reactive protein elevated beyond the upper limit of normal or b) sacroiliitis reported on MRI. PLAQUE PSORIASIS (PP) [A or B]: A) tried at least one traditional systemic agent (e.g., methotrexate [MTX], cyclosporine, acitretin, PUVA) for at least 3 months, unless intolerant (Note: a trial of at least one biologic that is not Cosentyx or a Cosentyx biosimilar also counts) or B) contraindication to MTX. CONTINUATION THERAPY: ALL INDICATIONS: patient has experienced benefit from the medication.
<b>Indications</b>	All FDA-approved Indications.
<b>Off-Label Uses</b>	N/A



PA Criteria	Criteria Details
Part B Prerequisite	No

# COTELLIC

---

## Products Affected

- COTELLIC

PA Criteria	Criteria Details
Exclusion Criteria	PENDING CMS REVIEW
Required Medical Information	PENDING CMS REVIEW
Age Restrictions	PENDING CMS REVIEW
Prescriber Restrictions	PENDING CMS REVIEW
Coverage Duration	PENDING CMS REVIEW
Other Criteria	PENDING CMS REVIEW
Indications	PENDING CMS REVIEW
Off-Label Uses	PENDING CMS REVIEW
Part B Prerequisite	PENDING CMS REVIEW

## CRESEMBA (ORAL)

### Products Affected

- CRESEMBA ORAL

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	Diagnosis
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	3 months
Other Criteria	N/A
Indications	All FDA-approved Indications, Some Medically-accepted Indications.
Off-Label Uses	Candidiasis of the esophagus - HIV infection, sepsis
Part B Prerequisite	No

## CYSTEAMINE (OPHTHALMIC)

### Products Affected

- CYSTARAN

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	Diagnosis
Age Restrictions	N/A
Prescriber Restrictions	Prescribed by or in consultation with an ophthalmologist or a metabolic disease specialist or specialist who focuses in the treatment of metabolic diseases
Coverage Duration	1 year
Other Criteria	Approve if the patient has corneal cysteine crystal deposits confirmed by slit-lamp examination
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

## CYSTEAMINE (ORAL)

### Products Affected

- CYSTAGON

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	Concomitant use of Cystagon and Procysbi
<b>Required Medical Information</b>	Diagnosis, genetic tests and lab results (as specified in the Other Criteria field)
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	Prescribed by or in consultation with a nephrologist or a metabolic disease specialist (or specialist who focuses in the treatment of metabolic diseases)
<b>Coverage Duration</b>	1 year
<b>Other Criteria</b>	Cystinosis, nephropathic-approve if the prescriber confirms the diagnosis was confirmed by genetic testing confirming a mutation of the CTNS gene OR white blood cell cystine concentration above the upper limit of the normal reference range for the reporting laboratory.
<b>Indications</b>	All FDA-approved Indications.
<b>Off-Label Uses</b>	N/A
<b>Part B Prerequisite</b>	No

# DALFAMPRIDINE

## Products Affected

- dalfampridine*

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	N/A
<b>Age Restrictions</b>	18 years and older (initial and continuation therapy)
<b>Prescriber Restrictions</b>	MS. If prescribed by, or in consultation with, a neurologist or MS specialist (initial and continuation).
<b>Coverage Duration</b>	Initial-4months, Continuation-1 year
<b>Other Criteria</b>	Initial-approve if the patient is ambulatory, the requested medication is being used to improve or maintain mobility in a patient with MS and the patient has impaired ambulation as evaluated by an objective measure (e.g., timed 25 foot walk and multiple sclerosis walking scale-12). Continuation-approve if the patient is ambulatory, the requested medication is being used to improve or maintain mobility in a patient with MS and the patient has responded to or is benefiting from therapy.
<b>Indications</b>	All FDA-approved Indications.
<b>Off-Label Uses</b>	N/A
<b>Part B Prerequisite</b>	No



# DAURISMO

## Products Affected

- DAURISMO ORAL TABLET 100 MG, 25 MG

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	Diagnosis, medications that will be used in combination, comorbidities
<b>Age Restrictions</b>	18 years and older
<b>Prescriber Restrictions</b>	N/A
<b>Coverage Duration</b>	1 year
<b>Other Criteria</b>	AML - approve if Daurismo will be used in combination with cytarabine.
<b>Indications</b>	All FDA-approved Indications.
<b>Off-Label Uses</b>	N/A
<b>Part B Prerequisite</b>	No

# DEFERASIROX

## Products Affected

- *deferasirox*

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	Serum ferritin level
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	Prescribed by or in consultation with a hematologist
<b>Coverage Duration</b>	1 year
<b>Other Criteria</b>	Transfusion-related chronic iron overload, initial therapy - approve if the patient is receiving blood transfusions at regular intervals for various conditions (eg, thalassemia syndromes, myelodysplastic syndrome, chronic anemia, sickle cell disease) AND prior to starting therapy, the serum ferritin level is greater than 1,000 mcg/L. Non-transfusion-dependent thalassemia syndromes chronic iron overload, initial therapy - approve if prior to starting therapy the serum ferritin level is greater than 300 mcg/L. Continuation therapy - approve is the patient is benefiting from therapy as confirmed by the prescribing physician.
<b>Indications</b>	All FDA-approved Indications.
<b>Off-Label Uses</b>	N/A
<b>Part B Prerequisite</b>	No

# DEFERIPRONE

## Products Affected

- *deferiprone*

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	Serum ferritin level
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	Prescribed by or in consultation with a hematologist
<b>Coverage Duration</b>	1 year
<b>Other Criteria</b>	Iron overload, chronic-transfusion related due to thalassemia syndrome or related to sickle cell disease or other anemias Initial therapy - approve. Continuation therapy - approve is the patient is benefiting from therapy as confirmed by the prescribing physician.
<b>Indications</b>	All FDA-approved Indications.
<b>Off-Label Uses</b>	N/A
<b>Part B Prerequisite</b>	No

## DIABETIC SUPPLY - ALCOHOL PADS

---

### Products Affected

- *alcohol pads*

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	N/A
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	12 months
Other Criteria	Approve if the prescriber confirms that the medical supply is being requested for a use that is directly associated with delivering insulin to the body.
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

## DIABETIC SUPPLY - GAUZE PADS

### Products Affected

- GAUZE PADS 2 X 2

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	N/A
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	12 months
Other Criteria	Approve if the prescriber confirms that the medical supply is being requested for a use that is directly associated with delivering insulin to the body.
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

## DIABETIC SUPPLY - NEEDLES

### Products Affected

- INSULIN SYRINGES (NON-PREFERRED BRANDS) SYRINGE 1 ML 29 GAUGE X 1/2"
- PEN NEEDLES (NON-PREFERRED BRANDS) NEEDLE 29 GAUGE X 1/2"

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	N/A
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	12 months
Other Criteria	Approve if the prescriber confirms that the medical supply is being requested for a use that is directly associated with delivering insulin to the body.
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

## DIABETIC SUPPLY - SYRINGES

### Products Affected

- INSULIN SYRINGE-NEEDLE U-100  
SYRINGE 0.3 ML 29 GAUGE, 1 ML 29  
GAUGE X 1/2", 1/2 ML 28 GAUGE

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	N/A
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	12 months
Other Criteria	Approve if the prescriber confirms that the medical supply is being requested for a use that is directly associated with delivering insulin to the body.
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

# DIACOMIT

## Products Affected

- DIACOMIT

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	Documentation of diagnosis.
<b>Age Restrictions</b>	6 months and older (initial therapy)
<b>Prescriber Restrictions</b>	Prescribed by or in consultation with an neurologist (initial therapy)
<b>Coverage Duration</b>	1 year
<b>Other Criteria</b>	Dravet Syndrome-Initial therapy-approve if the patient is concomitantly receiving clobazam or is unable to take clobazam due to adverse events. Dravet Syndrome-Continuation-approve if the patient is responding to therapy.
<b>Indications</b>	All FDA-approved Indications.
<b>Off-Label Uses</b>	N/A
<b>Part B Prerequisite</b>	No



# DIMETHYL FUMARATE

## Products Affected

- *dimethyl fumarate oral capsule, delayed release(dr/ec) 120 mg, 120 mg (14)- 240 mg (46), 240 mg*

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	Concurrent use with other disease-modifying agents used for multiple sclerosis (MS).
<b>Required Medical Information</b>	Relapsing form of Multiple Sclerosis (MS), to include clinically-isolated syndrome, relapsing-remitting disease, and active secondary progressive disease
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	Prescribed by or in consultation with a neurologist or MS specialist.
<b>Coverage Duration</b>	Authorization will be for 1 year.
<b>Other Criteria</b>	N/A
<b>Indications</b>	All FDA-approved Indications.
<b>Off-Label Uses</b>	N/A
<b>Part B Prerequisite</b>	No

# DOPTELET

## Products Affected

- DOPTELET (10 TAB PACK)
- DOPTELET (15 TAB PACK)
- DOPTELET (30 TAB PACK)

PA Criteria	Criteria Details
Exclusion Criteria	PENDING CMS REVIEW
Required Medical Information	PENDING CMS REVIEW
Age Restrictions	PENDING CMS REVIEW
Prescriber Restrictions	PENDING CMS REVIEW
Coverage Duration	PENDING CMS REVIEW
Other Criteria	PENDING CMS REVIEW
Indications	PENDING CMS REVIEW
Off-Label Uses	PENDING CMS REVIEW
Part B Prerequisite	PENDING CMS REVIEW

# DROXIDOPA

## Products Affected

- *droxidopa*

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	Medication history (as described in Other Criteria field)
<b>Age Restrictions</b>	18 years of age and older
<b>Prescriber Restrictions</b>	Prescribed by or in consultation with a cardiologist or a neurologist
<b>Coverage Duration</b>	12 months
<b>Other Criteria</b>	NOH, approve if the patient meets ALL of the following criteria: a) Patient has been diagnosed with symptomatic NOH due to primary autonomic failure (Parkinson's disease, multiple system atrophy, pure autonomic failure), dopamine beta-hydroxylase deficiency, or non-diabetic autonomic neuropathy, AND b) Patient has tried midodrine
<b>Indications</b>	All FDA-approved Indications.
<b>Off-Label Uses</b>	N/A
<b>Part B Prerequisite</b>	No

# DUAL OREXIN RECEPTOR ANTAGONIST

---

## Products Affected

- BELSOMRA

PA Criteria	Criteria Details
Exclusion Criteria	PENDING CMS REVIEW
Required Medical Information	PENDING CMS REVIEW
Age Restrictions	PENDING CMS REVIEW
Prescriber Restrictions	PENDING CMS REVIEW
Coverage Duration	PENDING CMS REVIEW
Other Criteria	PENDING CMS REVIEW
Indications	PENDING CMS REVIEW
Off-Label Uses	PENDING CMS REVIEW
Part B Prerequisite	PENDING CMS REVIEW

# DUPIXENT

## Products Affected

- DUPIXENT PEN SUBCUTANEOUS PEN INJECTOR 200 MG/1.14 ML, 300 MG/2 ML
- DUPIXENT SYRINGE SUBCUTANEOUS SYRINGE 100 MG/0.67 ML, 200 MG/1.14 ML, 300 MG/2 ML

PA Criteria	Criteria Details
Exclusion Criteria	PENDING CMS REVIEW
Required Medical Information	PENDING CMS REVIEW
Age Restrictions	PENDING CMS REVIEW
Prescriber Restrictions	PENDING CMS REVIEW
Coverage Duration	PENDING CMS REVIEW
Other Criteria	PENDING CMS REVIEW
Indications	PENDING CMS REVIEW
Off-Label Uses	PENDING CMS REVIEW
Part B Prerequisite	PENDING CMS REVIEW

# EMGALITY

## Products Affected

- EMGALITY PEN
- EMGALITY SUBCUTANEOUS SYRINGE 120 MG/ML

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	Combination therapy with another cGRP inhibitor for migraine headache prevention
<b>Required Medical Information</b>	Diagnosis, number of migraine or cluster headaches per month
<b>Age Restrictions</b>	18 years and older
<b>Prescriber Restrictions</b>	N/A
<b>Coverage Duration</b>	Cluster headache tx-6 months, migraine prevention-1 year
<b>Other Criteria</b>	Migraine headache prevention-Approve if the patient meets the following criteria (A and B): A) Patient has greater than or equal to 4 migraine headache days per month (prior to initiating a migraine-preventative medication), AND B) If pt is currently taking Emgality, pt has had significant clinical benefit from the medication. Episodic cluster headache treatment-approve if the patient has between one headache every other day and eight headaches per day.
<b>Indications</b>	All FDA-approved Indications.
<b>Off-Label Uses</b>	N/A
<b>Part B Prerequisite</b>	No

# ENBREL

## Products Affected

- ENBREL MINI
- ENBREL SUBCUTANEOUS SOLUTION
- ENBREL SUBCUTANEOUS SYRINGE
- ENBREL SURECLICK

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	Concurrent use with biologic therapy or targeted synthetic DMARD
<b>Required Medical Information</b>	Diagnosis, concurrent medications, previous therapies tried.
<b>Age Restrictions</b>	PP-4 years and older (initial therapy)
<b>Prescriber Restrictions</b>	Initial only-RA/AS/JIA/JRA,prescribed by or in consult w/ rheumatologist. Psoriatic arthritis, prescribed by or in consultation w/ rheumatologist or dermatologist.Plaque psoriasis (PP), prescribed by or in consult w/ dermatologist.GVHD,prescribed by or in consult w/ oncologist,hematologist,or physician affiliated w/ transplant center.Behcet's disease,prescribed by or in consult w/ rheumatologist,dermatologist,ophthalmologist,gastroenterologist,or neurologist.
<b>Coverage Duration</b>	Approve through end of plan year
<b>Other Criteria</b>	INITIAL THERAPY: RHEUMATOID ARTHRITIS (RA): tried one conventional synthetic DMARD for at least 3 months (see Note 1). JUVENILE IDIOPATHIC ARTHRITIS (JIA)/JRA (one of A, B, C, or D): A) patient has aggressive disease, B) tried one other systemic therapy (e.g., methotrexate [MTX], sulfasalazine, leflunomide, NSAID, or a biologic that is not a biosimilar of the requested product), C) patient will be started on Enbrel concurrently with MTX, sulfasalazine, or leflunomide, or D) patient has an absolute contraindication to MTX, sulfasalazine, or leflunomide. PLAQUE PSORIASIS (PP) [A or B]: A) tried at least one traditional systemic agent for at least 3 months, unless intolerant (e.g., MTX, cyclosporine, Soriatane, oral methoxsalen plus PUVA) [see Note 1] or B) patient has a contraindication to one oral agent for psoriasis such as MTX. GRAFT VERSUS HOST DISEASE (GVHD): approve. BEHCET'S: tried

PA Criteria	Criteria Details
	at least one conventional therapy (e.g., systemic corticosteroid, immunosuppressant, interferon alfa, mycophenolate), adalimumab, or infliximab. CONTINUATION THERAPY: ALL INDICATIONS: patient had a response to therapy. Note 1: a biologic that is not a biosimilar of the requested product will also count.
<b>Indications</b>	All FDA-approved Indications, Some Medically-accepted Indications.
<b>Off-Label Uses</b>	Graft versus host disease (GVHD), Behcet's disease
<b>Part B Prerequisite</b>	No



## ENDARI

### Products Affected

- *glutamine (sickle cell)*

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	Diagnosis, prescriber specialty
<b>Age Restrictions</b>	Greater than or equal to 5 years of age
<b>Prescriber Restrictions</b>	Prescribed by, or in consultation with, a physician who specializes in sickle cell disease (e.g., a hematologist)
<b>Coverage Duration</b>	Authorization will be for 1 year.
<b>Other Criteria</b>	N/A
<b>Indications</b>	All FDA-approved Indications.
<b>Off-Label Uses</b>	N/A
<b>Part B Prerequisite</b>	No

# EPIDIOLEX

## Products Affected

- EPIDIOLEX

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	Diagnosis, previous therapies
<b>Age Restrictions</b>	Patients 1 year and older (initial therapy)
<b>Prescriber Restrictions</b>	Prescribed by or in consultation with a neurologist (initial therapy)
<b>Coverage Duration</b>	1 year
<b>Other Criteria</b>	Dravet Syndrome-approve if the patient has tried or is concomitantly receiving at least two other antiseizure drugs or if the patient has tried or is concomitantly receiving one of Diacomit or clobazam or Fintepla. Lennox Gastaut Syndrome-approve if the patient has tried or is concomitantly receiving at least two other antiseizure drugs. Tuberous Sclerosis Complex-approve if the patient has tried or is concomitantly receiving at least two other antiseizure drugs. Continuation of therapy-approve if the patient is responding to therapy.
<b>Indications</b>	All FDA-approved Indications.
<b>Off-Label Uses</b>	N/A
<b>Part B Prerequisite</b>	No

# EPOETIN ALFA

## Products Affected

- PROCREDIT INJECTION SOLUTION                      20,000 UNIT/ML, 3,000 UNIT/ML, 4,000 UNIT/ML, 10,000 UNIT/ML, 2,000 UNIT/ML, 40,000 UNIT/ML
- RETACRIT

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	Diagnosis
<b>Age Restrictions</b>	MDS anemia = 18 years of age and older
<b>Prescriber Restrictions</b>	MDS anemia, myelofibrosis- prescribed by or in consultation with, a hematologist or oncologist.
<b>Coverage Duration</b>	Chemo-6m,Transfus-1m, CKD-1yr, Myelofibrosis-init-3 mo, cont-1 yr, all others-1 yr
<b>Other Criteria</b>	Anemia in a pt with Chronic Kidney Disease (CKD) not on dialysis- for initial therapy, approve if hemoglobin (Hb) is less than 10.0 g/dL for adults or less than or equal to 11 g/dL for children, or for continuation of therapy in a pt currently on an erythropoiesis-stimulating agent (ESA) approve if Hb is less than or equal to 12 g/dL. Anemia in a pt with cancer due to chemotherapy- approve if pt is currently receiving myelosuppressive chemo as a non-curative treatment and (for initial therapy) Hb is less than 10.0 g/dL or (if currently on ESA) Hb is less than or equal to 12.0 g/dL. Anemia in HIV with zidovudine- for initial therapy, approve if Hb is less than 10.0 g/dL or serum erythropoietin level is 500 mU/mL or less, or for continuation of therapy in a pt currently on ESA, approve if Hb is 12.0 g/dL or less. Surgical pts to reduce RBC transfusions - Approve if Hb is less than or equal to 13, AND surgery is elective, nonvascular and non-cardiac AND pt is unwilling or unable to donate autologous blood prior to surgery. MDS- for initial therapy, approve if Hb is less than 10 g/dL or serum erythropoietin level is 500 mU/mL or less, or for continuation of therapy in a pt currently on ESA approve if Hb is 12.0 g/dL or less. Myelofibrosis- for Initial therapy approve if patient has a Hb less than 10

PA Criteria	Criteria Details
	or serum erythropoietin less than or equal to 500 mU/mL, or for continuation of therapy in pt currently on ESA hemoglobin is less than or equal to 12g/dL. Anemia in patients with chronic renal failure on dialysis - deny under Medicare Part D (claim should be submitted under the ESRD bundled payment benefit).
<b>Indications</b>	All FDA-approved Indications, Some Medically-accepted Indications.
<b>Off-Label Uses</b>	Anemia due to myelodysplastic syndrome (MDS), Myelofibrosis
<b>Part B Prerequisite</b>	No

# ERIVEDGE

## Products Affected

- ERIVEDGE

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	BCC (La or Met) - must not have had disease progression while on Odomzo.
<b>Required Medical Information</b>	N/A
<b>Age Restrictions</b>	18 years and older
<b>Prescriber Restrictions</b>	N/A
<b>Coverage Duration</b>	1 year
<b>Other Criteria</b>	Basal cell carcinoma, locally advanced-patients new to therapy-approve if the patient has tried Odomzo. Central nervous system cancer (this includes brain and spinal cord tumors)-approve if the patient has medulloblastoma, the patient has tried at least one chemotherapy agent and according to the prescriber, the patient has a mutation of the sonic hedgehog pathway. Basal cell carcinoma, metastatic (this includes primary or recurrent nodal metastases and distant metastases)-approve.
<b>Indications</b>	All FDA-approved Indications, Some Medically-accepted Indications.
<b>Off-Label Uses</b>	Central nervous System Cancer
<b>Part B Prerequisite</b>	No

# ERLEADA

## Products Affected

- ERLEADA ORAL TABLET 240 MG, 60 MG

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	Diagnosis
<b>Age Restrictions</b>	18 years and older
<b>Prescriber Restrictions</b>	N/A
<b>Coverage Duration</b>	1 year
<b>Other Criteria</b>	Prostate cancer-non-metastatic, castration resistant and prostate cancer-metastatic, castration sensitive-approve if the requested medication will be used in combination with a gonadotropin-releasing hormone (GnRH) analog [for example: leuprolide acetate, Lupron Depot (leuprolide acetate intramuscular injection), Trelstar (triptorelin pamoate intramuscular injection), Zoladex (goserelin acetate subcutaneous implant), Vantas (histrelin acetate subcutaneous implant), Firmagon (degarelix subcutaneous injection), Orgovyx (relugolix tablets)] or if the patient has had a bilateral orchiectomy.
<b>Indications</b>	All FDA-approved Indications.
<b>Off-Label Uses</b>	N/A
<b>Part B Prerequisite</b>	No

# ERLOTINIB

## Products Affected

- *erlotinib oral tablet 100 mg, 150 mg, 25 mg*

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	Diagnosis
<b>Age Restrictions</b>	18 years and older
<b>Prescriber Restrictions</b>	N/A
<b>Coverage Duration</b>	Authorization will be for 1 year
<b>Other Criteria</b>	Advanced or Metastatic NSCLC, approve if the patient has EGFR mutation positive non-small cell lung cancer as detected by an approved test. Note-Examples of EGFR mutation-positive non-small cell lung cancer include the following mutations: exon 19 deletions, exon 21 (L858R) substitution mutations, L861Q, G719X and S768I. RCC, approve if the patient has recurrent or advanced non-clear cell histology RCC or if the patient had hereditary leiomyomatosis and renal cell carcinoma and erlotinib will be used in combination with bevacizumab. Bone cancer-approve if the patient has chordoma and has tried at least one previous therapy. Pancreatic cancer-approve if the medication is used in combination with gemcitabine and if the patient has locally advanced, metastatic or recurrent disease. Vulvar cancer-approve if the patient has advanced, recurrent or metastatic disease.
<b>Indications</b>	All FDA-approved Indications, Some Medically-accepted Indications.
<b>Off-Label Uses</b>	Renal Cell Carcinoma, vulvar cancer and Bone Cancer-Chordoma.
<b>Part B Prerequisite</b>	No

# EVEROLIMUS

## Products Affected

- *everolimus (antineoplastic) oral tablet*
- *everolimus (antineoplastic) oral tablet for suspension 2 mg, 3 mg, 5 mg*
- *torpenz*

PA Criteria	Criteria Details
Exclusion Criteria	PENDING CMS REVIEW
Required Medical Information	PENDING CMS REVIEW
Age Restrictions	PENDING CMS REVIEW
Prescriber Restrictions	PENDING CMS REVIEW
Coverage Duration	PENDING CMS REVIEW
Other Criteria	PENDING CMS REVIEW
Indications	PENDING CMS REVIEW
Off-Label Uses	PENDING CMS REVIEW
Part B Prerequisite	PENDING CMS REVIEW



# FASENRA

## Products Affected

- FASENRA PEN
- FASENRA SUBCUTANEOUS SYRINGE 10 MG/0.5 ML, 30 MG/ML

PA Criteria	Criteria Details
Exclusion Criteria	PENDING CMS REVIEW
Required Medical Information	PENDING CMS REVIEW
Age Restrictions	PENDING CMS REVIEW
Prescriber Restrictions	PENDING CMS REVIEW
Coverage Duration	PENDING CMS REVIEW
Other Criteria	PENDING CMS REVIEW
Indications	PENDING CMS REVIEW
Off-Label Uses	PENDING CMS REVIEW
Part B Prerequisite	PENDING CMS REVIEW

# FINGOLIMOD

## Products Affected

- *fingolimod*

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	Concurrent use of fingolimod with other disease-modifying agents used for multiple sclerosis (MS).
<b>Required Medical Information</b>	Relapsing form of Multiple Sclerosis (MS), to include clinically-isolated syndrome, relapsing-remitting disease, and active secondary progressive disease
<b>Age Restrictions</b>	10 years and older
<b>Prescriber Restrictions</b>	Prescribed by, or in consultation with, a neurologist or an MS specialist.
<b>Coverage Duration</b>	Authorization will be for 1 year.
<b>Other Criteria</b>	N/A
<b>Indications</b>	All FDA-approved Indications.
<b>Off-Label Uses</b>	N/A
<b>Part B Prerequisite</b>	No

# FINTEPLA

## Products Affected

- FINTEPLA

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	Diagnosis
<b>Age Restrictions</b>	2 years and older (initial therapy)
<b>Prescriber Restrictions</b>	Prescribed by or in consultation with an neurologist (initial therapy)
<b>Coverage Duration</b>	1 year
<b>Other Criteria</b>	Dravet Syndrome-Initial therapy-approve if the patient has tried or is concomitantly receiving at least two other antiepileptic drugs or patient has tried or is concomitantly receiving Epidiolex, Clobazam or Diacomit. Dravet Syndrome-Continuation-approve if the patient is responding to therapy. Lennox-Gastaut Syndrome, initial-approve if the patient has tried or is concomitantly receiving at least two other antiepileptic drugs. Lennox-Gastaut Syndrome, continuation-approve if the patient is responding to therapy.
<b>Indications</b>	All FDA-approved Indications.
<b>Off-Label Uses</b>	N/A
<b>Part B Prerequisite</b>	No

# FIRMAGON

## Products Affected

- FIRMAGON KIT W DILUENT SYRINGE

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	Diagnosis
Age Restrictions	N/A
Prescriber Restrictions	Prescribed by or in consultation with a oncologist
Coverage Duration	1 year
Other Criteria	Part B versus Part D determination will be made at time of prior authorization review per CMS guidance. Patients new to therapy, are required to try Eligard or Orgovyx prior to approval of Firmagon.
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

# FOTIVDA

## Products Affected

- FOTIVDA

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	Diagnosis, other therapies
<b>Age Restrictions</b>	18 years and older
<b>Prescriber Restrictions</b>	N/A
<b>Coverage Duration</b>	Authorization will be for 1 year
<b>Other Criteria</b>	Renal Cell Carcinoma (RCC)-approve if the patient has relapsed or Stage IV disease and has tried at least two other systemic regimens. Note: Examples of systemic regimens for renal cell carcinoma include axitinib tablets, axitinib + pembrolizumab injection, cabozantinib tablets, cabozantinib + nivolumab injection, sunitinib malate capsules, pazopanib tablets, sorafenib tablets, and lenvatinib capsules + everolimus.
<b>Indications</b>	All FDA-approved Indications.
<b>Off-Label Uses</b>	N/A
<b>Part B Prerequisite</b>	No

# FRUZAQLA

## Products Affected

- FRUZAQLA ORAL CAPSULE 1 MG, 5 MG

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	Diagnosis
<b>Age Restrictions</b>	18 years of age and older
<b>Prescriber Restrictions</b>	N/A
<b>Coverage Duration</b>	1 year
<b>Other Criteria</b>	Colon cancer, rectal cancer, or appendiceal cancer-Approve if the patient meets the following (A and B): A.Patient has advanced or metastatic disease, AND B.Patient has previously been treated with the following (i, ii, and iii): i.Fluoropyrimidine-, oxaliplatin-, and irinotecan-based chemotherapy, Note: Examples of fluoropyrimidine agents include 5-fluorouracil (5-FU) and capecitabine. AND ii.An anti-vascular endothelial growth factor (VEGF) agent, Note: Examples of anti-VEGF agents include bevacizumab. AND iii. If the tumor is RAS wild-type (KRAS wild-type and NRAS wild-type) [that is, the tumor or metastases are KRAS and NRAS mutation negative], the patient meets ONE of the following (a or b): a.According to the prescriber, anti-epidermal growth factor receptor (EGFR) therapy is NOT medically appropriate, OR b. The patient has received an anti-EGFR therapy. Note: Examples of anti-EGFR therapy includes Erbitux (cetuximab intravenous infusion) and Vectibix (panitumumab intravenous infusion).
<b>Indications</b>	All FDA-approved Indications, Some Medically-accepted Indications.
<b>Off-Label Uses</b>	Appendiceal cancer

PA Criteria	Criteria Details
Part B Prerequisite	No

# FULPHILA

## Products Affected

- FULPHILA

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	Diagnosis
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	Cancer patients receiving chemotherapy-prescribed by or in consultation with an oncologist or hematologist. PBPC - prescribed by or in consultation with an oncologist, hematologist, or physician that specializes in transplantation.
<b>Coverage Duration</b>	Cancer pts receiving chemo-6 mo. PBPC-30 days.
<b>Other Criteria</b>	Cancer patients receiving chemotherapy, approve if the patient meets one of the following: 1) is receiving myelosuppressive anti-cancer medications that are associated with a high risk of febrile neutropenia (the risk is at least 20% based on the chemotherapy regimen), 2) patient is receiving myelosuppressive anti-cancer medications that are associated with a risk of febrile neutropenia but the risk is less than 20% based on the chemotherapy regimen and the patient has one or more risk factors for febrile neutropenia according to the prescribing physician (eg, aged greater than or equal to 65 years, prior chemotherapy or radiation therapy, persistent neutropenia, bone marrow involvement by tumor, recent surgery and/or open wounds, liver and/or renal dysfunction, poor performance status or HIV infection, or 3) patient has had a neutropenic complication from prior chemotherapy and did not receive prophylaxis with a colony stimulating factor and a reduced dose or frequency of chemotherapy may compromise treatment.
<b>Indications</b>	All FDA-approved Indications, Some Medically-accepted Indications.
<b>Off-Label Uses</b>	Patients undergoing PBPC collection and therapy



PA Criteria	Criteria Details
Part B Prerequisite	No

# GATTEX

## Products Affected

- GATTEX 30-VIAL

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	Diagnosis
<b>Age Restrictions</b>	1 year and older
<b>Prescriber Restrictions</b>	Prescribed by or in consultation with a gastroenterologist (initial and continuation)
<b>Coverage Duration</b>	1 year
<b>Other Criteria</b>	Initial-approve if the patient is currently receiving parenteral nutrition on 3 or more days per week or according to the prescriber, the patient is unable to receive adequate total parenteral nutrition required for caloric needs. Continuation-approve if the patient has experienced improvement.
<b>Indications</b>	All FDA-approved Indications.
<b>Off-Label Uses</b>	N/A
<b>Part B Prerequisite</b>	No

# GAVRETO

## Products Affected

- GAVRETO

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	Diagnosis
<b>Age Restrictions</b>	NSCLC-18 years and older, thyroid cancer-12 years and older
<b>Prescriber Restrictions</b>	N/A
<b>Coverage Duration</b>	1 year
<b>Other Criteria</b>	NSCLC-approve if the patient has advanced, recurrent, or metastatic disease and rearranged during transfection (RET) fusion-positive disease detected by an Food and Drug Administration (FDA) approved test. Differentiated Thyroid Cancer- pt has unresectable, recurrent, or metastatic disease AND pt has RET fusion-positive or RET-mutation-positive disease AND disease requires treatment with systemic therapy AND the disease is radioactive iodine-refractory. Anaplastic thyroid cancer or Medullary Thyroid Cancer- pt has unresectable, recurrent, or metastatic disease AND pt has RET fusion-positive or RET-mutation-positive disease.
<b>Indications</b>	All FDA-approved Indications, Some Medically-accepted Indications.
<b>Off-Label Uses</b>	Medullary Thyroid Cancer, Anaplastic Thyroid Cancer
<b>Part B Prerequisite</b>	No

# GEFITINIB

---

## Products Affected

- *gefitinib*

PA Criteria	Criteria Details
Exclusion Criteria	PENDING CMS REVIEW
Required Medical Information	PENDING CMS REVIEW
Age Restrictions	PENDING CMS REVIEW
Prescriber Restrictions	PENDING CMS REVIEW
Coverage Duration	PENDING CMS REVIEW
Other Criteria	PENDING CMS REVIEW
Indications	PENDING CMS REVIEW
Off-Label Uses	PENDING CMS REVIEW
Part B Prerequisite	PENDING CMS REVIEW

# GILOTRIF

## Products Affected

- GILOTRIF

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	For NSCLC - EGFR exon deletions or mutations, or if NSCLC is squamous cell type
<b>Age Restrictions</b>	18 years and older
<b>Prescriber Restrictions</b>	N/A
<b>Coverage Duration</b>	Authorization will be for 1 year
<b>Other Criteria</b>	NSCLC EGFR pos - For the treatment of advanced or metastatic non small cell lung cancer (NSCLC)-approve if the patient has sensitizing EGFR mutation-positive NSCLC as detected by an approved test. Note: examples of sensitizing EGFR mutation-positive NSCLC include the following mutations : exon 19 deletions, exon 21 (L858R) substitution mutations, L861Q, G719X and S768I. NSCLC metastatic squamous cell must have disease progression after treatment with platinum based chemotherapy. Head and neck cancer-approve if the patient has non-nasopharyngeal head and neck cancer and the patient has disease progression on or after platinum based chemotherapy. (Part B before Part D Step Therapy - applies only to beneficiaries enrolled in an MA-PD plan)
<b>Indications</b>	All FDA-approved Indications, Some Medically-accepted Indications.
<b>Off-Label Uses</b>	Head and neck cancer
<b>Part B Prerequisite</b>	Yes

# GLATIRAMER

## Products Affected

- *glatiramer subcutaneous syringe 20 mg/ml, 40 mg/ml*
- *glatopa subcutaneous syringe 20 mg/ml, 40 mg/ml*

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	Concurrent use with other disease-modifying agent used for multiple sclerosis
<b>Required Medical Information</b>	Relapsing form of Multiple Sclerosis (MS), to include clinically-isolated syndrome, relapsing-remitting disease, and active secondary progressive disease
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	Prescribed by or after consultation with a neurologist or an MS specialist.
<b>Coverage Duration</b>	Authorization will be for 1 year.
<b>Other Criteria</b>	N/A
<b>Indications</b>	All FDA-approved Indications.
<b>Off-Label Uses</b>	N/A
<b>Part B Prerequisite</b>	No

## GLUCAGON-LIKE PEPTIDE-1 AGONISTS

### Products Affected

- BYDUREON BCISE
- BYETTA SUBCUTANEOUS PEN INJECTOR 10 MCG/DOSE(250 MCG/ML) 2.4 ML, 5 MCG/DOSE (250 MCG/ML) 1.2 ML
- MOUNJARO
- OZEMPIC SUBCUTANEOUS PEN INJECTOR 0.25 MG OR 0.5 MG (2 MG/3 ML), 1 MG/DOSE (4 MG/3 ML), 2 MG/DOSE (8 MG/3 ML)
- RYBELSUS
- TRULICITY

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	Diagnosis
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Authorization will be for 1 year
Other Criteria	N/A
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

# GONADOTROPIN-RELEASING HORMONE AGONISTS - ONCOLOGY

## Products Affected

- ELIGARD
- ELIGARD (3 MONTH)
- ELIGARD (4 MONTH)
- ELIGARD (6 MONTH)
- *leuprolide subcutaneous kit*

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	Diagnosis
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	Prostate cancer- prescribed by or in consultation with an oncologist or urologist. Head and neck-salivary gland tumors- prescribed by or in consultation with an oncologist.
<b>Coverage Duration</b>	1 year
<b>Other Criteria</b>	Head and neck cancer-salivary gland tumor- approve if pt has recurrent, unresectable, or metastatic disease AND androgen receptor-positive disease.
<b>Indications</b>	All FDA-approved Indications, Some Medically-accepted Indications.
<b>Off-Label Uses</b>	Head and neck cancer- salivary gland tumors (Eligard only)
<b>Part B Prerequisite</b>	No



## GRALISE/HORIZANT/LYRICA CR

### Products Affected

- gabapentin oral tablet extended release 24 hr 300 mg, 600 mg

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	N/A
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Authorization will be for 12 months.
Other Criteria	N/A
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

# GROWTH HORMONES

## Products Affected

- OMNITROPE

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	GHD in Children/Adolescents. Pt meets one of the following-1-had 2 GH stim tests with the following-levodopa, insulin-induced hypoglycemia, arginine, clonidine, or glucagon and both are less than 10ng/mL OR had at least 1 GH test and results are less than 10ng/mL and has at least one risk factor for GHD (e.g., ht for age curve deviated down across 2 major height percentiles [e.g., from above the 25 percentile to below the 10 percentile], growth rate is less than the expected normal growth rate based on age and gender, low IGF-1 and/or IGFBP-3 levels). 2.brain radiation or tumor resection and pt has 1 GH stim test less than 10ng/mL or has def in at least 1 other pituitary hormone (that is, ACTH, TSH, gonadotropin deficiency [LH and/or FSH] are counted as 1 def], or prolactin).3. congenital hypopituitarism and has one GH stim test less than 10ng/mL OR def in at least one other pituitary hormone and/or the patient has the imaging triad of ectopic posterior pituitary and pituitary hypoplasia with abnormal pituitary stalk 4.pt has multiple pituitary deficiencies and pt has 3 or more pituitary hormone deficiencies or pt has had one GH test less than 10ng/mL 5.pt had a hypophysectomy. Cont-pt responding to therapy
<b>Age Restrictions</b>	ISS 5 y/o or older, SGA 2 y/o or older, SBS 18 y/o or older
<b>Prescriber Restrictions</b>	GHD (Initial tx children or adolescents w/o hypophysectomy), GHD adults or transitional adolescents, Noonan (initial), Prader Willi (initial for child/adult and cont tx in adults), SHOX (initial), SGA (initial) - prescribed by or in consultation with an endocrinologist. CKD (initial) endocrinologist or nephrologist.
<b>Coverage Duration</b>	ISS - 6 mos initial, 12 months cont tx, SBS 1 month, others 12 mos
<b>Other Criteria</b>	GHD initial in adults and adolescents 1. endocrine must certify not being prescribed for anti-aging or to enhance athletic performance, 2. has either childhood onset or adult onset resulting from GHD alone, multiple

PA Criteria	Criteria Details
	<p>hormone deficiency from pituitary dx, hypothalamic dz, pituitary surgery, cranial radiation tx, tumor treatment, TBI or subarachnoid hemorrhage, AND 3. meets one of the following - A. has known perinatal insults, congenital or genetic defects or structural hypothalamic pituitary defects, B. 3 or more pituitary hormone def (ACTH, TSH, LH/FSH, or prolactin, age and gender adjusted IGF1 below the lower limits of the normal reference range AND other causes of low serum IGF-1 have been excluded, C. Neg response to ONE preferred GH stim test (insulin peak response less than or equal to 5 mcg/L, Glucagon peak less than or equal to 3 mcg/L (BMI is less than or equal to 25), less than or equal to 3 and BMI is greater than or equal to 25 and less than or equal to 30 with a high pretest probability of GH deficiency, less than or equal to 1 and BMI is greater than or equal to 25 and less than or equal to 30 with a low pretest probability of GH deficiency or less than or equal to 1 mcg/L (BMI is greater than 30), if insulin and glucagon contraindicated then Arginine test with peak of less than or equal to 0.4 mcg/L, or Macrilen peak less than 2.8 ng/ml AND BMI is less than or equal to 40 AND if a transitional adolescent must be off tx for at least one month before retesting. Cont tx - endocrine must certify not being prescribed for anti-aging or to enhance athletic performance. ISS initial - baseline ht less than the 1.2 percentile or a standard deviation score (SDS) less than -2.25 for age and gender, open epiphyses, does not have CDGP and height velocity is either growth rate (GR) is a. less than 4 cm/yr for pts greater than or equal to 5 or b. growth velocity is less than 10th percentile for age/gender. Cont tx - prescriber confirms response to therapy. CKD initial - CKD defined by abnormal CrCl. Noonan initial - baseline height less than 5th percentile. PW cont tx in adults or adolescents who don't meet child requir - physician certifies not being used for anti-aging or to enhance athletic performance. SHOX initial - SHOX def by chromo analysis, open epiphyses, height less than 3rd percentile for age/gender. SGA initial -baseline ht less than 5th percentile for age/gender and born SGA (birth weight/length that is more than 2 SD below mean for gestational age/gender and didn't have sufficient catch up growth by 2-4 y/o). Cont tx - prescriber confirms response to therapy. Cont Tx for CKD, Noonan, PW in child/adoles, SHOX, and TS - prescriber confirms response to therapy. SBS initial pt receiving specialized nutritional support. Cont tx - 2nd course if pt responded to tx with a decrease in the requirement for specialized nutritional support.</p>
<b>Indications</b>	All FDA-approved Indications, Some Medically-accepted Indications.
<b>Off-Label Uses</b>	Short bowel syndrome

PA Criteria	Criteria Details
Part B Prerequisite	No

## HIGH RISK MEDICATIONS - BENZODIAZEPINES

### Products Affected

- *clorazepate dipotassium oral tablet 15 mg, 3.75 mg, 7.5 mg*
- *diazepam intensol*
- *diazepam oral solution 5 mg/5 ml (1 mg/ml)*
- *diazepam oral tablet*
- *lorazepam intensol*
- *lorazepam oral tablet 0.5 mg, 1 mg, 2 mg*

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	N/A
<b>Age Restrictions</b>	Patients aged less than 65 years, approve. Patients aged 65 years and older, other criteria apply.
<b>Prescriber Restrictions</b>	N/A
<b>Coverage Duration</b>	Procedure-related sedation = 1mo. All other conditions = 12 months.
<b>Other Criteria</b>	All medically accepted indications other than insomnia, approve if the physician has assessed risk versus benefit in using the High Risk Medication (HRM) in this patient and has confirmed that he/she would still like to initiate/continue therapy. Insomnia, may approve lorazepam if the patient has had a trial with two of the following: ramelteon, doxepin 3mg or 6 mg, eszopiclone, zolpidem, or zaleplon and the physician has assessed risk versus benefit in using the HRM in this patient and has confirmed that he/she would still like initiate/continue therapy.
<b>Indications</b>	All Medically-accepted Indications.
<b>Off-Label Uses</b>	N/A
<b>Part B Prerequisite</b>	No

## HIGH RISK MEDICATIONS - BENZTROPINE

### Products Affected

- *benztropine oral*

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	N/A
<b>Age Restrictions</b>	Patients aged less than 65 years, approve. Patients aged 65 years and older, other criteria apply
<b>Prescriber Restrictions</b>	N/A
<b>Coverage Duration</b>	12 months
<b>Other Criteria</b>	For all medically-accepted indications, approve if the prescriber confirms he/she has assessed risk versus benefit in prescribing benztropine for the patient and he/she would still like to initiate/continue therapy
<b>Indications</b>	All Medically-accepted Indications.
<b>Off-Label Uses</b>	N/A
<b>Part B Prerequisite</b>	No

# HIGH RISK MEDICATIONS - CYCLOBENZAPRINE

## Products Affected

- *cyclobenzaprine oral tablet 10 mg, 5 mg*

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	N/A
<b>Age Restrictions</b>	Patients aged less than 65 years, approve.
<b>Prescriber Restrictions</b>	N/A
<b>Coverage Duration</b>	Authorization will be for 12 months.
<b>Other Criteria</b>	The physician has assessed risk versus benefit in using this High Risk Medication (HRM) in this patient and has confirmed that he/she would still like to initiate/continue therapy.
<b>Indications</b>	All Medically-accepted Indications.
<b>Off-Label Uses</b>	N/A
<b>Part B Prerequisite</b>	No

# HIGH RISK MEDICATIONS - FIRST GENERATION ANTIHISTAMINES

## Products Affected

- *hydroxyzine hcl oral tablet*
- *promethazine oral*

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	N/A
<b>Age Restrictions</b>	Patients aged less than 65 years, approve. Patients aged 65 years and older, other criteria apply.
<b>Prescriber Restrictions</b>	N/A
<b>Coverage Duration</b>	Authorization will be for 12 months.
<b>Other Criteria</b>	For hydroxyzine hydrochloride, authorize use without a previous drug trial for all FDA-approved indications other than anxiety. Approve hydroxyzine hydrochloride if the patient has tried at least two other FDA-approved products for the management of anxiety. Prior to approval of promethazine and hydroxyzine, approve if the physician must have assessed risk versus benefit in prescribing the requested HRM for the patient and must confirm that he/she would still like to initiate/continue therapy.
<b>Indications</b>	All Medically-accepted Indications.
<b>Off-Label Uses</b>	N/A
<b>Part B Prerequisite</b>	No



## HIGH RISK MEDICATIONS - PHENOBARBITAL

### Products Affected

- *phenobarbital*

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	Coverage is not provided for use in sedation/insomnia.
<b>Required Medical Information</b>	N/A
<b>Age Restrictions</b>	Patients aged less than 65 years, approve. Patients aged 65 years and older, other criteria apply.
<b>Prescriber Restrictions</b>	N/A
<b>Coverage Duration</b>	12 months
<b>Other Criteria</b>	For the treatment of seizures, approve only if the patient is currently taking phenobarbital.
<b>Indications</b>	All Medically-accepted Indications.
<b>Off-Label Uses</b>	N/A
<b>Part B Prerequisite</b>	No

## HIGH RISK MEDICATIONS- ESTROGENS

### Products Affected

- *dotti*
- *estradiol oral*
- *estradiol transdermal patch semiweekly*
- *estradiol transdermal patch weekly*
- *estradiol-norethindrone acet*
- *fyavolv*
- *jinteli*
- *lyllana*
- *mimvey*
- *norethindrone ac-eth estradiol oral tablet*  
0.5-2.5 mg-mcg, 1-5 mg-mcg

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	Previous medication use
<b>Age Restrictions</b>	Patients aged less than 65 years, approve. Patients aged 65 years and older, other criteria apply.
<b>Prescriber Restrictions</b>	N/A
<b>Coverage Duration</b>	Authorization will be for 12 months
<b>Other Criteria</b>	For the treatment of Vulvar Vaginal Atrophy, approve if the patient has had a trial of one of the following for vulvar vaginal atrophy (brand or generic): Estradiol Vaginal Cream, Invexxy, Premarin Vaginal Cream or estradiol valerate injection. For prophylaxis of Postmenopausal Osteoporosis, approve if the patient has had a trial of one of the following (brand or generic): alendronate, ibandronate, risedronate or Raloxifene. The physician has assessed risk versus benefit in using this High Risk medication (HRM) in this patient and has confirmed that he/she would still like to initiate/continue therapy.
<b>Indications</b>	All Medically-accepted Indications.
<b>Off-Label Uses</b>	N/A
<b>Part B Prerequisite</b>	No

# IBRANCE

## Products Affected

- IBRANCE

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	N/A
<b>Age Restrictions</b>	18 years and older
<b>Prescriber Restrictions</b>	N/A
<b>Coverage Duration</b>	1 year
<b>Other Criteria</b>	Breast cancer - approve recurrent or metastatic, hormone receptor positive (HR+) [i.e., estrogen receptor positive- [ER+] and/or progesterone receptor positive [PR+]] disease, and HER2-negative breast cancer when the pt meets ONE of the following 1. Pt is postmenopausal and Ibrance will be used in combination with anastrozole, exemestane, or letrozole 2, pt is premenopausal or perimenopausal and is receiving ovarian suppression/ablation with GnRH agonists, or has had surgical bilateral oophorectomy, or ovarian irradiation AND meets one of the following conditions: Ibrance will be used in combination with anastrozole, exemestane, or letrozole or Ibrance will be used in combination with fulvestrant 3. pt is a man (a man is defined as an individual with the biological traits of a man, regardless of the individual's gender identity or gender expression) who is receiving GnRH analog AND Ibrance with be used in combination with anastrozole, exemestane or letrozole or Ibrance will be used in combination with fulvestrant 4. Pt is postmenopausal and Ibrance will be used in combination with fulvestrant. In addition, patients new to therapy must have a trial of Kisqali, Kisqali Femara Co-Pack or Verzenio prior to approval of Ibrance. Liposarcoma-approve if the patient has well-differentiated/dedifferentiated liposarcoma (WD-DDLS).
<b>Indications</b>	All FDA-approved Indications, Some Medically-accepted Indications.

PA Criteria	Criteria Details
Off-Label Uses	Liposarcoma
Part B Prerequisite	No

# ICATIBANT

## Products Affected

- *icatibant*
- *sajazir*

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	N/A
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	Prescribed by, or in consultation with, an allergist/immunologist or a physician that specializes in the treatment of HAE or related disorders
<b>Coverage Duration</b>	Authorization will be for 1 year.
<b>Other Criteria</b>	Hereditary Angioedema (HAE) Due to C1 Inhibitor (C1-INH) Deficiency [Type I or Type II] - Treatment of Acute Attacks, Initial Therapy-the patient has HAE type I or type II as confirmed by the following diagnostic criteria (i and ii): i. the patient has low levels of functional C1-INH protein (less than 50 percent of normal) at baseline, as defined by the laboratory reference values AND ii. the patient has lower than normal serum C4 levels at baseline, as defined by the laboratory reference values. Patients who have treated previous acute HAE attacks with icatibant - the patient has treated previous acute HAE type I or type II attacks with icatibant AND according to the prescribing physician, the patient has had a favorable clinical response (e.g., decrease in the duration of HAE attacks, quick onset of symptom relief, complete resolution of symptoms, decrease in HAE acute attack frequency or severity) with icatibant treatment.
<b>Indications</b>	All FDA-approved Indications.
<b>Off-Label Uses</b>	N/A
<b>Part B Prerequisite</b>	No

# ICLUSIG

## Products Affected

- ICLUSIG

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	Diagnosis the Philadelphia chromosome (Ph) status of the leukemia must be reported. T315I status
<b>Age Restrictions</b>	18 years and older
<b>Prescriber Restrictions</b>	N/A
<b>Coverage Duration</b>	Authorization will be for 1 year
<b>Other Criteria</b>	Acute lymphoblastic leukemia, Philadelphia chromosome positive or chronic myeloid leukemia-approve. GIST - approve if the patient tried all of the following therapies first to align with NCCN recommendations which include: Imatinib or Ayyakit (avapritinib tablets), AND Sunitinib or Sprycel (dasatinib tablets), AND Stivarga (regorafenib tablets), AND Qinlock (ripretinib tablets). Myeloid/Lymphoid Neoplasms with Eosinophilia - approve if the tumor has ABL1 rearrangement or FGFR1 rearrangement.
<b>Indications</b>	All FDA-approved Indications, Some Medically-accepted Indications.
<b>Off-Label Uses</b>	Gastrointestinal Stromal Tumor, Myeloid/Lymphoid Neoplasms with Eosinophilia
<b>Part B Prerequisite</b>	No

## IDHIFA

### Products Affected

- IDHIFA

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	IDH2-mutation status
Age Restrictions	18 years and older
Prescriber Restrictions	N/A
Coverage Duration	Authorization will be for 1 year
Other Criteria	AML - approve if the patient is IDH2-mutation status positive as detected by an approved test
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

# IMATINIB

## Products Affected

- *imatinib oral tablet 100 mg, 400 mg*

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	Diagnosis. For indications of CML and ALL, the Philadelphia chromosome (Ph) status of the leukemia must be reported.
<b>Age Restrictions</b>	ASM, DFSP, HES, MDS/MPD/Myeloid/Lymphoid Neoplasms/Kaposi Sarcoma/Cutaneous Melanoma-18 years and older
<b>Prescriber Restrictions</b>	N/A
<b>Coverage Duration</b>	1 year
<b>Other Criteria</b>	For ALL/CML, must have Ph-positive for approval of imatinib. Kaposi's Sarcoma-approve if the patient has tried at least one regimen AND has relapsed or refractory disease. Pigmented villonodular synovitis/tenosynovial giant cell tumor (PVNS/TGCT)-patient has tried Turalio or according to the prescriber, the patient cannot take Turalio. Myelodysplastic/myeloproliferative disease-approve if the condition is associated with platelet-derived growth factor receptor (PDGFR) gene rearrangements.Graft versus host disease, chronic-approve if the patient has tried at least one conventional systemic treatment (e.g., imbruvica). Cutaneous melanoma-approve if the patient has an activating KIT mutation, metastatic or unresectable melanoma, and has tried at least one systemic regimen. Myeloid/lymphoid neoplasms with eosinophilia-approve if the tumor has an ABL1 rearrangement or an FIP1L1-PDGFR or PDGFRB rearrangement.
<b>Indications</b>	All FDA-approved Indications, Some Medically-accepted Indications.
<b>Off-Label Uses</b>	Chordoma, desmoid tumors (aggressive fibromatosis), cKit positive metastatic or unresectable cutaneous melanoma, Kaposi's Sarcoma and



PA Criteria	Criteria Details
	pigmented Villonodular Synovitis/Tenosynovial Giant Cell Tumor, myeloid/lymphoid neoplasms with eosinophilia, GVHD, chronic.
<b>Part B Prerequisite</b>	No

# IMBRUVICA

## Products Affected

- IMBRUVICA ORAL CAPSULE 140 MG, 70 MG
- IMBRUVICA ORAL TABLET 140 MG, 280 MG, 420 MG
- IMBRUVICA ORAL SUSPENSION

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	Diagnosis
<b>Age Restrictions</b>	GVHD-1 year and older, other-18 years and older
<b>Prescriber Restrictions</b>	N/A
<b>Coverage Duration</b>	1 year
<b>Other Criteria</b>	CLL- Approve. GVHD-Approve if the patient has tried one conventional systemic treatment for graft versus host disease (e.g., corticosteroids [methylprednisolone, prednisone], imatinib, low-dose methotrexate, sirolimus, mycophenolate mofetil, Jakafi [ruxolitinib tablets]). B-cell lymphoma-approve if the patient has tried at least one systemic regimen (e.g., cisplatin, cytarabine, rituximab, oxaliplatin, gemcitabine, ifosfamide, carboplatin, etoposide, or rituximab). Central nervous system Lymphoma (primary)- approve if the patient is not a candidate for or is intolerant to high-dose methotrexate OR has tried at least one therapy (e.g., methotrexate, rituximab, vincristine, procarbazine, cytarabine, thiotepa, carmustine, intrathecal methotrexate, cytarabine, or rituximab). Hairy Cell Leukemia - approve if the patient has tried at least two systemic regimens (cladribine, Nipent [pentostatin injection], rituximab, or Pegasys [peginterferon alfa-2a subcutaneous injection]).
<b>Indications</b>	All FDA-approved Indications, Some Medically-accepted Indications.
<b>Off-Label Uses</b>	Central Nervous System Lymphoma (Primary), Hairy Cell Leukemia, B-Cell Lymphoma

PA Criteria	Criteria Details
Part B Prerequisite	No

# INBRIJA

## Products Affected

- INBRIJA INHALATION CAPSULE, W/INHALATION DEVICE

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	Asthma, COPD, other chronic underlying lung disease
<b>Required Medical Information</b>	Diagnosis, medications that will be used in combination
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	Prescribed by or in consultation with a neurologist
<b>Coverage Duration</b>	1 year
<b>Other Criteria</b>	Approve if the patient is currently taking carbidopa-levodopa and is experiencing off episodes.
<b>Indications</b>	All FDA-approved Indications.
<b>Off-Label Uses</b>	N/A
<b>Part B Prerequisite</b>	No

# INGREZZA

## Products Affected

- INGREZZA
- INGREZZA INITIATION PK(TARDIV)
- INGREZZA SPRINKLE

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	Diagnosis
Age Restrictions	18 years and older
Prescriber Restrictions	TD - Prescribed by or in consultation with a neurologist or psychiatrist. Chorea HD - prescribed by or in consultation with a neurologist
Coverage Duration	1 year
Other Criteria	Chorea associated with Huntington's Disease- approve if diagnosis is confirmed by genetic testing (for example, an expanded HTT CAG repeat sequence of at least 36).
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

# INJECTABLE TESTOSTERONE PRODUCTS

## Products Affected

- *testosterone cypionate*
- *testosterone enanthate*

PA Criteria	Criteria Details
Exclusion Criteria	PENDING CMS REVIEW
Required Medical Information	PENDING CMS REVIEW
Age Restrictions	PENDING CMS REVIEW
Prescriber Restrictions	PENDING CMS REVIEW
Coverage Duration	PENDING CMS REVIEW
Other Criteria	PENDING CMS REVIEW
Indications	PENDING CMS REVIEW
Off-Label Uses	PENDING CMS REVIEW
Part B Prerequisite	PENDING CMS REVIEW

# INLYTA

## Products Affected

- INLYTA ORAL TABLET 1 MG, 5 MG

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	N/A
Age Restrictions	18 years and older
Prescriber Restrictions	N/A
Coverage Duration	Authorization will be for 1 year
Other Criteria	Advanced Renal cell carcinoma-approve. Differentiated thyroid cancer, approve if patient is refractory to radioactive iodine therapy. Soft tissue sarcoma-approve if the patient has alveolar soft part sarcoma and the medication will be used in combination with Keytruda (pembrolizumab).
Indications	All FDA-approved Indications, Some Medically-accepted Indications.
Off-Label Uses	Differentiated (i.e., papillary, follicular, and oncocytic) Thyroid Carcinoma, Soft tissue sarcoma
Part B Prerequisite	No

## INPEFA

### Products Affected

- INPEFA

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	Diagnosis
<b>Age Restrictions</b>	18 years and older
<b>Prescriber Restrictions</b>	N/A
<b>Coverage Duration</b>	1 year
<b>Other Criteria</b>	Heart Failure, to reduce the risk of cardiovascular death, hospitalization for heart failure, and urgent heart failure visit-approve. Type 2 Diabetes, to reduce the risk of cardiovascular death, hospitalization for heart failure, and urgent heart failure visit-approve if the patient has chronic kidney disease AND has one or more cardiovascular risk factor(s).Note: Patients with heart failure should be reviewed under criteria for Heart Failure.
<b>Indications</b>	All FDA-approved Indications.
<b>Off-Label Uses</b>	N/A
<b>Part B Prerequisite</b>	No



# INQOVI

## Products Affected

- INQOVI

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	Diagnosis
Age Restrictions	18 years and older
Prescriber Restrictions	N/A
Coverage Duration	1 year
Other Criteria	N/A
Indications	All FDA-approved Indications, Some Medically-accepted Indications.
Off-Label Uses	Myelodysplastic Syndrome/Myeloproliferative Neoplasm Overlap Neoplasms
Part B Prerequisite	No

# INREBIC

## Products Affected

- INREBIC

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	Diagnosis
Age Restrictions	18 years and older
Prescriber Restrictions	N/A
Coverage Duration	1 year
Other Criteria	Myelofibrosis (MF), including Primary MF, Post-Polycythemia Vera MF, and Post-Essential Thrombocythemia MF-approve if the patient has intermediate-2 or high-risk disease. Myeloid/Lymphoid Neoplasms with Eosinophilia-approve if the tumor has a JAK2 rearrangement.
Indications	All FDA-approved Indications, Some Medically-accepted Indications.
Off-Label Uses	Myeloid/Lymphoid Neoplasms with Eosinophilia
Part B Prerequisite	No

## IVERMECTIN (ORAL)

### Products Affected

- *ivermectin oral*

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	Diagnosis
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	N/A
<b>Coverage Duration</b>	30 days
<b>Other Criteria</b>	Pediculosis-approve if the patient has infection caused by pediculus humanus capitis (head lice), pediculus humanus corporis (body lice), or has pediculosis pubis caused by phthirus pubis (pubic lice). Scabies-approve if the patient has classic scabies, treatment resistant scabies, is unable to tolerate topical treatment, has crusted scabies or is using ivermectin tablets for prevention and/or control of scabies.
<b>Indications</b>	All FDA-approved Indications, Some Medically-accepted Indications.
<b>Off-Label Uses</b>	Ascariasis, Enterobiasis (pinworm infection), Hookworm-related cutaneous larva migrans, Mansonella ozzardi infection, Mansonella streptocerca infection, Pediculosis, Scabies. Trichuriasis, Wucheria bancrofti infection
<b>Part B Prerequisite</b>	No

## IVIG

### Products Affected

- PRIVIGEN

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	N/A
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Authorization will be for 12 months.
Other Criteria	Part B versus D determination per CMS guidance to establish if drug used for PID in pt's home.
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

# IWILFIN

## Products Affected

- IWILFIN

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	Diagnosis
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	1 year
Other Criteria	Neuroblastoma-Approve if the patient meets the following (A, B and C): A) Patient has high-risk disease, AND B) The medication is being used to reduce the risk of relapse, AND C) Patient has had at least a partial response to prior multiagent, multimodality therapy including anti-GD2 immunotherapy. Note: Examples of anti-GD2 immunotherapy includes Unituxin (dinutuximab intravenous infusion).
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

# JAKAFI

## Products Affected

- JAKAFI

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	N/A
<b>Age Restrictions</b>	ALL-less than 21 years of age, GVHD-12 and older, MF/PV/CMML-2/essential thrombo/myeloid/lymphoid neoplasm/T-cell Lymphoma-18 and older
<b>Prescriber Restrictions</b>	N/A
<b>Coverage Duration</b>	1 year
<b>Other Criteria</b>	For polycythemia vera patients must have tried hydroxyurea or peginterferon alfa-2a or Besremi (ropeginterferon alfa-2b-njft subcutaneous injection). ALL-approve if the mutation/pathway is Janus associated kinase (JAK)-related. GVHD, chronic-approve if the patient has tried one conventional systemic treatment for graft versus host disease (for example: prednisone, ibrutinib capsules/tablets). GVHD, acute-approve if the patient has tried one systemic corticosteroid. Atypical chronic myeloid leukemia-approve if the patient has a CSF3R mutation or a janus associated kinase 2 (JAK2) mutation. Chronic monomyelocytic leukemia-2 (CMML-2)-approve if the patient is also receiving a hypomethylating agent. Essential thrombocythemia-approve if the patient has tried hydroxyurea, peginterferon alfa-2a or anagrelide. Myeloid/lymphoid neoplasms-approve if the patient has eosinophilia and the tumor has a janus associated kinase 2 (JAK2) rearrangement. T-Cell Lymphoma - approve if pt has T-cell prolymphocytic leukemia or T-cell large granular lymphocytic leukemia AND pt has tried at least one systemic regimen.
<b>Indications</b>	All FDA-approved Indications, Some Medically-accepted Indications.

PA Criteria	Criteria Details
Off-Label Uses	Acute lymphoblastic leukemia, atypical chronic myeloid leukemia, chronic monomyelocytic leukemia-2 (CMML-2), essential thrombocythemia, myeloid/lymphoid neoplasms, T-Cell lymphoma
Part B Prerequisite	No

# JAYPIRCA

## Products Affected

- JAYPIRCA ORAL TABLET 100 MG, 50 MG

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	Diagnosis
<b>Age Restrictions</b>	18 years and older
<b>Prescriber Restrictions</b>	N/A
<b>Coverage Duration</b>	1 year
<b>Other Criteria</b>	<p>Mantle cell lymphoma-approve if the patient has tried at least one systemic regimen or patient is not a candidate for a systemic regimen (i.e., an elderly patient who is frail), AND the patient has tried one Bruton tyrosine kinase inhibitor (BTK) for mantle cell lymphoma. Note: Examples of a systemic regimen contain one or more of the following products: rituximab, dexamethasone, cytarabine, carboplatin, cisplatin, oxaliplatin, cyclophosphamide, doxorubicin, vincristine, prednisone, methotrexate, bendamustine, Velcade (bortezomib intravenous or subcutaneous injection), lenalidomide, gemcitabine, and Venclexta (venetoclax tablets). Note: Examples of BTK inhibitors indicated for mantle cell lymphoma include Brukinsa (zanubrutinib capsules), Calquence (acalabrutinib capsules), and Imbruvica (ibrutinib capsules, tablets, and oral suspension). CLL/SLL-patient meets (A or B): A) patient has resistance or intolerance to Imbruvica (ibrutinib tablets, capsules, or oral solution), Calquence (acalabrutinib tablets), or Brukinsa (zanubrutinib capsules) or B) patient has relapsed or refractory disease and has tried a Bruton tyrosine kinase (BTK) inhibitor and Venclexta (venetoclax tablet)-based regimen. Examples of BTK inhibitor include: Imbruvica (ibrutinib tablets, capsules, or oral solution), Calquence (acalabrutinib tablets), or Brukinsa</p>



PA Criteria	Criteria Details
	(zanubrutinib capsules). Richter's Transformation to DLBCL- pt has tried at least one chemotherapy regimen or is not a candidate for a chemotherapy regimen.
<b>Indications</b>	All FDA-approved Indications, Some Medically-accepted Indications.
<b>Off-Label Uses</b>	Richter's Transformation to Diffuse Large B-Cell Lymphoma
<b>Part B Prerequisite</b>	No

# KALYDECO

## Products Affected

- KALYDECO

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	Combination use with Orkambi, Trikafta or Symdeko
<b>Required Medical Information</b>	N/A
<b>Age Restrictions</b>	1 month of age and older
<b>Prescriber Restrictions</b>	prescribed by or in consultation with a pulmonologist or a physician who specializes in CF
<b>Coverage Duration</b>	1 year
<b>Other Criteria</b>	CF - must meet A, B, and C: A) pt must have one mutation in the CFTR gene that is considered to be pathogenic or likely pathogenic B) pt must have positive CF newborn screening test or family history of CF or clinical presentation consistent with signs and symptoms of CF and C) evidence of abnormal CFTR function as demonstrated by i, ii, or iii: (i) elevated sweat chloride test or (ii) two CFTR mutations or (iii) abnormal nasal potential difference.
<b>Indications</b>	All FDA-approved Indications.
<b>Off-Label Uses</b>	N/A
<b>Part B Prerequisite</b>	No

# KERENDIA

## Products Affected

- KERENDIA

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	Concomitant use with spironolactone or eplerenone
<b>Required Medical Information</b>	Diagnosis
<b>Age Restrictions</b>	18 years and older (initial and continuation therapy)
<b>Prescriber Restrictions</b>	N/A
<b>Coverage Duration</b>	1 year
<b>Other Criteria</b>	<p>Diabetic kidney disease-initial-approve if the patient meets the following criteria (i, ii and iii): i. Patient has a diagnosis of type 2 diabetes AND ii. Patient meets one of the following (a or b): a) Patient is currently receiving a maximally tolerated labeled dosage of an angiotensin converting enzyme (ACE) inhibitor or angiotensin receptor blocker (ARB) OR b) According to the prescriber, patient has a contraindication to ACE inhibitor and ARB therapy AND iii. At baseline (prior to the initiation of Kerendia), patient meets all of the following (a, b, and c): a) Estimated glomerular filtration rate greater than or equal to 25 mL/min/1.73 m<sup>2</sup> AND b) Urine albumin-to-creatinine ratio greater than or equal to 30 mg/g AND c) Serum potassium level less than or equal to 5.0 mEq/L.</p> <p>Diabetic kidney disease-continuation-approve if the patient meets the following criteria (i and ii): i. Patient has a diagnosis of type 2 diabetes AND ii. Patient meets one of the following (a or b): a) Patient is currently receiving a maximally tolerated angiotensin converting enzyme (ACE) inhibitor or angiotensin receptor blocker (ARB) OR b) According to the prescriber, patient has a contraindication to ACE inhibitor and ARB therapy.</p>
<b>Indications</b>	All FDA-approved Indications.
<b>Off-Label Uses</b>	N/A

PA Criteria	Criteria Details
Part B Prerequisite	No

# KESIMPTA

---

## Products Affected

- KESIMPTA PEN

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	Concurrent use with other disease-modifying agents used for multiple sclerosis (MS)
<b>Required Medical Information</b>	Relapsing form of Multiple Sclerosis (MS), to include, clinically-isolated syndrome, relapsing-remitting disease, and active secondary progressive disease.
<b>Age Restrictions</b>	18 years and older
<b>Prescriber Restrictions</b>	Prescribed by or in consultation with a neurologist or a physician who specializes in the treatment of multiple sclerosis
<b>Coverage Duration</b>	Authorization will be for 1 year
<b>Other Criteria</b>	N/A
<b>Indications</b>	All FDA-approved Indications.
<b>Off-Label Uses</b>	N/A
<b>Part B Prerequisite</b>	No

# KISQALI

## Products Affected

- KISQALI FEMARA CO-PACK ORAL TABLET 200 MG/DAY(200 MG X 1)-2.5 MG, 400 MG/DAY(200 MG X 2)-2.5 MG, 600 MG/DAY(200 MG X 3)-2.5 MG
- KISQALI ORAL TABLET 200 MG/DAY (200 MG X 1), 400 MG/DAY (200 MG X 2), 600 MG/DAY (200 MG X 3)

PA Criteria	Criteria Details
Exclusion Criteria	PENDING CMS REVIEW
Required Medical Information	PENDING CMS REVIEW
Age Restrictions	PENDING CMS REVIEW
Prescriber Restrictions	PENDING CMS REVIEW
Coverage Duration	PENDING CMS REVIEW
Other Criteria	PENDING CMS REVIEW
Indications	PENDING CMS REVIEW
Off-Label Uses	PENDING CMS REVIEW
Part B Prerequisite	PENDING CMS REVIEW

# KORLYM

## Products Affected

- *mifepristone oral tablet 300 mg*

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	Diagnosis, prior surgeries
<b>Age Restrictions</b>	18 years and older
<b>Prescriber Restrictions</b>	Prescribed by or in consultation with an endocrinologist or a physician who specializes in the treatment of Cushing's syndrome
<b>Coverage Duration</b>	1 year
<b>Other Criteria</b>	Endogenous Cushing's Syndrome-Approve if mifepristone is being used to control hyperglycemia secondary to hypercortisolism in patients who have type 2 diabetes mellitus or glucose intolerance AND pt meets (i, ii or iii): i) patient is not a candidate for surgery or surgery has not been curative, or (ii) patient is awaiting surgery for endogenous Cushing's Syndrome or (iii) patient is awaiting therapeutic response after radiotherapy for endogenous Cushing's Syndrome.
<b>Indications</b>	All FDA-approved Indications.
<b>Off-Label Uses</b>	N/A
<b>Part B Prerequisite</b>	No

# KOSELUGO

## Products Affected

- KOSELUGO

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	Diagnosis
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	1 year
Other Criteria	<p>Neurofibromatosis Type 1-approve if prior to starting Koselugo, the patient has symptomatic, inoperable plexiform neurofibromas and if the patient is 2 to 18 years old OR if the patient is 19 years or older if the patient started on therapy with Koselugo prior to becoming 19. Circumscribed Glioma-approve if the patient has recurrent, refractory or progressive disease AND the tumor is BRAF fusion positive OR BRAF V600E activating mutation positive OR patient has neurofibromatosis type 1 mutated glioma AND this medication will be used as a single agent AND the patient is 3-21 years of age OR is greater 21 and has been previously started on therapy with Koselugo prior to becoming 21 years of age. Langerhans Cell Histiocytosis- approve if the patient meets the following criteria (A and B):</p> <p>A) Patient meets one of the following (i, ii, iii, or iv): i. Patient meets both of the following (a and b): a) Patient has multisystem Langerhans cell histiocytosis, AND b) Patient has symptomatic disease or impending organ dysfunction, OR ii. Patient has single system lung Langerhans cell histiocytosis, OR iii. Patient meets all of the following (a, b, and c): a) Patient has single system bone disease, AND b) Patient has not responded to treatment with a bisphosphonate, AND c) Patient has more than 2 bone lesions, OR iv. Patient has central nervous system disease, AND B) The medication is used as a single agent.</p>



PA Criteria	Criteria Details
<b>Indications</b>	All FDA-approved Indications, Some Medically-accepted Indications.
<b>Off-Label Uses</b>	Circumscribed Glioma, Langerhans Cell Histiocytosis
<b>Part B Prerequisite</b>	No

# KRAZATI

---

## Products Affected

- KRAZATI

PA Criteria	Criteria Details
Exclusion Criteria	PENDING CMS REVIEW
Required Medical Information	PENDING CMS REVIEW
Age Restrictions	PENDING CMS REVIEW
Prescriber Restrictions	PENDING CMS REVIEW
Coverage Duration	PENDING CMS REVIEW
Other Criteria	PENDING CMS REVIEW
Indications	PENDING CMS REVIEW
Off-Label Uses	PENDING CMS REVIEW
Part B Prerequisite	PENDING CMS REVIEW

# LAPATINIB

## Products Affected

- *lapatinib*

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	Diagnosis for which lapatinib is being used. Metastatic breast cancer, HER2 status or hormone receptor (HR) status.
<b>Age Restrictions</b>	18 years and older
<b>Prescriber Restrictions</b>	N/A
<b>Coverage Duration</b>	1 year
<b>Other Criteria</b>	HER2-positive recurrent or metastatic breast cancer, approve if lapatinib will be used in combination with capecitabine OR trastuzumab and the patient has tried at least two prior anti-HER2 based regimens OR the medication will be used in combination with an aromatase inhibitor and and the patient has HR+ disease and the patient is a postmenopausal woman or the patient is premenopausal or perimenopausal woman and is receiving ovarian suppression/ablation with a GnRH agonist, surgical bilateral oophorectomy or ovarian irradiation OR the patient is a man and is receiving a GnRH analog. Colon or rectal cancer-approve if the patient has unresectable advanced or metastatic disease that is human epidermal receptor 2 (HER2) amplified and with wild-type RAS and BRAF disease and the patient has tried at least one chemotherapy regimen or is not a candidate for intensive therapy and the medication is used in combination with trastuzumab (Part B before Part D Step Therapy - applies only to beneficiaries enrolled in an MA-PD plan) and the patient has not been previously treated with a HER2-inhibitor. Bone Cancer-approve if the patient has recurrent chordoma and if the patient has epidermal growth-factor receptor (EGFR)-positive disease.
<b>Indications</b>	All FDA-approved Indications, Some Medically-accepted Indications.

PA Criteria	Criteria Details
Off-Label Uses	Bone cancer-chordoma, colon or rectal cancer, breast cancer in pre/perimenopausal women and men
Part B Prerequisite	Yes

## LEDIPASVIR/SOFOSBUVIR

### Products Affected

- LEDIPASVIR-SOFOSBUVIR

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	Combination use with other direct acting antivirals, excluding ribavirin
<b>Required Medical Information</b>	Diagnosis
<b>Age Restrictions</b>	3 years or older
<b>Prescriber Restrictions</b>	Prescribed by or in consultation w/ GI, hepatologist, ID, or a liver transplant MD
<b>Coverage Duration</b>	Criteria will be applied consistent with current AASLD/IDSA guidance.
<b>Other Criteria</b>	Criteria will be applied consistent with current AASLD/IDSA guidance.
<b>Indications</b>	All FDA-approved Indications, Some Medically-accepted Indications.
<b>Off-Label Uses</b>	Indications consistent with current AASLD/IDSA guidance
<b>Part B Prerequisite</b>	No

# LENALIDOMIDE

## Products Affected

- *lenalidomide*
- REVLIMID

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	Diagnosis and previous therapies or drug regimens tried.
<b>Age Restrictions</b>	18 years and older (except Kaposi's Sarcoma, Castleman's Disease, CNS Lymphoma)
<b>Prescriber Restrictions</b>	N/A
<b>Coverage Duration</b>	Authorization will be for 1 year
<b>Other Criteria</b>	Follicular lymphoma-approve if the patient is using lenalidomide in combination with rituximab or has tried at least one prior therapy. MCL-approve -if the patient is using lenalidomide in combination with rituximab or has tried at least two other therapies or therapeutic regimens. MZL-approve if the patient is using lenalidomide in combination with rituximab or has tried at least one other therapy or therapeutic regimen. Multiple myeloma-approve. MDS-approve if the patient meets one of the following: 1) Pt has symptomatic anemia, OR 2) Pt has transfusion-dependent anemia, OR 3) Pt has anemia that is not controlled with an erythroid stimulating agent (eg, Epogen, Procrit [epoetin alfa injection], Aranesp [darbepoetin alfa injection]). B-cell-lymphoma (other)-approve if the pt has tried at least one prior therapy. Myelofibrosis-approve if according to the prescriber the patient has anemia and the pt has serum erythropoietin levels greater than or equal to 500 mU/mL or according to the prescriber the patient has anemia, has serum erythropoietin levels less than 500 mU/mL and patient has experienced no response or loss of response to erythropoietic stimulating agents. Peripheral T-Cell Lymphoma or T-Cell Leukemia/Lymphoma-approve if the pt has tried at least one other therapy or regimen. CNS lymphoma-approve if according to the prescriber the

PA Criteria	Criteria Details
	<p>patient has relapsed or refractory disease. Hodgkin lymphoma, classical-approve if the patient has tried at least three other regimens. Castleman's disease-approve if the patient has relapsed/refractory or progressive disease. Kaposi's Sarcoma-approve if the patient has tried at least one regimen or therapy and the patient has relapsed or refractory disease. Systemic light chain amyloidosis-approve if lenalidomide is used in combination with dexamethasone.</p>
<b>Indications</b>	All FDA-approved Indications, Some Medically-accepted Indications.
<b>Off-Label Uses</b>	<p>Systemic Amyloidosis Light Chain, Diffuse Large B Cell Lymphoma (Non-Hodgkin's Lymphoma), Myelofibrosis. Castleman's Disease, Hodgkin lymphoma (Classical), Peripheral T-Cell Lymphoma, T-Cell Leukemia/Lymphoma, Central nervous system Lymphoma, Kaposi's sarcoma.</p>
<b>Part B Prerequisite</b>	No

# LENVIMA

## Products Affected

- LENVIMA ORAL CAPSULE 10 MG/DAY (10 MG X 1), 12 MG/DAY (4 MG X 3), 14 MG/DAY(10 MG X 1-4 MG X 1), 18 MG/DAY (10 MG X 1-4 MG X2), 20 MG/DAY (10 MG X 2), 24 MG/DAY(10 MG X 2-4 MG X 1), 4 MG, 8 MG/DAY (4 MG X 2)

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	N/A
<b>Age Restrictions</b>	18 years and older
<b>Prescriber Restrictions</b>	N/A
<b>Coverage Duration</b>	1 year
<b>Other Criteria</b>	<p>DTC - must be refractory to radioactive iodine treatment for approval. RCC - approve if the pt meets ALL of the following criteria: 1) pt has advanced disease AND if the patient meets i or ii-i. Lenvima is is being used in combination with Keytruda OR ii. Lenvima is used in combination with Afinitor/Afinitor Disperz and the patient meets a or b-a. Patient has clear cell histology and patient has tried one antiangiogenic therapy or b. patient has non-clear cell histology. MTC-approve if the patient has tried at least one systemic therapy. Endometrial Carcinoma-Approve if the patient meets the following criteria (A, B, C, and D): A) The patient has advanced endometrial carcinoma that is not microsatellite instability-high (MSI-H) or mismatch repair deficient (dMMR) AND B) The medication is used in combination with Keytruda (pembrolizumab for intravenous injection) AND C)the disease has progressed on at least one prior systemic therapy AND D) The patient is not a candidate for curative surgery or radiation. HCC-approve if the patient has unresectable or metastatic disease. Thymic carcinoma-approve if the patient has tried at least one chemotherapy regimen. Melanoma - approve if the patient has unresectable or metastatic melanoma AND the medication will be used in combination with Keytruda</p>



PA Criteria	Criteria Details
	(pembrolizumab intravenous injection) AND the patient has disease progression on anti-programmed death receptor-1 (PD-1)/programmed death-ligand 1 (PD-L1)-based therapy.
<b>Indications</b>	All FDA-approved Indications, Some Medically-accepted Indications.
<b>Off-Label Uses</b>	Patients with Medullary Thyroid Carcinoma (MTC), thymic carcinoma, Renal cell carcinoma with non-clear cell histology and Melanoma
<b>Part B Prerequisite</b>	No

# LIBERVANT

## Products Affected

- LIBERVANT

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	Diagnosis
Age Restrictions	2 to 5 years of age
Prescriber Restrictions	N/A
Coverage Duration	1 year
Other Criteria	N/A
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

# LIDOCAINE PATCH

## Products Affected

- *lidocaine topical adhesive patch, medicated 5 %*
- *lidocan iii*
- *tridacaine ii*

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	N/A
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	N/A
<b>Coverage Duration</b>	Authorization will be for 12 months
<b>Other Criteria</b>	N/A
<b>Indications</b>	All FDA-approved Indications, Some Medically-accepted Indications.
<b>Off-Label Uses</b>	Diabetic neuropathic pain, chronic back pain
<b>Part B Prerequisite</b>	No

# LIVTENCITY

## Products Affected

- LIVTENCITY

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	Concomitant use with ganciclovir or valganciclovir
<b>Required Medical Information</b>	Diagnosis
<b>Age Restrictions</b>	12 years and older
<b>Prescriber Restrictions</b>	Prescribed by or in consultation with a hematologist, infectious diseases specialist, oncologist, or a physician affiliated with a transplant center.
<b>Coverage Duration</b>	2 months
<b>Other Criteria</b>	Cytomegalovirus Infection, Treatment-approve if the patient meets the following criteria (A, B, and C): A) Patient weighs greater than or equal to 35 kg, AND B) Patient is post-transplant, AND Note: This includes patients who are post hematopoietic stem cell transplant or solid organ transplant. C) Patient has cytomegalovirus infection/disease that is refractory to treatment with at least one of the following: cidofovir, foscarnet, ganciclovir, or valganciclovir or patient has a significant intolerance to ganciclovir or valganciclovir.
<b>Indications</b>	All FDA-approved Indications.
<b>Off-Label Uses</b>	N/A
<b>Part B Prerequisite</b>	No

## LONG ACTING OPIOIDS

### Products Affected

- BELBUCA
- buprenorphine transdermal patch
- hydromorphone oral tablet extended release 24 hr
- methadone oral solution 10 mg/5 ml, 5 mg/5 ml
- methadone oral tablet 10 mg, 5 mg
- morphine oral tablet extended release
- OXYCONTIN ORAL TABLET, ORAL ONLY, EXT.REL. 12 HR 10 MG, 15 MG, 20 MG, 30 MG, 40 MG, 60 MG, 80 MG

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	Acute (ie, non-chronic) pain
<b>Required Medical Information</b>	Pain type (chronic vs acute), prior pain medications/therapies tried, concurrent pain medications/therapies
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	N/A
<b>Coverage Duration</b>	Authorization will be for 12 months.
<b>Other Criteria</b>	For pain severe enough to require daily, around-the-clock, long-term opioid treatment, approve if all of the following criteria are met: 1) patient is not opioid naive, AND 2) non-opioid therapies have been tried and are being used in conjunction with opioid therapy according to the prescribing physician, AND 3) the prescribing physician has checked the patient's history of controlled substance prescriptions using state prescription drug monitoring program (PDMP), AND 4) the prescribing physician has discussed risks (eg, addiction, overdose) and realistic benefits of opioid therapy with the patient, AND 5) according to the prescriber physician there is a treatment plan (including goals for pain and function) in place and reassessments are scheduled at regular intervals. Patients with cancer, in hospice, sickle cell disease or who reside in a long term care facility are not required to meet above criteria. Clinical criteria incorporated into the quantity limit edits for all oral long-acting opioids require confirmation that the indication is intractable pain (ie, FDA labeled use) prior to reviewing for quantity exception.

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Indications</b>	All FDA-approved Indications.
<b>Off-Label Uses</b>	N/A
<b>Part B Prerequisite</b>	No

# LONSURF

## Products Affected

- LONSURF

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	N/A
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	N/A
<b>Coverage Duration</b>	Authorization will be for 1 year
<b>Other Criteria</b>	Gastric or Gastroesophageal Junction Adenocarcinoma, approve if the patient has been previously treated with at least two chemotherapy regimens for gastric or gastroesophageal junction adenocarcinoma. Colon and rectal cancer-approve per labeling if the patient has been previously treated with a fluoropyrimidine, oxaliplatin, irinotecan and if the patient's tumor or metastases are wild-type RAS (KRAS wild type and NRAS wild type) they must also try Erbitux or Vectibix.
<b>Indications</b>	All FDA-approved Indications.
<b>Off-Label Uses</b>	N/A
<b>Part B Prerequisite</b>	No

# LORBRENA

## Products Affected

- LORBRENA ORAL TABLET 100 MG, 25 MG

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	Diagnosis, ALK status, ROS1 status, previous therapies
<b>Age Restrictions</b>	18 years and older
<b>Prescriber Restrictions</b>	N/A
<b>Coverage Duration</b>	1 year
<b>Other Criteria</b>	Erdheim Chester disease-approve if the patient has anaplastic lymphoma kinase (ALK) rearrangement/fusion-positive disease. Inflammatory myofibroblastic tumor (IMT)-approve if the patient has IMT with ALK translocation. NSCLC - Approve if the patient has ALK-positive advanced or metastatic NSCLC, as detected by an approved test. In addition, patients new to therapy must also have a trial of Alecensa prior to approval of Lorbrena. NSCLC-ROS1 Rearrangement-Positive, advanced or metastatic NSCLC-approve if the patient has tried at least one of crizotinib, entrectinib or ceritinib.
<b>Indications</b>	All FDA-approved Indications, Some Medically-accepted Indications.
<b>Off-Label Uses</b>	Non-small cell lung cancer (NSCLC)-ROS1 Rearrangement-Positive, Erdheim Chester Disease, Inflammatory Myofibroblastic Tumor (IMT)
<b>Part B Prerequisite</b>	No



# LUMAKRAS

## Products Affected

- LUMAKRAS ORAL TABLET 120 MG, 320 MG

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	Diagnosis
<b>Age Restrictions</b>	18 years and older
<b>Prescriber Restrictions</b>	N/A
<b>Coverage Duration</b>	Authorization will be for 1 year
<b>Other Criteria</b>	Non-Small Cell Lung Cancer (NSCLC)-approve if the patient has KRAS G12C-mutated locally advanced or metastatic NSCLC, as determined by an FDA-approved test AND has been previously treated with at least one systemic regimen. Pancreatic Adenocarcinoma- approve if patient has KRAS G12C-mutated disease, as determined by an approved test AND either (i or ii): (i) patient has locally advanced or metastatic disease and has been previously treated with at least one systemic regimen OR (ii) patient has recurrent disease after resection.
<b>Indications</b>	All FDA-approved Indications, Some Medically-accepted Indications.
<b>Off-Label Uses</b>	Pancreatic Adenocarcinoma
<b>Part B Prerequisite</b>	No

# LUPRON DEPOT

## Products Affected

- LUPRON DEPOT

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	Diagnosis
<b>Age Restrictions</b>	Premenstrual disorders - 18 years and older
<b>Prescriber Restrictions</b>	Prostate cancer-prescribed by/consultation with oncologist or urologist. Other cancer diagnosis- prescribed by/consultation with an oncologist. Gender dysphoria/reassignment- prescribed by/consultation with endocrinologist or physician who specializes in treatment of transgender patients
<b>Coverage Duration</b>	uterine leiomyomata - 3 months, abnormal uterine bleeding - 6 months, all others - 12 months
<b>Other Criteria</b>	Endometriosis-approve if the pt has tried one of the following, unless contraindicated: a contraceptive, an oral progesterone or depo-medroxyprogesterone injection. An exception can be made if the pt has previously tried a gonadotropin-releasing hormone [GnRH] agonist (e.g. Lupron Depot) or antagonist (e.g. Orilissa). Head and neck cancer-salivary gland tumor- approve if pt has recurrent, unresectable, or metastatic disease AND androgen receptor-positive disease. Premenstrual disorders including PMS and PMDD- approve if pt has severe refractory premenstrual symptoms AND pt has tried an SSRI or combined oral contraceptive. Prostate cancer - for patients new to therapy requesting Lupron Depot 7.5mg, patients are required to try Orgovyx or Eligard prior to approval.
<b>Indications</b>	All FDA-approved Indications, Some Medically-accepted Indications.
<b>Off-Label Uses</b>	abnormal uterine bleeding, breast cancer, gender dysphoria/gender reassignment, head and neck cancer-salivary gland tumors, ovarian cancer including fallopian tube and primary peritoneal cancers, premenstrual disorders including premenstrual syndrome and premenstrual dysphoric

PA Criteria	Criteria Details
	disorder, prophylaxis or treatment of uterine bleeding or menstrual suppression in pts with hematologic malignancy or undergoing cancer treatment or prior to bone marrow or stem cell transplant, uterine cancer
<b>Part B Prerequisite</b>	No

# LYNPARZA

## Products Affected

- LYNPARZA

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	N/A
Age Restrictions	18 years and older
Prescriber Restrictions	N/A
Coverage Duration	1 year
Other Criteria	Ovarian, Fallopian Tube, or Primary Peritoneal Cancer - Maintenance monotherapy-Approve if the patient has a germline or somatic BRCA mutation-positive disease as confirmed by an approved test AND The patient is in complete or partial response to at least one platinum-based chemotherapy regimen (e.g., carboplatin with gemcitabine, carboplatin with paclitaxel, cisplatin with gemcitabine). Ovarian, fallopian tube, or primary peritoneal cancer-maintenance, combination therapy-approve if the medication is used in combination with bevacizumab, the patient has homologous recombination deficiency (HRD)-positive disease, as confirmed by an approved test and the patient is in complete or partial response to first-line platinum-based chemotherapy regimen. Breast cancer, adjuvant-approve if the patient has germline BRCA mutation-positive, HER2-negative breast cancer and the patient has tried neoadjuvant or adjuvant therapy. Breast cancer, recurrent or metastatic disease-approve if the patient has recurrent or metastatic disease and has germline BRCA mutation-positive breast cancer. Pancreatic Cancer-maintenance therapy-approve if the patient has a germline BRCA mutation-positive metastatic disease and the disease has not progressed on at least 16 weeks of treatment with a first-line platinum-based chemotherapy regimen. Prostate cancer-castration resistant-approve if the patient has metastatic disease, the

PA Criteria	Criteria Details
	<p>medication is used concurrently with a gonadotropin-releasing hormone (GnRH) analog or the patient has had a bilateral orchiectomy, and the patient meets either (i or ii): i) the patient has germline or somatic homologous recombination repair (HRR) gene-mutated disease, as confirmed by an approved test and the patient has been previously treated with at least one androgen receptor directed therapy or ii) the patient has a BRCA mutation and the medication is used in combination with abiraterone plus one of prednisone or prednisolone. Uterine Leiomyosarcoma-approve if the patient has BRCA2-altered disease and has tried one systemic regimen.</p>
<b>Indications</b>	All FDA-approved Indications, Some Medically-accepted Indications.
<b>Off-Label Uses</b>	Uterine Leiomyosarcoma
<b>Part B Prerequisite</b>	No

# LYTGOBI

## Products Affected

- LYTGOBI ORAL TABLET 12 MG/DAY (4 MG X 3), 16 MG/DAY (4 MG X 4), 20 MG/DAY (4 MG X 5)

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	Diagnosis
<b>Age Restrictions</b>	18 years and older
<b>Prescriber Restrictions</b>	N/A
<b>Coverage Duration</b>	1 year
<b>Other Criteria</b>	Cholangiocarcinoma-approve if the patient has unresectable locally advanced or metastatic disease, tumor has fibroblast growth factor receptor 2 (FGFR2) gene fusions or other rearrangements as detected by an approved test and if the patient has been previously treated with at least one systemic regimen. Note: Examples of systemic regimens include gemcitabine + cisplatin, 5-fluorouracil + oxaliplatin or cisplatin, capecitabine + cisplatin or oxaliplatin, gemcitabine + Abraxane (albumin-bound paclitaxel) or capecitabine or oxaliplatin, and gemcitabine + cisplatin + Abraxane.
<b>Indications</b>	All FDA-approved Indications.
<b>Off-Label Uses</b>	N/A
<b>Part B Prerequisite</b>	No

# MAVYRET

## Products Affected

- MAVYRET ORAL PELLETS IN PACKET
- MAVYRET ORAL TABLET

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	Diagnosis
Age Restrictions	3 years or older
Prescriber Restrictions	Prescribed by or in consultation with a gastroenterologist, hepatologist, infectious diseases physician, or a liver transplant physician
Coverage Duration	Criteria will be applied consistent with current AASLD/IDSA guidance.
Other Criteria	Criteria will be applied consistent with current AASLD/IDSA guidance.
Indications	All FDA-approved Indications, Some Medically-accepted Indications.
Off-Label Uses	Indications consistent with current AASLD/IDSA guidance
Part B Prerequisite	No

# MEGESTROL

## Products Affected

- *megestrol oral suspension 400 mg/10 ml (40 mg/ml), 625 mg/5 ml (125 mg/ml)*
- *megestrol oral tablet*

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	Coverage is not provided for weight gain for cosmetic reasons.
<b>Required Medical Information</b>	N/A
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	N/A
<b>Coverage Duration</b>	12 months
<b>Other Criteria</b>	N/A
<b>Indications</b>	All Medically-accepted Indications.
<b>Off-Label Uses</b>	N/A
<b>Part B Prerequisite</b>	No



# MEKINIST

## Products Affected

- MEKINIST ORAL RECON SOLN
- MEKINIST ORAL TABLET 0.5 MG, 2 MG

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	Diagnosis for which Mekinist is being used. For melanoma, thyroid cancer and NSCLC must have documentation of BRAF V600 mutations
<b>Age Restrictions</b>	1 year and older
<b>Prescriber Restrictions</b>	N/A
<b>Coverage Duration</b>	Authorization will be for 1 year
<b>Other Criteria</b>	Melanoma must be used in patients with BRAF V600 mutation, and patient has unresectable, advanced (including Stage III or Stage IV disease), or metastatic melanoma. Note-This includes adjuvant treatment in patients with Stage III disease with no evidence of disease post-surgery. For NSCLC requires BRAF V600E Mutation and use in combination with Tafenlar. Thyroid cancer, anaplastic-patient has locally advanced or metastatic anaplastic disease AND Mekinist will be taken in combination with Tafenlar, unless intolerant AND the patient has BRAF V600-positive disease. Ovarian/fallopian tube/primary peritoneal cancer-approve if the patient has recurrent disease and the medication is used for low-grade serous carcinoma or the patient has BRAF V600 mutation positive disease and the medication will be taken in combination with Tafenlar. Biliary Tract Cancer-approve if the patient has tried at least one systemic chemotherapy regimen, patient has BRAF V600 mutation positive disease and the medication will be taken in combination with Tafenlar. Central Nervous System Cancer-approve if the medication is being used for one of the following situations (i, ii, or iii): i. Adjuvant treatment of one of the following conditions (a, b, or c): a) Pilocytic astrocytoma OR b) Pleomorphic xanthoastrocytoma OR c) Ganglioglioma OR ii. Recurrent or

PA Criteria	Criteria Details
	<p>progressive disease for one of the following conditions (a, b, c or d): a) glioma OR b) Isocitrate dehydrogenase-2 (IDH2)-mutant astrocytoma OR c) Glioblastoma or d) Oligodendroglioma OR iii. Melanoma with brain metastases AND patient has BRAF V600 mutation-positive disease AND medication will be taken in combination with Tafinlar (dabrafenib). Histiocytic neoplasm-approve if patient has Langerhans cell histiocytosis and one of the following (a, b, or c): a) Multisystem disease OR b) Pulmonary disease OR c) Central nervous system lesions OR patient has Erdheim Chester disease or Rosai-Dorfman disease. Metastatic or Solid Tumors-Approve if the patient meets the following (A, B, and C): A) Patient has BRAF V600 mutation-positive disease, AND B) The medication will be taken in combination with Tafinlar (dabrafenib capsules), AND C) Patient has no satisfactory alternative treatment options. Hairy Cell Leukemia, approve if pt has not previously been treated with a BRAF inhibitor therapy and this will be used for relapsed/refractory disease and will be taken in combination with Tafinlar. Small bowel adenocarcinoma, approve if pt has BRAF V600E mutation-positive advanced or metastatic disease and this will be used with Tafinlar AND (i or ii): i) this will be used as initial therapy and pt has received previous FOLFOX/CAPEOX therapy in the adjuvant setting within the past 12 months or has a contraindication, or (ii) this will be used as second-line and subsequent therapy.</p>
<b>Indications</b>	All FDA-approved Indications, Some Medically-accepted Indications.
<b>Off-Label Uses</b>	Histiocytic Neoplasm, Hairy Cell Leukemia
<b>Part B Prerequisite</b>	No

# MEKTOVI

## Products Affected

- MEKTOVI

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	Diagnosis, BRAF V600 status, concomitant medications
<b>Age Restrictions</b>	18 years and older
<b>Prescriber Restrictions</b>	N/A
<b>Coverage Duration</b>	1 year
<b>Other Criteria</b>	Melanoma - approve if the patient has unresectable, advanced or metastatic melanoma AND has a BRAF V600 mutation AND Mektovi will be used in combination with Braftovi. In addition, patients new to therapy must have a trial with Cotellic or Mekinist prior to approval of Mektovi. Histiocytic neoplasm-approve if the patient has Langerhans cell histiocytosis and one of the following (i, ii, or iii): i. multisystem disease OR, ii. pulmonary disease or, iii. central nervous system lesions. NSCLC-approve if pt has BRAF V600E mutation-positive metastatic disease AND this medication will be taken in combination with Braftovi (encorafenib capsules).
<b>Indications</b>	All FDA-approved Indications, Some Medically-accepted Indications.
<b>Off-Label Uses</b>	Histiocytic Neoplasms
<b>Part B Prerequisite</b>	No

# MEMANTINE

## Products Affected

- *memantine oral capsule, sprinkle, er 24hr*
- *memantine oral solution*
- *memantine oral tablet*
- NAMZARIC

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	Indication for which memantine is being prescribed.
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	N/A
<b>Coverage Duration</b>	Authorization will be for 12 months.
<b>Other Criteria</b>	N/A
<b>Indications</b>	All FDA-approved Indications, Some Medically-accepted Indications.
<b>Off-Label Uses</b>	Patients with mild to moderate vascular dementia.
<b>Part B Prerequisite</b>	No

## MODAFINIL/ARMODAFINIL

### Products Affected

- *armodafinil*
- *modafinil oral tablet 100 mg, 200 mg*

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	N/A
<b>Age Restrictions</b>	18 years and older
<b>Prescriber Restrictions</b>	Excessive daytime sleepiness associated with narcolepsy-prescribed by or in consultation with a sleep specialist physician or neurologist
<b>Coverage Duration</b>	Authorization will be for 12 months.
<b>Other Criteria</b>	Excessive sleepiness associated with Shift Work Sleep Disorder (SWSD) - approve if patient is working at least 5 overnight shifts per month. Adjunctive/augmentation treatment for depression in adults if the patient is concurrently receiving other medication therapy for depression. Excessive daytime sleepiness associated with obstructive sleep apnea/hypoapnea syndrome-approve. Excessive daytime sleepiness associated with Narcolepsy-approve if narcolepsy has been confirmed with polysomnography and a multiple sleep latency test (MSLT).
<b>Indications</b>	All FDA-approved Indications, Some Medically-accepted Indications.
<b>Off-Label Uses</b>	Excessive daytime sleepiness (EDS) associated with myotonic dystrophy (modafinil only). Adjunctive/augmentation for treatment of depression in adults (modafinil only).
<b>Part B Prerequisite</b>	No

# MYFEMBREE

## Products Affected

- MYFEMBREE

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	Diagnosis, test results
<b>Age Restrictions</b>	18 years and older
<b>Prescriber Restrictions</b>	Fibroids-Prescribed by or in consultation with an obstetrician-gynecologist or a health care practitioner who specializes in the treatment of women's health
<b>Coverage Duration</b>	24 months of total therapy between Myfembree or Oriahnn
<b>Other Criteria</b>	Uterine Fibroids (Leiomyomas)-approve if the patient is premenopausal (before menopause) and is experiencing heavy menstrual bleeding associated with the uterine fibroids, the uterine fibroids have been confirmed by a pelvic ultrasound, including transvaginal ultrasonography or sonohysterography, hysteroscopy, or magnetic resonance imaging. Endometriosis-approve if the patient is premenopausal and patient has previously tried one of the following (i or ii): i. A contraceptive (e.g., combination oral contraceptives, levonorgestrel-releasing intrauterine systems, a depo-medroxyprogesterone injection), unless contraindicated OR ii. An oral progesterone (e.g., norethindrone tablets), unless contraindicated. Note: An exception to this requirement can be made if the patient has previously used a gonadotropin-releasing hormone agonist (e.g., Lupron Depot [leuprolide depot suspension]) or Orilissa (elagolix tablets).
<b>Indications</b>	All FDA-approved Indications.
<b>Off-Label Uses</b>	N/A
<b>Part B Prerequisite</b>	No

# NAYZILAM

## Products Affected

- NAYZILAM

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	Diagnosis, other medications used at the same time
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	Prescribed by or in consultation with a neurologist
<b>Coverage Duration</b>	1 year
<b>Other Criteria</b>	Intermittent Episodes of Frequent Seizure Activity (i.e., seizure clusters, acute repetitive seizures)-approve if the patient is currently receiving maintenance antiepileptic medication(s).
<b>Indications</b>	All FDA-approved Indications.
<b>Off-Label Uses</b>	N/A
<b>Part B Prerequisite</b>	No

# NERLYNX

## Products Affected

- NERLYNX

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	Stage of cancer, HER2 status, previous or current medications tried
<b>Age Restrictions</b>	18 years and older
<b>Prescriber Restrictions</b>	N/A
<b>Coverage Duration</b>	Adjuvant tx-Approve for 1 year (total), advanced or metastatic disease-1 year
<b>Other Criteria</b>	Breast cancer, adjuvant therapy - approve if the patient meets all of the following criteria: patient will not be using this medication in combination with HER2 antagonists, patient has HER2-positive breast cancer AND Patient has completed one year of adjuvant therapy with trastuzumab OR could not tolerate one year of therapy. Breast cancer, recurrent or metastatic disease-approve if the patient has HER-2 positive breast cancer, Nerlynx will be used in combination with capecitabine and the patient has tried at least two prior anti-HER2 based regimens.
<b>Indications</b>	All FDA-approved Indications.
<b>Off-Label Uses</b>	N/A
<b>Part B Prerequisite</b>	No



# NEXLETOL

## Products Affected

- NEXLETOL

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	LDL-C and response to other agents, prior therapies tried, medication adverse event history, medical history (as specified in the Other Criteria field)
<b>Age Restrictions</b>	18 years and older
<b>Prescriber Restrictions</b>	N/A
<b>Coverage Duration</b>	Authorization will be for 1 year
<b>Other Criteria</b>	<p>HYPERLIPIDEMIA WITH HeFH (both A and B): A) meets (a, b, c, or d): a) untreated LDL-C greater than or equal to 190 mg/dL, b) phenotypic confirmation (Note 1) of HeFH, c) Dutch Lipid Network criteria score greater than 5, or d) Simon Broome criteria met threshold for definite or possible/probable, AND B) meets (a or b): a) tried one high-intensity statin (throughout, see Definition 1 below) and ezetimibe and LDL-C remains 70 mg/dL or higher or b) statin intolerant (throughout, see Definition 2 below). ESTABLISHED CVD (both A and B): A) patient has/had one of the following conditions: prior MI, ACS, angina, CVA or TIA, CAD, PAD, coronary or other arterial revascularization procedure, B) meets (a or b): a) tried one high-intensity statin and ezetimibe and LDL-C remains 55 mg/dL or higher or b) statin intolerant. PRIMARY HYPERLIPIDEMIA (not associated with established CVD or HeFH) [A or B]: A) tried one high-intensity statin and ezetimibe for 8 weeks or longer and LDL-C remains 70 mg/dL or higher or B) statin intolerant. Note 1: Examples include mutations at the low-density lipoprotein receptor (LDLR), apolipoprotein B (apo B), proprotein convertase subtilisin kexin type 9 (PCSK) or low-density lipoprotein receptor adaptor protein (LDLRAP1) gene. Definition 1: High intensity statin defined as atorvastatin greater than or equal to 40 mg daily or rosuvastatin greater than or equal to 20 mg daily. Definition 2:</p>

PA Criteria	Criteria Details
	Statin intolerance defined as experiencing statin related rhabdomyolysis or skeletal-related muscle symptoms while receiving separate trials of atorvastatin and rosuvastatin and during both trials the skeletal-related symptoms resolved upon discontinuation of the statin.
<b>Indications</b>	All FDA-approved Indications.
<b>Off-Label Uses</b>	N/A
<b>Part B Prerequisite</b>	No

# NEXLIZET

## Products Affected

- NEXLIZET

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	LDL-C and response to other agents, prior therapies tried, medication adverse event history, medical history (as specified in the Other Criteria field)
<b>Age Restrictions</b>	18 years and older
<b>Prescriber Restrictions</b>	N/A
<b>Coverage Duration</b>	Authorization will be for 1 year
<b>Other Criteria</b>	<p>HYPERLIPIDEMIA WITH HeFH (both A and B): A) meets (a, b, c, or d): a) untreated LDL-C greater than or equal to 190 mg/dL, b) phenotypic confirmation (Note 1) of HeFH, c) Dutch Lipid Network criteria score greater than 5, or d) Simon Broome criteria met threshold for definite or possible/probable, AND B) meets (a or b): a) tried one high-intensity statin (throughout, see Definition 1 below) and LDL-C remains 70 mg/dL or higher or b) statin intolerant (throughout, see Definition 2 below).</p> <p>ESTABLISHED CVD (both A and B): A) patient has/had one of the following conditions: prior MI, ACS, angina, CVA or TIA, CAD, PAD, coronary or other arterial revascularization procedure, B) meets (a or b): a) tried one high-intensity statin and LDL-C remains 55 mg/dL or higher or b) statin intolerant.</p> <p>PRIMARY HYPERLIPIDEMIA (not associated with established CVD or HeFH) [A or B]: A) tried one high-intensity statin for 8 weeks or longer and LDL-C remains 70 mg/dL or higher or B) statin intolerant.</p> <p>Note 1: Examples include mutations at the low-density lipoprotein receptor (LDLR), apolipoprotein B (apo B), proprotein convertase subtilisin kexin type 9 (PCSK) or low-density lipoprotein receptor adaptor protein (LDLRAP1) gene. Definition 1: High intensity statin defined as atorvastatin greater than or equal to 40 mg daily or rosuvastatin greater than or equal to 20 mg daily. Definition 2: Statin</p>

PA Criteria	Criteria Details
	intolerance defined as experiencing statin related rhabdomyolysis or skeletal-related muscle symptoms while receiving separate trials of atorvastatin and rosuvastatin and during both trials the skeletal-related symptoms resolved upon discontinuation of the statin.
<b>Indications</b>	All FDA-approved Indications.
<b>Off-Label Uses</b>	N/A
<b>Part B Prerequisite</b>	No

# NILUTAMIDE

## Products Affected

- *nilutamide*

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	Diagnosis
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	1 year
Other Criteria	Prostate cancer-approve if nilutamide is used concurrently with a luteinizing hormone-releasing hormone (LHRH) agonist.
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

# NINLARO

---

## Products Affected

- NINLARO

PA Criteria	Criteria Details
Exclusion Criteria	PENDING CMS REVIEW
Required Medical Information	PENDING CMS REVIEW
Age Restrictions	PENDING CMS REVIEW
Prescriber Restrictions	PENDING CMS REVIEW
Coverage Duration	PENDING CMS REVIEW
Other Criteria	PENDING CMS REVIEW
Indications	PENDING CMS REVIEW
Off-Label Uses	PENDING CMS REVIEW
Part B Prerequisite	PENDING CMS REVIEW

# NITISINONE

## Products Affected

- *nitisinone*

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	Concomitant therapy with nitisinone products
<b>Required Medical Information</b>	Diagnosis, genetic tests and lab results (as specified in the Other Criteria field)
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	Prescribed by or in consultation with a metabolic disease specialist (or specialist who focuses in the treatment of metabolic diseases)
<b>Coverage Duration</b>	1 year
<b>Other Criteria</b>	Hereditary Tyrosinemia, Type 1-approve if the prescriber confirms the diagnosis was confirmed by genetic testing confirming biallelic pathogenic/likely pathogenic variants in the FAH gene OR elevated levels of succinylacetone in the serum or urine.
<b>Indications</b>	All FDA-approved Indications.
<b>Off-Label Uses</b>	N/A
<b>Part B Prerequisite</b>	No

# NIVESTYM

## Products Affected

- NIVESTYM

PA Criteria	Criteria Details
Exclusion Criteria	PENDING CMS REVIEW
Required Medical Information	PENDING CMS REVIEW
Age Restrictions	PENDING CMS REVIEW
Prescriber Restrictions	PENDING CMS REVIEW
Coverage Duration	PENDING CMS REVIEW
Other Criteria	PENDING CMS REVIEW
Indications	PENDING CMS REVIEW
Off-Label Uses	PENDING CMS REVIEW
Part B Prerequisite	PENDING CMS REVIEW



## NON-INJECTABLE TESTOSTERONE PRODUCTS

### Products Affected

- testosterone transdermal gel in metered-dose pump 12.5 mg/ 1.25 gram (1 %), 20.25 mg/1.25 gram (1.62 %)      1.62 % (20.25 mg/1.25 gram), 1.62 % (40.5 mg/2.5 gram)
- testosterone transdermal gel in packet 1 % (25 mg/2.5gram), 1 % (50 mg/5 gram),      • testosterone transdermal solution in metered pump w/app

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	Diagnosis of primary hypogonadism (congenital or acquired) in males. Diagnosis of secondary (hypogonadotropic) hypogonadism (congenital or acquired) in males. Hypogonadism (primary or secondary) in males, serum testosterone level. [Man is defined as an individual with the biological traits of a man, regardless of the individual's gender identity or gender expression.]
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	Gender-Dysphoric/Gender-Incongruent Persons, Persons Undergoing Female-to-Male (FTM) Gender Reassignment (i.e., Endocrinologic Masculinization)-prescribed by or in consultation with an endocrinologist or a physician who specializes in the treatment of transgender patients.
<b>Coverage Duration</b>	Authorization will be for 12 months.
<b>Other Criteria</b>	Hypogonadism (primary or secondary) in males - initial therapy, approve if all of the following criteria are met: 1) patient has persistent signs and symptoms of androgen deficiency (pre-treatment) [eg, depressed mood, decreased energy, progressive decrease in muscle mass, osteoporosis, loss of libido, AND 2) patient has had two pre-treatment serum testosterone (total or available) measurements, each taken in the morning on two separate days, AND 3) the two serum testosterone levels are both low, as defined by the normal laboratory reference values. Hypogonadism (primary or secondary) in males - continuing therapy, approve if the patient meets all of the following criteria: 1) patient has persistent signs and symptoms of androgen deficiency (pre-treatment) AND 2) patient had at least one pre-treatment serum testosterone level that was low. [Note: male is defined as

PA Criteria	Criteria Details
	<p>an individual with the biological traits of a man, regardless of the individual's gender identity or gender expression.] Gender-Dysphoric/Gender-Incongruent Persons, Persons Undergoing Female-to-Male (FTM) Gender Reassignment (i.e., Endocrinologic Masculinization)-approve.Note: For a patient who has undergone gender reassignment, use this FTM criterion for hypogonadism indication.</p>
<b>Indications</b>	All FDA-approved Indications, Some Medically-accepted Indications.
<b>Off-Label Uses</b>	Gender-Dysphoric/Gender-Incongruent Persons, Persons Undergoing Female-to-Male (FTM) Gender Reassignment (i.e., Endocrinologic Masculinization)
<b>Part B Prerequisite</b>	No

# NUBEQA

## Products Affected

- NUBEQA

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	Diagnosis
<b>Age Restrictions</b>	18 years and older
<b>Prescriber Restrictions</b>	N/A
<b>Coverage Duration</b>	1 year
<b>Other Criteria</b>	Prostate cancer - non-metastatic, castration resistant-approve if the requested medication will be used concurrently with a gonadotropin-releasing hormone (GnRH) agonist or if the patient has had a bilateral orchiectomy or if the medication is used concurrently with Firmagon. Prostate cancer-metastatic, castration sensitive-approve if (A and B): A) the medication is used in combination with docetaxel or patient has completed docetaxel therapy, and B) the medication will be used in combination with a GnRH agonist or in combination with Firmagon or if the patient had a bilateral orchiectomy.
<b>Indications</b>	All FDA-approved Indications.
<b>Off-Label Uses</b>	N/A
<b>Part B Prerequisite</b>	No

# NUCALA

## Products Affected

- NUCALA SUBCUTANEOUS AUTO-INJECTOR
- NUCALA SUBCUTANEOUS RECON SOLN
- NUCALA SUBCUTANEOUS SYRINGE 100 MG/ML, 40 MG/0.4 ML

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	Concurrent use with another monoclonal antibody therapy.
<b>Required Medical Information</b>	N/A
<b>Age Restrictions</b>	Asthma-6 years of age and older. EGPA/Polyps-18 years of age and older. HES-12 years and older.
<b>Prescriber Restrictions</b>	Asthma-Prescribed by or in consultation with an allergist, immunologist, or pulmonologist. EGPA-prescribed by or in consultation with an allergist, immunologist, pulmonologist or rheumatologist. HES-prescribed by or in consultation with an allergist, immunologist, hematologist, pulmonologist or rheumatologist. Polyps-prescribed by or in consult with allergist, immunologist or Otolaryngologist.
<b>Coverage Duration</b>	Initial-Asthma/EGPA/polyps-6 months, HES-8 months. 12 months continuation.
<b>Other Criteria</b>	Asthma initial - must have blood eosinophil level of greater than or equal to 150 cells per microliter within the previous 6 wks (prior to tx with Nucala or another monoclonal antibody therapy that may lower blood eosinophil levels) AND has received combo tx w/inhaled corticosteroid AND at least 1 additional asthma controller/maintenance med (Examples: LAMA, LABA, leukotriene receptor antagonist, monoclonal antibody) AND pt's asthma cont to be uncontrolled, or was uncontrolled prior to starting Nucala or another monoclonal antibody therapy for asthma as defined by 1 of following-pt experi 2 or more asthma exacer req tx w/systemic corticosteroids in prev yr, pt experienced 1 or more asthma exacer requiring hospitalization, urgent care visit or ED visit in the prev yr, pt has a FEV1 less than 80 percent predicted, Pt has FEV1/FVC less than 0.80, or Pt's asthma worsens upon taper of oral (systemic) corticosteroid

PA Criteria	Criteria Details
	<p>therapy. Cont-pt responded to Nucala tx as determined by the prescribing physician AND Pt cont to receive tx with an inhaled corticosteroid. EGPA initial-approve if pt has active, non-severe disease, has/had a blood eosinophil level of greater than or equal to 150 cells per microliter within the previous 6 wks or within 6 wks prior to any monoclonal antibody that may lower blood eosinophil levels. Cont-pt responded to Nucala tx as determined by the prescribing physician.HES initial-pt has had hypereosinophilic synd for greater than or equal to 6 months AND has FIP1L1-PDGFRalpha-negative dis AND pt does NOT have identifiable non-hematologic secondary cause of hypereosinophilic synd AND prior to initiating tx with monoclonal antibody that may lower blood eosinophil levels, pt has/had a blood eosinophil level of greater than or equal to 1,000 cells per microliter. Cont-approve if the patient has responded to Nucala tx. Nasal polyps, initial-approve if pt meets ALL of the following criteria(A, B, C and D):A) pt has chronic rhinosinusitis w/nasal polyposis as evidenced by direct examination, endoscopy, or sinus CT scan AND B)pt experienced 2 or more of the following sympt for at least 6 months:nasal congest/obstruct/discharge, and/or reduction/loss of smell AND C)pt meets BOTH of the following (a and b): a)Pt has received tx with intranasal corticosteroid AND b)Pt will continue to receive tx with intranasal corticosteroid concomitantly with Nucala AND D)pt meets 1 of the following (a, b or c): a)Pt has received at least 1 course of tx with a systemic corticosteroid for 5 days or more within the previous 2 years, OR b)Pt has a contraindication to systemic corticosteroid tx, OR c)Pt had prior surgery for nasal polyps.Cont-approve if the pt has received at least 6 months of therapy, continues to receive tx with an intranasal corticosteroid and has responded to tx.</p>
<b>Indications</b>	All FDA-approved Indications.
<b>Off-Label Uses</b>	N/A
<b>Part B Prerequisite</b>	No

# NUEDEXTA

## Products Affected

- NUEDEXTA

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	N/A
Age Restrictions	N/A
Prescriber Restrictions	Prescribed by or in consultation with a neurologist
Coverage Duration	1 year
Other Criteria	N/A
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

# NUPLAZID

## Products Affected

- NUPLAZID

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	Diagnosis
Age Restrictions	N/A
Prescriber Restrictions	Prescribed by or in consultation with a neurologist
Coverage Duration	1 year
Other Criteria	N/A
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

# NURTEC

## Products Affected

- NURTEC ODT

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	Concurrent use with another calcitonin gene-related peptide (CGRP) inhibitor being prescribed for migraine headache prevention if Nurtec ODT is being taking for the preventive treatment of episodic migraine.
<b>Required Medical Information</b>	Diagnosis
<b>Age Restrictions</b>	18 years and older
<b>Prescriber Restrictions</b>	N/A
<b>Coverage Duration</b>	1 year
<b>Other Criteria</b>	Migraine, Acute treatment-approve if the patient has tried at least one triptan or has a contraindication to triptans. Preventive treatment of episodic migraine-approve if the patient has greater than or equal to 4 but less than 15 migraine headache days per month (prior to initiating a migraine preventive medication) and if the patient is currently taking Nurtec ODT, the patient has had a significant clinical benefit from the medication.
<b>Indications</b>	All FDA-approved Indications.
<b>Off-Label Uses</b>	N/A
<b>Part B Prerequisite</b>	No



# NYVEPRIA

## Products Affected

- NYVEPRIA

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	Diagnosis
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	Cancer patients receiving chemotherapy-prescribed by or in consultation with an oncologist or hematologist. PBPC-prescribed by or in consultation with an oncologist, hematologist, or physician that specializes in transplantation.
<b>Coverage Duration</b>	Cancer pts receiving chemo-6 mo. PBPC-1 mo
<b>Other Criteria</b>	Cancer patients receiving chemotherapy, approve if the patient meets one of the following: 1) is receiving myelosuppressive anti-cancer medications that are associated with a high risk of febrile neutropenia (the risk is at least 20% based on the chemotherapy regimen), 2) patient is receiving myelosuppressive anti-cancer medications that are associated with a risk of febrile neutropenia but the risk is less than 20% based on the chemotherapy regimen and the patient has one or more risk factors for febrile neutropenia according to the prescribing physician (eg, aged greater than or equal to 65 years, prior chemotherapy or radiation therapy, persistent neutropenia, bone marrow involvement by tumor, recent surgery and/or open wounds, liver and/or renal dysfunction, poor performance status or HIV infection, or 3) patient has had a neutropenic complication from prior chemotherapy and did not receive prophylaxis with a colony stimulating factor and a reduced dose or frequency of chemotherapy may compromise treatment.
<b>Indications</b>	All FDA-approved Indications, Some Medically-accepted Indications.
<b>Off-Label Uses</b>	Patients undergoing PBPC collection and therapy

PA Criteria	Criteria Details
Part B Prerequisite	No

# OCALIVA

## Products Affected

- OCALIVA

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	Prescriber specialty, lab values, prior medications used for diagnosis and length of trials
<b>Age Restrictions</b>	18 years and older (initial)
<b>Prescriber Restrictions</b>	Prescribed by or in consultation with a gastroenterologist, hepatologist, or liver transplant physician (initial)
<b>Coverage Duration</b>	6 months initial, 1 year cont.
<b>Other Criteria</b>	Initial treatment of PBC-Patient must meet both 1 and 2-1. Patient has a diagnosis of PBC as defined by TWO of the following:a)Alkaline phosphatase (ALP) elevated above the upper limit of normal as defined by normal laboratory reference values b)Positive anti-mitochondrial antibodies (AMAs) or other PBC-specific auto-antibodies, including sp100 or gp210, if AMA is negative c)Histologic evidence of primary biliary cholangitis (PBC) from a liver biopsy 2. Patient meets ONE of the following: a) Patient has been receiving ursodiol therapy for greater than or equal to 1 year and has had an inadequate response. b) Patient is unable to tolerate ursodiol therapy. Cont tx - approve if the patient has responded to Ocaliva therapy as determined by the prescribing physician (e.g., improved biochemical markers of PBC (e.g., alkaline phosphatase [ALP], bilirubin, gamma-glutamyl transpeptidase [GGT], aspartate aminotransferase [AST], alanine aminotransferase [ALT] levels)). Patients new to therapy and continuing therapy must not have cirrhosis or must have compensated cirrhosis without evidence of portal hypertension.
<b>Indications</b>	All FDA-approved Indications.
<b>Off-Label Uses</b>	N/A

PA Criteria	Criteria Details
Part B Prerequisite	No

# OCTREOTIDE INJECTABLE

## Products Affected

- *octreotide acetate injection solution*

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	Diagnosis
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	Acromegaly-prescr/consult w/endocrinologist. NETs-prescr/consult w/oncologist, endocrinologist, or gastroenterologist. Pheochromocytoma/paraganglioma-prescr/consult w/endo/onc/neuro.Meningioma-prescr/consult w/oncologist, radiologist, neurosurg/thymoma/thymic carcinoma-prescr/consult with oncologist
<b>Coverage Duration</b>	1 year
<b>Other Criteria</b>	ACROMEGALY (A or B): A) inadequate response to surgery and/or radiotherapy or patient not an appropriate candidate or B) patient is experiencing negative effects due to tumor size (e.g., optic nerve compression) and has pre-treatment (baseline) insulin-like growth factor-1 (IGF-1) level above the upper limit of normal based on age and gender.
<b>Indications</b>	All FDA-approved Indications, Some Medically-accepted Indications.
<b>Off-Label Uses</b>	Meningioma, thymoma and thymic carcinoma, pheochromocytoma and paraganglioma
<b>Part B Prerequisite</b>	No

# ODOMZO

## Products Affected

- ODOMZO

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	BCC - Must not have had disease progression while on Erivedge (vismodegib).
<b>Age Restrictions</b>	18 years and older
<b>Prescriber Restrictions</b>	N/A
<b>Coverage Duration</b>	1 year
<b>Other Criteria</b>	Locally advanced BCC approve if the BCC has recurred following surgery/radiation therapy or if the patient is not a candidate for surgery AND the patient is not a candidate for radiation therapy, according to the prescribing physician. Metastatic BCC - approve, if the disease is limited to nodal metastases. (Note-This includes primary or recurrent nodal metastases. A patient with distant metastasis does not meet this requirement.)
<b>Indications</b>	All FDA-approved Indications, Some Medically-accepted Indications.
<b>Off-Label Uses</b>	Metastatic BCC
<b>Part B Prerequisite</b>	No

## OFEV

### Products Affected

- OFEV

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	N/A
<b>Age Restrictions</b>	18 years of age and older
<b>Prescriber Restrictions</b>	IPF-Prescribed by or in consultation with a pulmonologist. Interstitial lung disease associated with systemic sclerosis-prescribed by or in consultation with a pulmonologist or rheumatologist.
<b>Coverage Duration</b>	1 year
<b>Other Criteria</b>	IDIOPATHIC PULMONARY FIBROSIS (IPF) [A and B]: A) diagnosis confirmed by presence of usual interstitial pneumonia (UIP) pattern on high-resolution computed tomography (HRCT) or surgical lung biopsy and B) forced vital capacity (FVC) greater than or equal to 40 percent of the predicted value. INTERSTITIAL LUNG DISEASE ASSOCIATED WITH SYSTEMIC SCLEROSIS (A and B): A) diagnosis confirmed by HRCT and B) FVC greater than or equal to 40 percent of the predicted value. CHRONIC FIBROSING INTERSTITIAL LUNG DISEASE (all of A, B and C): A) FVC greater than or equal to 45 percent of the predicted value, B) fibrosing lung disease impacting more than 10 percent of lung volume on HRCT, and C) clinical signs of progression.
<b>Indications</b>	All FDA-approved Indications.
<b>Off-Label Uses</b>	N/A
<b>Part B Prerequisite</b>	No

# OGSIVEO

---

## Products Affected

- OGSIVEO ORAL TABLET 100 MG, 150 MG, 50 MG

PA Criteria	Criteria Details
Exclusion Criteria	PENDING CMS REVIEW
Required Medical Information	PENDING CMS REVIEW
Age Restrictions	PENDING CMS REVIEW
Prescriber Restrictions	PENDING CMS REVIEW
Coverage Duration	PENDING CMS REVIEW
Other Criteria	PENDING CMS REVIEW
Indications	PENDING CMS REVIEW
Off-Label Uses	PENDING CMS REVIEW
Part B Prerequisite	PENDING CMS REVIEW



## OJEMDA

### Products Affected

- OJEMDA ORAL SUSPENSION FOR RECONSTITUTION MG/WEEK (100 MG X 5), 600 MG/WEEK (100 MG X 6)
- OJEMDA ORAL TABLET 400 MG/WEEK (100 MG X 4), 500

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	Diagnosis
Age Restrictions	6 months of age and older
Prescriber Restrictions	N/A
Coverage Duration	1 year
Other Criteria	PEDIATRIC LOW GRADE GLIOMA-patient has relapsed or refractory disease and the tumor is positive for one of the following: BRAF fusion, BRAF rearrangement or BRAF V600 mutation.
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

# OJJAARA

## Products Affected

- OJJAARA

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	Diagnosis
<b>Age Restrictions</b>	18 years of age and older
<b>Prescriber Restrictions</b>	N/A
<b>Coverage Duration</b>	1 year
<b>Other Criteria</b>	Myelofibrosis-approve if the patient has intermediate-risk or high-risk disease and (a or b): a) the patient has anemia, defined as hemoglobin less than 10g/dL and has symptomatic splenomegaly and/or constitutional symptoms, or b) the patient has platelet count greater than or equal to 50x10 <sup>9</sup> /L.
<b>Indications</b>	All FDA-approved Indications.
<b>Off-Label Uses</b>	N/A
<b>Part B Prerequisite</b>	No

# ONUREG

## Products Affected

- ONUREG

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	Diagnosis
Age Restrictions	18 years and older
Prescriber Restrictions	N/A
Coverage Duration	1 year
Other Criteria	AML - Approve if the medication is used for post-remission maintenance therapy AND the patient has intermediate or poor-risk cytogenetics AND allogeneic hematopoietic stem cell transplant is not planned.
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

# OPSUMIT

## Products Affected

- OPSUMIT

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	PAH WHO group, right heart catheterization results
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	PAH - must be prescribed by or in consultation with a cardiologist or a pulmonologist.
<b>Coverage Duration</b>	Authorization will be for 1 year
<b>Other Criteria</b>	Pulmonary arterial hypertension (PAH) WHO Group 1 patients are required to have had a right-heart catheterization to confirm the diagnosis of PAH to ensure appropriate medical assessment.
<b>Indications</b>	All FDA-approved Indications.
<b>Off-Label Uses</b>	N/A
<b>Part B Prerequisite</b>	No

# OPSYNVI

## Products Affected

- OPSYNVI

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	Concurrent use with guanylate cyclase stimulators
<b>Required Medical Information</b>	Diagnosis
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	Prescribed by or in consultation with a cardiologist or a pulmonologist (initial/continuation)
<b>Coverage Duration</b>	1 year
<b>Other Criteria</b>	Pulmonary arterial hypertension (PAH) WHO Group 1-approve if patient has had a right-heart catheterization to confirm the diagnosis.
<b>Indications</b>	All FDA-approved Indications.
<b>Off-Label Uses</b>	N/A
<b>Part B Prerequisite</b>	No

# ORENCIA

## Products Affected

- ORENCIA CLICKJECT
- ORENCIA SUBCUTANEOUS SYRINGE 125 MG/ML, 50 MG/0.4 ML, 87.5 MG/0.7 ML

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	Concurrent use with a Biologic DMARD or Targeted Synthetic DMARD.
<b>Required Medical Information</b>	Diagnosis, concurrent medications, previous drugs tried.
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	Initial therapy only-RA and JIA/JRA prescribed by or in consultation with a rheumatologist. PsA-prescribed by or in consultation with a rheumatologist or dermatologist.
<b>Coverage Duration</b>	Approve through end of plan year
<b>Other Criteria</b>	RA initial, patient has tried one conventional synthetic DMARD for at least 3 months (note: patients who have already had a 3-month trial of a biologic for RA are not required to step back and try a conventional synthetic DMARD). PsA-approve. Juvenile idiopathic arthritis (JIA) [or Juvenile Rheumatoid Arthritis (JRA)]-approve if the patient has tried one other agent for this condition or the patient will be starting on Orencia concurrently with methotrexate, sulfasalazine or leflunomide or the patient has an absolute contraindication to methotrexate, sulfasalazine or leflunomide or the patient has aggressive disease. Cont tx - responded to therapy.
<b>Indications</b>	All FDA-approved Indications.
<b>Off-Label Uses</b>	N/A
<b>Part B Prerequisite</b>	No

# ORGOVYX

## Products Affected

- ORGOVYX

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	Diagnosis
Age Restrictions	18 years and older
Prescriber Restrictions	N/A
Coverage Duration	1 year
Other Criteria	Prostate Cancer-approve.
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

# ORKAMBI

## Products Affected

- ORKAMBI ORAL GRANULES IN PACKET
- ORKAMBI ORAL TABLET

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	Combination use with Kalydeco, Trikafta or Symdeko.
<b>Required Medical Information</b>	N/A
<b>Age Restrictions</b>	1 year of age and older
<b>Prescriber Restrictions</b>	prescribed by or in consultation with a pulmonologist or a physician who specializes in CF
<b>Coverage Duration</b>	1 year
<b>Other Criteria</b>	CF - Approve if the pt meets A, B and C: A) pt has two copies of the F508del mutation in the CFTR gene, and B) pt must have positive CF newborn screening test or family history of CF or clinical presentation consistent with signs and symptoms of CF and C) evidence of abnormal CFTR function as demonstrated by i, ii, or iii: (i) elevated sweat chloride test or (ii) two CFTR mutations or (iii) abnormal nasal potential difference.
<b>Indications</b>	All FDA-approved Indications.
<b>Off-Label Uses</b>	N/A
<b>Part B Prerequisite</b>	No



# ORSERDU

## Products Affected

- ORSERDU ORAL TABLET 345 MG, 86 MG

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	Diagnosis
<b>Age Restrictions</b>	18 years and older
<b>Prescriber Restrictions</b>	N/A
<b>Coverage Duration</b>	1 year
<b>Other Criteria</b>	Breast cancer in postmenopausal women or Men-approve if the patient meets the following criteria (A, B, C, D, and E): A) Patient has recurrent or metastatic disease, AND B) Patient has estrogen receptor positive (ER+) disease, AND C) Patient has human epidermal growth factor receptor 2 (HER2)-negative disease, AND D) Patient has estrogen receptor 1 gene (ESR1)-mutated disease, AND E) Patient has tried at least one endocrine therapy. Note: Examples of endocrine therapy include fulvestrant, anastrozole, exemestane, letrozole, and tamoxifen.
<b>Indications</b>	All FDA-approved Indications.
<b>Off-Label Uses</b>	N/A
<b>Part B Prerequisite</b>	No

# OTEZLA

## Products Affected

- OTEZLA MG (51), 10 MG (4)-20 MG (4)-30 MG
- OTEZLA STARTER ORAL (47)
- TABLETS,DOSE PACK 10 MG (4)- 20

PA Criteria	Criteria Details
Exclusion Criteria	PENDING CMS REVIEW
Required Medical Information	PENDING CMS REVIEW
Age Restrictions	PENDING CMS REVIEW
Prescriber Restrictions	PENDING CMS REVIEW
Coverage Duration	PENDING CMS REVIEW
Other Criteria	PENDING CMS REVIEW
Indications	PENDING CMS REVIEW
Off-Label Uses	PENDING CMS REVIEW
Part B Prerequisite	PENDING CMS REVIEW

# OXERVATE

## Products Affected

- OXERVATE

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	Treatment duration greater than 16 weeks per affected eye(s)
<b>Required Medical Information</b>	Diagnosis
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	Prescribed by an ophthalmologist or an optometrist.
<b>Coverage Duration</b>	Initial-8 weeks, continuation-approve for an additional 8 weeks
<b>Other Criteria</b>	Patients who have already received Oxervate-approve if the patient has previously received less than or equal to 8 weeks of treatment per affected eye(s) and the patient has a recurrence of neurotrophic keratitis.
<b>Indications</b>	All FDA-approved Indications.
<b>Off-Label Uses</b>	N/A
<b>Part B Prerequisite</b>	No

# PANRETIN

## Products Affected

- PANRETIN

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	Diagnosis
Age Restrictions	N/A
Prescriber Restrictions	Prescribed by or in consultation with a dermatologist, oncologist, or infectious disease specialist
Coverage Duration	1 year
Other Criteria	Kaposi Sarcoma-approve if the patient is not receiving systemic therapy for Kaposi Sarcoma.
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

# PEMAZYRE

## Products Affected

- PEMAZYRE

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	Diagnosis, prior therapies
<b>Age Restrictions</b>	18 years and older
<b>Prescriber Restrictions</b>	N/A
<b>Coverage Duration</b>	Authorization will be for 1 year
<b>Other Criteria</b>	Cholangiocarcinoma-approve if the patient has unresectable locally advanced or metastatic disease and the tumor has a fibroblast growth factor receptor 2 (FGFR2) fusion or other rearrangement, as detected by an approved test AND the patient has been previously treated with at least one systemic therapy regimen. Myeloid/lymphoid neoplasms-approve if the patient has eosinophilia and the cancer has fibroblast growth factor receptor 1 (FGFR1) rearrangement, as detected by an approved test and the cancer is in chronic phase or blast phase.
<b>Indications</b>	All FDA-approved Indications.
<b>Off-Label Uses</b>	N/A
<b>Part B Prerequisite</b>	No

# PENICILLAMINE

## Products Affected

- *penicillamine oral tablet*

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	Diagnosis
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	Wilson's Disease-Prescribed by or in consultation with a gastroenterologist, hepatologist or liver transplant physician
<b>Coverage Duration</b>	1 year
<b>Other Criteria</b>	Cystinuria-approve. Wilson's disease-approve if diagnosis is confirmed by ONE of the following (i or ii): i. Genetic testing results confirming biallelic pathogenic ATP7B mutations (in either symptomatic or asymptomatic individuals), OR ii. Confirmation of at least two of the following (a, b, c, d): a) Presence of Kayser-Fleischer rings, b) Serum ceruloplasmin level less than 20 mg/dL, c) Liver biopsy findings consistent with Wilson's disease, d) 24-hour urinary copper greater than 40 mcg/24 hours.
<b>Indications</b>	All FDA-approved Indications.
<b>Off-Label Uses</b>	N/A
<b>Part B Prerequisite</b>	No

# PHENYL BUTYRATE

## Products Affected

- sodium phenylbutyrate*

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	Concomitant therapy with more than one phenylbutyrate product
<b>Required Medical Information</b>	Diagnosis, genetic tests and lab results (as specified in the Other Criteria field)
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	Prescribed by or in consultation with a metabolic disease specialist (or specialist who focuses in the treatment of metabolic diseases)
<b>Coverage Duration</b>	Pt meets criteria with no genetic test - 3 mo approval. Pt had genetic test - 12 mo approval
<b>Other Criteria</b>	Urea cycle disorders-approve if genetic testing confirmed a mutation resulting in a urea cycle disorder or if the patient has hyperammonemia diagnosed with an ammonia level above the upper limit of the normal reference range for the reporting laboratory.
<b>Indications</b>	All FDA-approved Indications.
<b>Off-Label Uses</b>	N/A
<b>Part B Prerequisite</b>	No

# PHEOCHROMOCYTOMA

## Products Affected

- *metyrosine*

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	Diagnosis, prior medication trials
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	Prescribed by or in consultation with an endocrinologist or a physician who specializes in the management of pheochromocytoma (initial and continuation therapy for metyrosine)
<b>Coverage Duration</b>	Authorization will be for 1 year
<b>Other Criteria</b>	If the requested drug is metyrosine for initial therapy, approve if the patient has tried a selective alpha blocker (e.g., doxazosin, terazosin or prazosin). If the requested drug is metyrosine for continuation therapy, approve if the patient is currently receiving metyrosine or has received metyrosine in the past.
<b>Indications</b>	All FDA-approved Indications.
<b>Off-Label Uses</b>	N/A
<b>Part B Prerequisite</b>	No



## PHOSPHODIESTERASE-5 INHIBITORS FOR PAH

### Products Affected

- *alyq*
- *sildenafil (pulmonary arterial hypertension) oral tablet*
- *tadalafil (pulmonary arterial hypertension) oral tablet 20 mg*

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	Concurrent Use With Guanylate Cyclase Stimulators.
<b>Required Medical Information</b>	Diagnosis, right heart cath results
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	For PAH, if prescribed by, or in consultation with, a cardiologist or a pulmonologist.
<b>Coverage Duration</b>	Authorization will be for 1 year.
<b>Other Criteria</b>	Pulmonary arterial hypertension (PAH) WHO Group 1, are required to have had a right-heart catheterization to confirm diagnosis of PAH to ensure appropriate medical assessment. Clinical criteria incorporated into the quantity limit edits for sildenafil 20 mg tablets require confirmation that the indication is PAH (ie, FDA labeled use) prior to reviewing for quantity exception.
<b>Indications</b>	All FDA-approved Indications.
<b>Off-Label Uses</b>	N/A
<b>Part B Prerequisite</b>	No

# PIQRAY

## Products Affected

- PIQRAY ORAL TABLET 200 MG/DAY (200 MG X 1), 250 MG/DAY (200 MG X 1), 300 MG/DAY (150 MG X 2)

PA Criteria	Criteria Details
Exclusion Criteria	PENDING CMS REVIEW
Required Medical Information	PENDING CMS REVIEW
Age Restrictions	PENDING CMS REVIEW
Prescriber Restrictions	PENDING CMS REVIEW
Coverage Duration	PENDING CMS REVIEW
Other Criteria	PENDING CMS REVIEW
Indications	PENDING CMS REVIEW
Off-Label Uses	PENDING CMS REVIEW
Part B Prerequisite	PENDING CMS REVIEW

# PIRFENIDONE

## Products Affected

- *pirfenidone oral capsule*
- *pirfenidone oral tablet 267 mg, 801 mg*

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	N/A
Age Restrictions	18 years of age and older
Prescriber Restrictions	Prescribed by or in consultation with a pulmonologist
Coverage Duration	1 year
Other Criteria	IPF - must have FVC greater than or equal to 40 percent of the predicted value AND IPF must be diagnosed with either findings on high-resolution computed tomography (HRCT) indicating usual interstitial pneumonia (UIP) or surgical lung biopsy demonstrating UIP.
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

# PLEGRIDY

## Products Affected

- PLEGRIDY SUBCUTANEOUS PEN INJECTOR 125 MCG/0.5 ML
- PLEGRIDY SUBCUTANEOUS SYRINGE 125 MCG/0.5 ML

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	Concurrent use of with other disease-modifying agents used for multiple sclerosis (MS).
<b>Required Medical Information</b>	Relapsing form of Multiple Sclerosis (MS), to include clinically-isolated syndrome, relapsing-remitting disease, and active secondary progressive disease
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	Prescribed by, or in consultation with, a neurologist or an MS specialist.
<b>Coverage Duration</b>	Authorization will be for 1 year
<b>Other Criteria</b>	N/A
<b>Indications</b>	All FDA-approved Indications.
<b>Off-Label Uses</b>	N/A
<b>Part B Prerequisite</b>	No

# POMALYST

## Products Affected

- POMALYST

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	Diagnosis
<b>Age Restrictions</b>	Kaposi Sarcoma/MM-18 years and older
<b>Prescriber Restrictions</b>	N/A
<b>Coverage Duration</b>	Authorization will be for 1 year
<b>Other Criteria</b>	CNS Lymphoma-approve if the patient has relapsed or refractory disease. Kaposi Sarcoma-Approve if the patient meets one of the following (i or ii): i. patient is Human Immunodeficiency Virus (HIV)-negative OR ii. patient meets both of the following (a and b): a) The patient is Human Immunodeficiency Virus (HIV)-positive AND b) The patient continues to receive highly active antiretroviral therapy (HAART). MM-approve if the patient has received at least one other Revlimid (lenalidomide tablets)-containing regimen.
<b>Indications</b>	All FDA-approved Indications, Some Medically-accepted Indications.
<b>Off-Label Uses</b>	Systemic Light Chain Amyloidosis, Central Nervous System (CNS) Lymphoma
<b>Part B Prerequisite</b>	No

## POSACONAZOLE (ORAL)

### Products Affected

- *posaconazole oral tablet, delayed release (dr/ec)*

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	Diagnosis
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	N/A
<b>Coverage Duration</b>	Aspergillus/Candida prophylaxis, mucormycosis-6 mo, all others-3 months
<b>Other Criteria</b>	N/A
<b>Indications</b>	All FDA-approved Indications, Some Medically-accepted Indications.
<b>Off-Label Uses</b>	mucomycosis - maintenance, fusariosis, invasive - treatment fungal infections (systemic) in patients with human immunodeficiency virus (HIV) infections (e.g., histoplasmosis, coccidioidomycosis) - treatment.
<b>Part B Prerequisite</b>	No

# PREVYMIS

---

## Products Affected

- PREVYMIS ORAL

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	Diagnosis
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	6 months
Other Criteria	N/A
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

# PROLIA

## Products Affected

- PROLIA

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	Concomitant use with other medications for osteoporosis
<b>Required Medical Information</b>	N/A
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	N/A
<b>Coverage Duration</b>	Authorization will be for 1 year.
<b>Other Criteria</b>	Treatment of postmenopausal osteoporosis/Treatment of osteoporosis in men (to increase bone mass) [a man is defined as an individual with the biological traits of a man, regardless of the individual's gender identity or gender expression], approve if the patient meets one of the following: 1. has had inadequate response after 12 months of therapy with an oral bisphosphonate, had osteoporotic fracture or fragility fracture while receiving an oral bisphosphonate, or intolerability to an oral bisphosphonate, OR 2. the patient cannot take an oral bisphosphonate because they cannot swallow or have difficulty swallowing, they cannot remain in an upright position, or they have a pre-existing GI medical condition, OR 3. pt has tried an IV bisphosphonate (ibandronate or zoledronic acid), OR 4. the patient has severe renal impairment (eg, creatinine clearance less than 35 mL/min) or chronic kidney disease, or if the patient has an osteoporotic fracture or fragility fracture . Treatment of bone loss in patient at high risk for fracture receiving ADT for nonmetastatic prostate cancer, approve if the patient has prostate cancer that is not metastatic to the bone and the patient is receiving ADT (eg, leuprolide, triptorelin, goserelin) or the patient has undergone a bilateral orchiectomy. Treatment of bone loss (to increase bone mass) in patients at high risk for fracture receiving adjuvant AI therapy for breast cancer,



PA Criteria	Criteria Details
	<p>approve if the patient has breast cancer that is not metastatic to the bone and in receiving concurrent AI therapy (eg, anastrozole, letrozole, exemestane). Treatment of GIO, approve if pt tried one oral bisphosphonate OR pt cannot take an oral bisphosphonate because the patient cannot swallow or has difficulty swallowing or the patient cannot remain in an upright position post oral bisphosphonate administration or has a pre-existing GI medical condition (eg, patient with esophageal lesions, esophageal ulcers, or abnormalities of the esophagus that delay esophageal emptying [stricture, achalasia]), OR pt has tried zoledronic acid (Reclast), OR pt has severe renal impairment (CrCL less than 35 mL/min) or has CKD or has had an osteoporotic fracture or fragility fracture.</p>
<b>Indications</b>	All FDA-approved Indications.
<b>Off-Label Uses</b>	N/A
<b>Part B Prerequisite</b>	No

# PROMACTA

---

## Products Affected

- PROMACTA

PA Criteria	Criteria Details
Exclusion Criteria	PENDING CMS REVIEW
Required Medical Information	PENDING CMS REVIEW
Age Restrictions	PENDING CMS REVIEW
Prescriber Restrictions	PENDING CMS REVIEW
Coverage Duration	PENDING CMS REVIEW
Other Criteria	PENDING CMS REVIEW
Indications	PENDING CMS REVIEW
Off-Label Uses	PENDING CMS REVIEW
Part B Prerequisite	PENDING CMS REVIEW

# PYRIMETHAMINE

## Products Affected

- *pyrimethamine*

PA Criteria	Criteria Details
Exclusion Criteria	PENDING CMS REVIEW
Required Medical Information	PENDING CMS REVIEW
Age Restrictions	PENDING CMS REVIEW
Prescriber Restrictions	PENDING CMS REVIEW
Coverage Duration	PENDING CMS REVIEW
Other Criteria	PENDING CMS REVIEW
Indications	PENDING CMS REVIEW
Off-Label Uses	PENDING CMS REVIEW
Part B Prerequisite	PENDING CMS REVIEW

# QINLOCK

## Products Affected

- QINLOCK

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	Diagnosis, other therapies tried
<b>Age Restrictions</b>	18 years and older
<b>Prescriber Restrictions</b>	N/A
<b>Coverage Duration</b>	Authorization will be for 1 year
<b>Other Criteria</b>	Gastrointestinal stromal tumor (GIST)-approve if the patient has tried imatinib or avapritinib tablets, AND the patient meets one of the following criteria (i, ii, or iii): i. Patient has tried sunitinib and regorafenib tablets, OR ii. Patient has tried dasatinib tablets, OR iii. Patient is intolerant of sunitinib. Melanoma, cutaneous-approve if the patient has metastatic or unresectable disease, AND the patient has an activating KIT mutation, AND the patient has tried at least one systemic regimen.
<b>Indications</b>	All FDA-approved Indications, Some Medically-accepted Indications.
<b>Off-Label Uses</b>	Melanoma, cutaneous
<b>Part B Prerequisite</b>	No

# QULIPTA

## Products Affected

- QULIPTA

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	Concurrent use with another calcitonin gene-related peptide (CGRP) inhibitor being prescribed for migraine headache prevention
<b>Required Medical Information</b>	Diagnosis
<b>Age Restrictions</b>	18 years and older
<b>Prescriber Restrictions</b>	N/A
<b>Coverage Duration</b>	1 year
<b>Other Criteria</b>	Preventive treatment of episodic migraine-approve if the patient meets (A and B): A) has greater than or equal to 4 and less than 15 migraine headache days per month (prior to initiating a migraine-preventative medication, and B) if the pt is currently taking Qulipta, the pt has had significant clinical benefit. Chronic migraine prevention-approve if the patient has greater than or equal to 15 migraine headache days per month (prior to initiating a migraine-preventative medication).
<b>Indications</b>	All FDA-approved Indications.
<b>Off-Label Uses</b>	N/A
<b>Part B Prerequisite</b>	No

## RADICAVA ORS

### Products Affected

- RADICAVA ORS STARTER KIT SUSP

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	ALSFRS-R score, FVC %, time elapsed since diagnosis.
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	Prescribed by or in consultation with a neurologist, a neuromuscular disease specialist, or a physician specializing in the treatment of ALS (initial and continuation).
<b>Coverage Duration</b>	Initial, 6 months. Continuation, 6 months
<b>Other Criteria</b>	ALS, initial therapy - approve if the patient meets ALL of the following criteria: 1. According to the prescribing physician, the patient has a definite or probable diagnosis of ALS, based on the application of the El Escorial or the revised Airlie house diagnostic criteria AND 2. Patient has a score of two points or more on each item of the ALS Functional Rating Scale - Revised (ALSFRS-R) [ie, has retained most or all activities of daily living], AND 3. Patient has a percent predicted FVC greater than or equal to 80% (ie, has normal respiratory function), AND 4. Patient has been diagnosed with ALS for less than or equal to 2 years. 5. Patient has received or is currently receiving riluzole tablets. Note-a trial of Tiglutik or Exservan would also count. ALS, continuation therapy - approve if, according to the prescribing physician, the patient continues to benefit from therapy.
<b>Indications</b>	All FDA-approved Indications.
<b>Off-Label Uses</b>	N/A
<b>Part B Prerequisite</b>	No

# RELEUKO

---

## Products Affected

- RELEUKO SUBCUTANEOUS

PA Criteria	Criteria Details
Exclusion Criteria	PENDING CMS REVIEW
Required Medical Information	PENDING CMS REVIEW
Age Restrictions	PENDING CMS REVIEW
Prescriber Restrictions	PENDING CMS REVIEW
Coverage Duration	PENDING CMS REVIEW
Other Criteria	PENDING CMS REVIEW
Indications	PENDING CMS REVIEW
Off-Label Uses	PENDING CMS REVIEW
Part B Prerequisite	PENDING CMS REVIEW

# REPATHA

## Products Affected

- REPATHA
- REPATHA PUSHTRONEX
- REPATHA SURECLICK

PA Criteria	Criteria Details
Exclusion Criteria	PENDING CMS REVIEW
Required Medical Information	PENDING CMS REVIEW
Age Restrictions	PENDING CMS REVIEW
Prescriber Restrictions	PENDING CMS REVIEW
Coverage Duration	PENDING CMS REVIEW
Other Criteria	PENDING CMS REVIEW
Indications	PENDING CMS REVIEW
Off-Label Uses	PENDING CMS REVIEW
Part B Prerequisite	PENDING CMS REVIEW



# RETEVMO

## Products Affected

- RETEVMO ORAL CAPSULE 40 MG, 80 MG
- RETEVMO ORAL TABLET 120 MG, 160 MG, 40 MG, 80 MG

PA Criteria	Criteria Details
Exclusion Criteria	PENDING CMS REVIEW
Required Medical Information	PENDING CMS REVIEW
Age Restrictions	PENDING CMS REVIEW
Prescriber Restrictions	PENDING CMS REVIEW
Coverage Duration	PENDING CMS REVIEW
Other Criteria	PENDING CMS REVIEW
Indications	PENDING CMS REVIEW
Off-Label Uses	PENDING CMS REVIEW
Part B Prerequisite	PENDING CMS REVIEW

# REZDIFFRA

---

## Products Affected

- REZDIFFRA

PA Criteria	Criteria Details
Exclusion Criteria	PENDING CMS REVIEW
Required Medical Information	PENDING CMS REVIEW
Age Restrictions	PENDING CMS REVIEW
Prescriber Restrictions	PENDING CMS REVIEW
Coverage Duration	PENDING CMS REVIEW
Other Criteria	PENDING CMS REVIEW
Indications	PENDING CMS REVIEW
Off-Label Uses	PENDING CMS REVIEW
Part B Prerequisite	PENDING CMS REVIEW

# REZLIDHIA

## Products Affected

- REZLIDHIA

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	Diagnosis
Age Restrictions	18 years and older
Prescriber Restrictions	N/A
Coverage Duration	1 year
Other Criteria	Acute myeloid leukemia-approve if the patient has relapsed or refractory disease and the patient has isocitrate dehydrogenase-1 (IDH1) mutation positive disease as detected by an approved test.
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

# REZUROCK

## Products Affected

- REZUROCK

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	Diagnosis
Age Restrictions	12 years and older
Prescriber Restrictions	N/A
Coverage Duration	1 year
Other Criteria	Graft-versus-host disease-approve if the patient has chronic graft-versus-host disease and has tried at least two conventional systemic treatments (e.g., ibrutinib, cyclosporine) for chronic graft-versus-host disease.
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

# RILUZOLE

## Products Affected

- *riluzole*

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	Diagnosis
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	Prescribed by or in consultation with a neurologist, a neuromuscular disease specialist, or a physician specializing in the treatment of ALS.
<b>Coverage Duration</b>	1 year
<b>Other Criteria</b>	N/A
<b>Indications</b>	All FDA-approved Indications.
<b>Off-Label Uses</b>	N/A
<b>Part B Prerequisite</b>	No

# RINVOQ

## Products Affected

- RINVOQ ORAL TABLET EXTENDED  
RELEASE 24 HR 15 MG, 30 MG, 45 MG

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	PENDING CMS REVIEW
<b>Required Medical Information</b>	PENDING CMS REVIEW
<b>Age Restrictions</b>	PENDING CMS REVIEW
<b>Prescriber Restrictions</b>	PENDING CMS REVIEW
<b>Coverage Duration</b>	PENDING CMS REVIEW
<b>Other Criteria</b>	PENDING CMS REVIEW
<b>Indications</b>	PENDING CMS REVIEW
<b>Off-Label Uses</b>	PENDING CMS REVIEW
<b>Part B Prerequisite</b>	PENDING CMS REVIEW

## ROFLUMILAST (ORAL)

### Products Affected

- roflumilast*

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	Chronic Obstructive Pulmonary Disease (COPD), medications tried.
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	N/A
<b>Coverage Duration</b>	Authorization will be for 1 year.
<b>Other Criteria</b>	COPD, approve in patients who meet all of the following conditions: Patients has severe COPD or very severe COPD, AND Patient has a history of exacerbations, AND Patient has tried a medication from two of the three following drug categories: long-acting beta2-agonist (LABA) [eg, salmeterol, indacaterol], long-acting muscarinic antagonist (LAMA) [eg, tiotropium], inhaled corticosteroid (eg, fluticasone).
<b>Indications</b>	All FDA-approved Indications.
<b>Off-Label Uses</b>	N/A
<b>Part B Prerequisite</b>	No

# ROZLYTREK

## Products Affected

- ROZLYTREK ORAL CAPSULE 100 MG, 200 MG
- ROZLYTREK ORAL PELLETS IN PACKET

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	Diagnosis
<b>Age Restrictions</b>	NSCLC-18 years and older, Solid Tumors-1 month and older, Pediatric Diffuse High-Grade Glioma-less than 18 years old
<b>Prescriber Restrictions</b>	N/A
<b>Coverage Duration</b>	1 year
<b>Other Criteria</b>	Solid Tumors-Approve if the patient's tumor is positive for neurotrophic receptor tyrosine kinase (NTRK) gene fusion AND the tumor is metastatic OR surgical resection of tumor will likely result in severe morbidity. Non-Small Cell Lung Cancer-Approve if the patient has ROS1-positive metastatic disease and the mutation was detected by an approved test. Pediatric Diffuse High-Grade Glioma- approve if the tumor is positive for neurotrophic receptor tyrosine kinase (NTRK) gene fusion AND the medication is used either as adjuvant therapy or for recurrent or progressive disease.
<b>Indications</b>	All FDA-approved Indications, Some Medically-accepted Indications.
<b>Off-Label Uses</b>	Pediatric Diffuse High-Grade Glioma
<b>Part B Prerequisite</b>	No



# RUBRACA

## Products Affected

- RUBRACA

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	Diagnosis for which Rubraca is being used. BRCA-mutation (germline or somatic) status. Other medications tried for the diagnosis provided
<b>Age Restrictions</b>	18 years and older
<b>Prescriber Restrictions</b>	N/A
<b>Coverage Duration</b>	1 year
<b>Other Criteria</b>	<p>Maintenance Therapy of Ovarian, Fallopian tube or Primary peritoneal cancer-Approve if the patient is in complete or partial response after a platinum-based chemotherapy regimen and the patient is in complete or partial response to first-line primary treatment or if the patient has recurrent disease and has a BRCA mutation. Castration-Resistant Prostate Cancer - Approve if the patient meets the following criteria (A, B, C, and D): A) The patient has metastatic disease that is BRCA-mutation positive (germline and/or somatic) AND B) The patient meets one of the following criteria (i or ii): i. The medication is used concurrently with a gonadotropin-releasing hormone (GnRH) analog OR ii. The patient has had a bilateral orchiectomy AND C) The patient has been previously treated with at least one androgen receptor-directed therapy AND D) The patient meets one of the following criteria (i or ii): i. The patient has been previously treated with at least one taxane-based chemotherapy OR ii. The patient is not a candidate or is intolerant to taxane-based chemotherapy. Pancreatic adenocarcinoma-approve if pt has a BRCA mutation or PALB2 mutation AND pt has tried platinum-based chemotherapy AND has not had disease progression following the most recent platinum-based chemotherapy. Uterine leiomyosarcoma-approve if the patient has BRCA2-altered disease and has tried one systemic regimen.</p>

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Indications</b>	All FDA-approved Indications, Some Medically-accepted Indications.
<b>Off-Label Uses</b>	Uterine Leiomyosarcoma, Pancreatic Adenocarcinoma
<b>Part B Prerequisite</b>	No

# RUFINAMIDE

## Products Affected

- *rufinamide*

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	Diagnosis
<b>Age Restrictions</b>	Patients 1 years of age or older
<b>Prescriber Restrictions</b>	Prescribed by or in consultation with a neurologist
<b>Coverage Duration</b>	1 year
<b>Other Criteria</b>	Initial therapy-approve if rufinamide is being used for adjunctive treatment. Continuation-approve if the patient is responding to therapy
<b>Indications</b>	All FDA-approved Indications, Some Medically-accepted Indications.
<b>Off-Label Uses</b>	Treatment-Refractory Seizures/Epilepsy
<b>Part B Prerequisite</b>	No

# RYDAPT

## Products Affected

- RYDAPT

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	For AML, FLT3 status
<b>Age Restrictions</b>	18 years and older
<b>Prescriber Restrictions</b>	N/A
<b>Coverage Duration</b>	1 year
<b>Other Criteria</b>	AML -approve if the patient is FLT3-mutation positive as detected by an approved test. Myeloid or lymphoid Neoplasms with eosinophilia-approve if the patient has an FGFR1 rearrangement or has an FLT3 rearrangement.
<b>Indications</b>	All FDA-approved Indications, Some Medically-accepted Indications.
<b>Off-Label Uses</b>	Myeloid or lymphoid Neoplasms with eosinophilia
<b>Part B Prerequisite</b>	No

# SAPROPTERIN

---

## Products Affected

- *sapropterin*

PA Criteria	Criteria Details
Exclusion Criteria	PENDING CMS REVIEW
Required Medical Information	PENDING CMS REVIEW
Age Restrictions	PENDING CMS REVIEW
Prescriber Restrictions	PENDING CMS REVIEW
Coverage Duration	PENDING CMS REVIEW
Other Criteria	PENDING CMS REVIEW
Indications	PENDING CMS REVIEW
Off-Label Uses	PENDING CMS REVIEW
Part B Prerequisite	PENDING CMS REVIEW

# SCSEMBLIX

## Products Affected

- SCSEMBLIX ORAL TABLET 100 MG, 20 MG, 40 MG

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	Diagnosis
<b>Age Restrictions</b>	18 years and older
<b>Prescriber Restrictions</b>	N/A
<b>Coverage Duration</b>	1 year
<b>Other Criteria</b>	Chronic Myeloid Leukemia (CML)-approve if the patient meets the following (A and B): A) Patient has Philadelphia chromosome-positive chronic myeloid leukemia, AND B) Patient meets one of the following (i or ii): i. The chronic myeloid leukemia is T315I-positive, OR ii. Patient has tried at least two other tyrosine kinase inhibitors indicated for use in Philadelphia chromosome-positive chronic myeloid leukemia. Note: Examples of tyrosine kinase inhibitors include imatinib tablets, Bosulif (bosutinib tablets), Iclusig (ponatinib tablets), Sprycel (dasatinib tablets), and Tasigna (nilotinib capsules). Myeloid/Lymphoid Neoplasms with Eosinophilia - approve if the tumor has an ABL1 rearrangement.
<b>Indications</b>	All FDA-approved Indications, Some Medically-accepted Indications.
<b>Off-Label Uses</b>	Myeloid/Lymphoid Neoplasms with Eosinophilia
<b>Part B Prerequisite</b>	No

# SIGNIFOR

## Products Affected

- SIGNIFOR

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	N/A
<b>Age Restrictions</b>	18 years and older (initial therapy)
<b>Prescriber Restrictions</b>	Prescribed by or in consultation with an endocrinologist or a physician or specializes in the treatment of Cushing's syndrome (initial therapy)
<b>Coverage Duration</b>	Cushing's disease/syndrome-Initial therapy - 4 months, Continuation therapy - 1 year.
<b>Other Criteria</b>	Cushing's disease, initial therapy - approve if, according to the prescribing physician, the patient is not a candidate for surgery, or surgery has not been curative. Cushing's disease, continuation therapy - approve if the patient has already been started on Signifor/Signifor LAR and, according to the prescribing physician, the patient has had a response and continuation of therapy is needed to maintain response.
<b>Indications</b>	All FDA-approved Indications.
<b>Off-Label Uses</b>	N/A
<b>Part B Prerequisite</b>	No

# SIRTURO

## Products Affected

- SIRTURO

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	Patients weighing less than 15 kg
<b>Required Medical Information</b>	Diagnosis, concomitant therapy
<b>Age Restrictions</b>	Patients 5 years of age or older
<b>Prescriber Restrictions</b>	Prescribed by, or in consultation with an infectious diseases specialist
<b>Coverage Duration</b>	9 months
<b>Other Criteria</b>	Tuberculosis (Pulmonary)-Approve if the patient has multidrug-resistant tuberculosis and the requested medication is prescribed as part of a combination regimen with other anti-tuberculosis agents
<b>Indications</b>	All FDA-approved Indications.
<b>Off-Label Uses</b>	N/A
<b>Part B Prerequisite</b>	No



# SKYRIZI

## Products Affected

- SKYRIZI SUBCUTANEOUS PEN INJECTOR
- SKYRIZI SUBCUTANEOUS SYRINGE 150 MG/ML
- SKYRIZI SUBCUTANEOUS WEARABLE INJECTOR 180 MG/1.2 ML (150 MG/ML), 360 MG/2.4 ML (150 MG/ML)

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	Concurrent Use with other Biologics or Targeted Synthetic Disease-Modifying Antirheumatic Drugs (DMARDs)
<b>Required Medical Information</b>	Diagnosis, Previous medication use
<b>Age Restrictions</b>	PP-18 years of age and older (initial therapy)
<b>Prescriber Restrictions</b>	PP-Prescribed by or in consultation with a dermatologist (initial therapy), PsA-prescribed by or in consultation with a rheumatologist or dermatologist (initial therapy), CD-presc/consult-gastro (initial therapy)
<b>Coverage Duration</b>	Approve through end of plan year
<b>Other Criteria</b>	INITIAL THERAPY: PLAQUE PSORIASIS (PP) [A or B]: A) tried at least one traditional systemic agent for psoriasis (e.g., methotrexate [MTX], cyclosporine, acitretin tablets, or PUVA) for at least 3 months, unless intolerant. (Note: a 3-month trial or previous intolerance to at least one biologic also counts) or B) contraindication to MTX. PSORIATIC ARTHRITIS (PsA): approve. CROHN'S DISEASE (CD) [one of A, B, C, or D]: A) tried or is currently taking corticosteroids, unless contraindicated, B) tried one other conventional systemic therapy (e.g., azathioprine, 6-mercaptopurine, MTX) [Notes: a trial of a biologic that is not a biosimilar of Skyrizi also counts. A trial of mesalamine does not count as a systemic agent], C) patient has enterocutaneous (perianal or abdominal) or rectovaginal fistulas, or D) patient had ileocolonic resection to reduce the chance of CD recurrence. ALL INDICATIONS: Patient must be receiving induction dosing with Skyrizi IV within 3 months of initiating therapy with Skyrizi subcutaneous. CONTINUATION THERAPY: ALL INDICATIONS: patient has responded to therapy.

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Indications</b>	All FDA-approved Indications.
<b>Off-Label Uses</b>	N/A
<b>Part B Prerequisite</b>	No

## SOFOSBUVIR/VELPATASVIR

### Products Affected

- SOFOSBUVIR-VELPATASVIR

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	Combination use with other direct acting antivirals, excluding ribavirin.
<b>Required Medical Information</b>	Diagnosis
<b>Age Restrictions</b>	3 years or older
<b>Prescriber Restrictions</b>	Prescribed by or in consultation with a gastroenterologist, hepatologist, infectious diseases physician, or a liver transplant physician
<b>Coverage Duration</b>	Criteria will be applied consistent with current AASLD/IDSA guidance.
<b>Other Criteria</b>	Criteria will be applied according to AASLD guidelines.
<b>Indications</b>	All FDA-approved Indications, Some Medically-accepted Indications.
<b>Off-Label Uses</b>	Indications consistent with current AASLD/IDSA guidance
<b>Part B Prerequisite</b>	No

# SOLARAZE

## Products Affected

- *diclofenac sodium topical gel 3 %*

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	N/A
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Authorization will be for 6 months.
Other Criteria	N/A
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

# SOMAVERT

## Products Affected

- SOMAVERT

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	Diagnosis, previous therapy, concomitant therapy
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	Prescribed by or in consultation with an endocrinologist
<b>Coverage Duration</b>	1 year
<b>Other Criteria</b>	ACROMEGALY (A or B): A) inadequate response to surgery and/or radiotherapy or patient not an appropriate candidate or B) patient is experiencing negative effects due to tumor size (e.g., optic nerve compression) and has pre-treatment (baseline) insulin-like growth factor-1 (IGF-1) level above the upper limit of normal based on age and gender.
<b>Indications</b>	All FDA-approved Indications.
<b>Off-Label Uses</b>	N/A
<b>Part B Prerequisite</b>	No

# SORAFENIB

## Products Affected

- *sorafenib*

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	N/A
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	N/A
<b>Coverage Duration</b>	Authorization will be for 1 year
<b>Other Criteria</b>	Bone cancer, approve if the patient has recurrent chordoma or has osteosarcoma and has tried one standard chemotherapy regimen. GIST, approve if the patient has tried TWO of the following: imatinib mesylate, avapritinib, sunitinib, dasatinib, ripretinib or regorafenib. Differentiated (ie, papillary, follicular, oncocytic) thyroid carcinoma (DTC), approve if the patient is refractory to radioactive iodine treatment. Medullary thyroid carcinoma, approve if the patient has tried at least one systemic therapy. AML - Approve if disease is FLT3-ITD mutation positive as detected by an approved test and the medication is used in combination with azacitidine or decitabine or patient has had an allogeneic stem cell transplant and is in remission. Renal cell carcinoma (RCC)-approve if the patient has relapsed or advanced clear cell histology and the patient has tried at least one systemic therapy (e.g., Inlyta, Votrient, Sutent Cabometyx). Ovarian, fallopian tube, primary peritoneal cancer-approve if the patient has platinum resistant disease and sorafenib is used in combination with topotecan. HCC-approve if the patient has unresectable or metastatic disease. Soft tissue sarcoma-approve if the patient has angiosarcoma or desmoid tumors (aggressive fibromatosis) or solitary fibrous tumor/hemangiopericytoma. Myeloid/lymphoid neoplasms with eosinophilia-approve if the tumor has an FLT3 rearrangement. Please note

<b>PA Criteria</b>	<b>Criteria Details</b>
	for all diagnoses: Part B before Part D Step Therapy applies only to beneficiaries enrolled in an MA-PD plan
<b>Indications</b>	All FDA-approved Indications, Some Medically-accepted Indications.
<b>Off-Label Uses</b>	Bone cancer, Soft tissue sarcoma, gastrointestinal stromal tumors (GIST), medullary thyroid carcinoma, Acute Myeloid Leukemia, ovarian, fallopian tube, primary peritoneal cancer, myeloid/lymphoid neoplasms with eosinophilia
<b>Part B Prerequisite</b>	Yes

# SOTYKTU

## Products Affected

- SOTYKTU

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	Concurrent use with other biologics or with targeted synthetic disease-modifying antirheumatic drugs (DMARDs). Concurrent use with other potent immunosuppressants, including methotrexate.
<b>Required Medical Information</b>	Diagnosis
<b>Age Restrictions</b>	18 years and older (initial)
<b>Prescriber Restrictions</b>	Prescribed by or in consultation with a dermatologist (initial)
<b>Coverage Duration</b>	Approve through end of plan year
<b>Other Criteria</b>	INITIAL THERAPY: PLAQUE PSORIASIS (PP) [A or B]: A) tried at least one traditional systemic agent (e.g., MTX, cyclosporine (CSA), acitretin, PUVA) for at least 3 months, unless intolerant (Note: a trial of a biologic will also count) or B) contraindication to MTX. CONTINUATION THERAPY: patient had a response to therapy.
<b>Indications</b>	All FDA-approved Indications.
<b>Off-Label Uses</b>	N/A
<b>Part B Prerequisite</b>	No



# SPRYCEL

## Products Affected

- SPRYCEL ORAL TABLET 100 MG, 140 MG, 20 MG, 50 MG, 70 MG, 80 MG

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	Diagnosis for which Sprycel is being used. For indications of CML and ALL, the Philadelphia chromosome (Ph) status of the leukemia must be reported. For melanoma, cutaneous- KIT mutation and previous therapies.
<b>Age Restrictions</b>	GIST/bone cancer/ melanoma, cutaneous-18 years and older
<b>Prescriber Restrictions</b>	N/A
<b>Coverage Duration</b>	Authorization will be for 1 year
<b>Other Criteria</b>	For CML, patient must have Ph-positive CML. For ALL, patient must have Ph-positive ALL. For Bone Cancer-approve if patient has chondrosarcoma or chordoma. GIST - approve if the patient has tried imatinib or avapritinib. For melanoma, cutaneous - approve if patient has metastatic or unresectable disease AND has an activating KIT mutation AND has tried at least one systemic regimen.
<b>Indications</b>	All FDA-approved Indications, Some Medically-accepted Indications.
<b>Off-Label Uses</b>	GIST, bone cancer, melanoma cutaneous
<b>Part B Prerequisite</b>	No

# STELARA

## Products Affected

- STELARA SUBCUTANEOUS SOLUTION
- STELARA SUBCUTANEOUS SYRINGE 45 MG/0.5 ML, 90 MG/ML

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	Ustekinumab should not be given in combination with a Biologic DMARD or Targeted Synthetic DMARD
<b>Required Medical Information</b>	Diagnosis, concurrent medications, previous drugs tried.
<b>Age Restrictions</b>	PP-6 years and older (initial therapy).
<b>Prescriber Restrictions</b>	Plaque psoriasis.Prescribed by or in consultation with a dermatologist (initial therapy). PsA-prescribed by or in consultation with a rheumatologist or dermatologist (initial therapy). CD/UC-prescribed by or in consultation with a gastroenterologist (initial therapy).
<b>Coverage Duration</b>	Approve through end of plan year
<b>Other Criteria</b>	INITIAL THERAPY for STELARA SC: PLAQUE PSORIASIS (PP) [A or B]: A) tried one traditional systemic agent (e.g., methotrexate [MTX], cyclosporine, acitretin, psoralen plus PUVA) for at least 3 months, unless intolerant or B) contraindication to MTX. (Note: a 3-month trial or intolerance of at least one biologic that is not Stelara or a Stelara biosimilar also counts.) CROHN'S DISEASE (CD) [A and B]: A) receiving/received single IV loading dose within 2 months of initiating therapy with Stelara SC, and B) (a, b, c or d): a) tried or is currently taking corticosteroids (CS), or CS are contraindicated, b) tried one conventional systemic therapy, c) has enterocutaneous (perianal or abdominal) or rectovaginal fistulas, or d) had ileocolonic resection to reduce the chance of CD recurrence. ULCERATIVE COLITIS (UC) [A and B]: A) receiving/received single IV loading dose within 2 months of initiating therapy with Stelara SC and B) meets one of the following (a or b): a) tried one systemic agent or b) has pouchitis and tried an antibiotic, probiotic, CS enema or mesalamine enema. INDUCTION THERAPY for STELARA IV: UC [A or B]: A) tried one systemic agent or B) has pouchitis and has tried an antibiotic,

PA Criteria	Criteria Details
	probiotic, CS enema or mesalamine enema. CD, approve single dose of IV if meets A, B, C, or D: A) tried or is currently taking CS, or CS are contraindicated, B) tried one other conventional systemic therapy (e.g., azathioprine, 6-MP, MTX, certolizumab, vedolizumab, adalimumab, infliximab), C) has enterocutaneous (perianal or abdominal) or rectovaginal fistulas, or D) had ileocolonic resection to reduce the chance of Crohn's disease recurrence. CONTINUATION THERAPY: PP/PsA/CD/UC: patient has responded to therapy.
<b>Indications</b>	All FDA-approved Indications.
<b>Off-Label Uses</b>	N/A
<b>Part B Prerequisite</b>	No

# STIVARGA

## Products Affected

- STIVARGA

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	Diagnosis for which Stivarga is being used. Prior therapies tried. For metastatic CRC, KRAS/NRAS mutation status.
<b>Age Restrictions</b>	18 years and older
<b>Prescriber Restrictions</b>	N/A
<b>Coverage Duration</b>	1 year
<b>Other Criteria</b>	For GIST, patient must have previously been treated with imatinib or Ayvakit and sunitinib or Sprycel. For HCC, patient must have previously been treated with at least one systemic regimen. Soft tissue sarcoma, advanced or metastatic disease-approve if the patient has non-adipocytic sarcoma, angiosarcoma, or pleomorphic rhabdomyosarcoma. Bone Cancer-approve if the patient has relapsed/refractory or metastatic disease AND the patient has tried one systemic chemotherapy regimen AND pt has Ewing sarcoma or osteosarcoma. Colon and Rectal cancer/Appendiceal cancer-approve if the patient has advanced or metastatic disease, has been previously treated with a fluoropyrimidine, oxaliplatin, irinotecan and if the patient meets one of the following (i or ii): i. patient's tumor or metastases are wild-type RAS (KRAS wild type and NRAS wild type), the patient has tried Erbitux or Vectibix or the patient's tumor did not originate on the left side of the colon (from the splenic fixture to rectum) or ii. the patient's tumor or metastases has a RAS mutation (either KRAS mutation or NRAS mutation). Glioblastoma-approve if the patient has recurrent or progressive disease.
<b>Indications</b>	All FDA-approved Indications, Some Medically-accepted Indications.
<b>Off-Label Uses</b>	Soft tissue Sarcoma, Bone Cancer, Glioblastoma, Appendiceal cancer

PA Criteria	Criteria Details
Part B Prerequisite	No

# SUCRAID

## Products Affected

- SUCRAID

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	Diagnosis, genetic and lab test results (as specified in the Other Criteria field)
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	Prescribed by or in consultation with a geneticist, gastroenterologist, a metabolic disorder sub-specialist, or a physician who specializes in the treatment of congenital diarrheal disorders
<b>Coverage Duration</b>	1 year
<b>Other Criteria</b>	Approve if the patient meets the following criteria (A and B): A) The diagnosis is established by one of the following (i or ii): i. Patient has endoscopic biopsy of the small bowel with disaccharidase levels consistent with congenital sucrose-isomaltase deficiency as evidenced by ALL of the following (a, b, c, and d): a) Decreased (usually absent) sucrase (normal reference: greater than 25 U/g protein), b) Decreased or normal isomaltase (palatinase) [normal reference: greater than 5 U/g protein], c) Decreased maltase (normal reference: greater than 100 U/g protein), d) Decreased or normal lactase (normal reference: greater than 15 U/g protein) OR ii. Patient has a molecular genetic test demonstrating homozygous or compound heterozygous pathogenic or likely pathogenic sucrase-isomaltase gene variant AND B) Patient has symptomatic congenital sucrose-isomaltase deficiency (e.g., diarrhea, bloating, abdominal cramping).
<b>Indications</b>	All FDA-approved Indications.
<b>Off-Label Uses</b>	N/A

PA Criteria	Criteria Details
Part B Prerequisite	No

## SUNITINIB

### Products Affected

- sunitinib malate*

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	N/A
<b>Age Restrictions</b>	18 years and older
<b>Prescriber Restrictions</b>	N/A
<b>Coverage Duration</b>	1 year
<b>Other Criteria</b>	Gastrointestinal stromal tumors (GIST), approve if the patient has tried imatinib or Ayvakit or if the patient has succinate dehydrogenase (SDH)-deficient GIST. Chordoma, approve if the patient has recurrent disease. Differentiated thyroid carcinoma, approve if the patient is refractory to radioactive iodine therapy. Medullary thyroid carcinoma, approve if the patient has tried at least one systemic therapy. Meningioma, approve if the patient has recurrent or progressive disease. Thymic carcinoma - has tried at least one systemic chemotherapy. Renal Cell Carcinoma (RCC) - approve if the patient has relapsed or advanced disease. Neuroendocrine tumors of the pancreas-approve for advanced or metastatic disease. Pheochromocytoma/paraganglioma-approve if the patient has unresectable or metastatic disease. Myeloid/lymphoid neoplasms with eosinophilia-approve if the tumor has an FLT3 rearrangement.
<b>Indications</b>	All FDA-approved Indications, Some Medically-accepted Indications.
<b>Off-Label Uses</b>	Chordoma, angiosarcoma, solitary fibrous tumor/hemangiopericytoma, alveolar soft part sarcoma (ASPS), differentiated (ie, papillary, follicular, and oncocytic carcinoma) thyroid carcinoma, medullary thyroid carcinoma, meningioma, thymic carcinoma, pheochromocytoma/paraganglioma,



PA Criteria	Criteria Details
	myeloid/lymphoid neoplasms with eosinophilia, GIST-unresectable succinate dehydrogenase (SDH)-deficient GIST, or use after avapritinib.
<b>Part B Prerequisite</b>	No

## SYMDEKO

### Products Affected

- SYMDEKO

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	Patients with unknown CFTR gene mutations, Combination therapy with Orkambi, Kalydeco or Trikafta
<b>Required Medical Information</b>	Diagnosis, specific CFTR gene mutations
<b>Age Restrictions</b>	Six years of age and older
<b>Prescriber Restrictions</b>	Prescribed by or in consultation with a pulmonologist or a physician who specializes in CF
<b>Coverage Duration</b>	1 year
<b>Other Criteria</b>	CF - Approve if the pt meets A, B and C: A) pt has at least one mutation in the CFTR gene that is considered to be pathogenic or likely pathogenic, and B) pt must have positive CF newborn screening test or family history of CF or clinical presentation consistent with signs and symptoms of CF and C) evidence of abnormal CFTR function as demonstrated by i, ii, or iii: (i) elevated sweat chloride test or (ii) two CFTR mutations or (iii) abnormal nasal potential difference.
<b>Indications</b>	All FDA-approved Indications.
<b>Off-Label Uses</b>	N/A
<b>Part B Prerequisite</b>	No

# SYMLIN

## Products Affected

- SYMLINPEN 120
- SYMLINPEN 60

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	N/A
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Authorization will be for 1 year
Other Criteria	N/A
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

# TABRECTA

## Products Affected

- TABRECTA

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	Diagnosis
Age Restrictions	18 years and older
Prescriber Restrictions	N/A
Coverage Duration	Authorization will be for 1 year
Other Criteria	Non-Small Cell Lung Cancer (NSCLC)-Approve if the patient has advanced or metastatic disease AND the tumor is positive for a mutation that leads to mesenchymal-epithelial transition (MET) exon 14 skipping or high-level MET amplification, as detected by an approved test.
Indications	All FDA-approved Indications, Some Medically-accepted Indications.
Off-Label Uses	Non-small cell lung cancer with high-level MET amplification.
Part B Prerequisite	No

# TADALAFIL

## Products Affected

- *tadalafil oral tablet 2.5 mg, 5 mg*

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	Indication for which tadalafil is being prescribed.
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	N/A
<b>Coverage Duration</b>	Authorization will be for 12 mos.
<b>Other Criteria</b>	Benign prostatic hyperplasia (BPH), after confirmation that tadalafil is being prescribed as once daily dosing, to treat the signs and symptoms of BPH and not for the treatment of erectile dysfunction (ED).
<b>Indications</b>	All FDA-approved Indications.
<b>Off-Label Uses</b>	N/A
<b>Part B Prerequisite</b>	No

# TAFAMIDIS

## Products Affected

- VYNDAMAX

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	Concomitant use with Onpattro or Tegsedi or Wainua. Concurrent use of Vyndaqel and Vyndamax.
<b>Required Medical Information</b>	Diagnosis, genetic tests and lab results (as specified in the Other Criteria field)
<b>Age Restrictions</b>	18 years and older
<b>Prescriber Restrictions</b>	Prescribed by or in consultation with a cardiologist or a physician who specializes in the treatment of amyloidosis
<b>Coverage Duration</b>	1 year
<b>Other Criteria</b>	Cardiomyopathy of Wild-Type or Hereditary Transthyretin Amyloidosis-approve if the diagnosis was confirmed by one of the following (i, ii or iii): i. A technetium pyrophosphate scan (i.e., nuclear scintigraphy), ii. Amyloid deposits are identified on cardiac biopsy OR iii. patient had genetic testing which, according to the prescriber, identified a TTR mutation AND Diagnostic cardiac imaging (e.g., echocardiogram, cardiac magnetic imaging) has demonstrated cardiac involvement (e.g., increased thickness of the ventricular wall or interventricular septum).
<b>Indications</b>	All FDA-approved Indications.
<b>Off-Label Uses</b>	N/A
<b>Part B Prerequisite</b>	No

# TAFINLAR

## Products Affected

- TAFINLAR ORAL CAPSULE
- TAFINLAR ORAL TABLET FOR SUSPENSION

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	Diagnosis for which Tafinlar is being used. BRAF V600 mutations
<b>Age Restrictions</b>	1 year and older
<b>Prescriber Restrictions</b>	N/A
<b>Coverage Duration</b>	Authorization will be for 1 year
<b>Other Criteria</b>	Melanoma with BRAF V600 mutation AND patient has unresectable, advanced (including Stage III or Stage IV disease) or metastatic melanoma. Note-This includes adjuvant treatment in patients with Stage III disease with no evidence of disease post-surgery. For NSCLC, must have BRAF V600E mutation. Thyroid Cancer, anaplastic-must have BRAF V600-positive disease AND Tafinlar will be taken in combination with Mekinist, unless intolerant AND the patient has locally advanced or metastatic anaplastic disease. Thyroid Cancer, differentiated (i.e., papillary, follicular, or Hurthle cell) AND the patient has disease that is refractory to radioactive iodine therapy AND the patient has BRAF-positive disease. Biliary Tract Cancer-approve if the patient has tried at least one systemic chemotherapy regimen, patient has BRAF V600 mutation positive disease and the medication will be taken in combination with Mekinist. Central Nervous System Cancer-approve if the medication is being used for one of the following situations (i, ii, or iii): i. Adjuvant treatment of one of the following conditions (a, b, or c): a) Pilocytic astrocytoma OR b) Pleomorphic xanthoastrocytoma OR c) Ganglioglioma OR ii. Recurrent or progressive disease for one of the following conditions (a, b, c, or d): a) glioma OR b) Isocitrate dehydrogenase-2 (IDH2)-mutant astrocytoma OR

PA Criteria	Criteria Details
	<p>c) Glioblastoma OR d)Oligodendroglioma OR iii. Melanoma with brain metastases AND patient has BRAF V600 mutation-positive disease AND medication will be taken in combination with Mekinist (trametinib tablets). Histiocytic neoplasm-approve if patient has Langerhans cell histiocytosis and one of the following (a, b, or c): a) Multisystem disease OR b) Pulmonary disease OR c) Central nervous system lesions OR patient has Erdheim Chester disease AND patient has BRAF V600-mutation positive disease. Metastatic or solid tumors-approve if BRAF V600 mutation-positive disease AND medication will be taken in combination with Mekinist (trametinib tablets) AND patient has no satisfactory alternative treatment options. Ovarian, Fallopian Tube, or Primary Peritoneal Cancer-approve if the patient meets the following (A, B, and C): A) Patient has recurrent disease, AND B) Patient has BRAF V600 mutation-positive disease, AND C) The medication will be taken in combination with Mekinist (trametinib tablets). Hairy Cell Leukemia, approve if pt has not previously been treated with a BRAF inhibitor therapy and this will be used for relapsed/refractory disease and will be taken in combination with Mekinist. Small bowel adenocarcinoma, approve if pt has BRAF V600E mutation-positive advanced or metastatic disease and this will be used with Mekinist AND (i or ii): i) this will be used as initial therapy and pt has received previous FOLFOX/CAPEOX therapy in the adjuvant setting within the past 12 months or has a contraindication, or (ii) this will be used as second-line and subsequent therapy.</p>
<b>Indications</b>	All FDA-approved Indications, Some Medically-accepted Indications.
<b>Off-Label Uses</b>	Patients with Differentiated Thyroid Cancer, Biliary tract cancer, central nervous system cancer, histiocytic neoplasm, Ovarian, Fallopian Tube, or Primary Peritoneal Cancer, hairy cell leukemia, small bowel adenocarcinoma
<b>Part B Prerequisite</b>	No



# TAGRISO

## Products Affected

- TAGRISSO

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	Diagnosis
<b>Age Restrictions</b>	18 years and older
<b>Prescriber Restrictions</b>	N/A
<b>Coverage Duration</b>	1 year
<b>Other Criteria</b>	<p>NSCLC-EGFR Mutation Positive (other than EGFR T790M-Positive Mutation)- approve if the patient has advanced or metastatic disease, has EGFR mutation-positive NSCLC as detected by an approved test. Note-examples of EGFR mutation-positive NSCLC include the following mutations: exon 19 deletions, exon 21 (L858R) substitution mutations, L861Q, G719X and S7681. NSCLC-EGFR T790M mutation positive-approve if the patient has metastatic EGFR T790M mutation-positive NSCLC as detected by an approved test and has progressed on treatment with at least one of the EGFR tyrosine kinase inhibitors. NSCLC-Post resection-approve if the patient has completely resected disease and has received previous adjuvant chemotherapy or if the patient is ineligible to receive platinum based chemotherapy and the patient has EGFR exon 19 deletions or exon 21 L858R substitution mutations, as detected by an approved test.</p>
<b>Indications</b>	All FDA-approved Indications.
<b>Off-Label Uses</b>	N/A
<b>Part B Prerequisite</b>	No

# TALZENNA

## Products Affected

- TALZENNA

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	Diagnosis
Age Restrictions	18 years and older
Prescriber Restrictions	N/A
Coverage Duration	1 year
Other Criteria	Recurrent or metastatic breast cancer-approve if the patient has germline BRCA mutation-positive disease. In addition, patients new to therapy are required to try Lynparza prior to approval of Talzenna. Prostate cancer - approve if the patient has metastatic castration resistant prostate cancer, AND is using this medication concurrently with a gonadotropin-releasing hormone (GnRH) analog or has had a bilateral orchiectomy AND the patient has homologous recombination repair (HRR) gene-mutated disease [Note: HRR gene mutations include ATM, ATR, BRCA1, BRCA2, CDK12, CHEK2, FANCA, MLH1, MRE11A, NBN, PALB2, or RAD51C] AND the medication is used in combination with Xtandi (enzalutamide capsules and tablets). In addition, patients new to therapy with BRCA-mutated-positive prostate cancer are required to try Lynparza prior to approval of Talzenna.
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

# TASIGNA

## Products Affected

- TASIGNA ORAL CAPSULE 150 MG, 200 MG, 50 MG

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	Diagnosis for which Tasigna is being used. For indication of CML and ALL, the Philadelphia chromosome (Ph) status of the leukemia must be reported. For indication of gastrointestinal stromal tumor (GIST) and melanoma, cutaneous, prior therapies tried. For melanoma, cutaneous, KIT mutation status.
<b>Age Restrictions</b>	ALL/GIST/Myeloid/lymphoid neoplasms/melanoma, cutaneous-18 years and older
<b>Prescriber Restrictions</b>	N/A
<b>Coverage Duration</b>	Authorization will be for 12 months.
<b>Other Criteria</b>	Acute lymphoblastic leukemia, philadelphia chromosome positive or chronic myeloid leukemia- approve. For GIST, approve if the patient has tried two of the following: imatinib, avapritinib, sunitinib, dasatinib, regorafenib or ripretinib. Myeloid/lymphoid neoplasms with eosinophilia-approve if the tumor has an ABL1 rearrangement. Pigmented villonodular synovitis/tenosynovial giant cell tumor-approve if the patient has tried Turalio or cannot take Turalio, according to the prescriber. For melanoma, cutaneous - approve if the patient has metastatic or unresectable disease AND has an activating KIT mutation AND has tried at least one systemic regimen.
<b>Indications</b>	All FDA-approved Indications, Some Medically-accepted Indications.
<b>Off-Label Uses</b>	Philadelphia positive Acute Lymphoblastic Leukemia (ALL) and Gastrointestinal Stromal Tumor (GIST), Pigmented villonodular synovitis/tenosynovial giant cell tumor, Myeloid/Lymphoid neoplasms with Eosinophilia, melanoma cutaneous.

PA Criteria	Criteria Details
Part B Prerequisite	No

# TAZAROTENE

## Products Affected

- *tazarotene topical cream 0.1 %*
- *tazarotene topical gel*

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	Cosmetic uses
<b>Required Medical Information</b>	N/A
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	N/A
<b>Coverage Duration</b>	1 year
<b>Other Criteria</b>	N/A
<b>Indications</b>	All Medically-accepted Indications.
<b>Off-Label Uses</b>	N/A
<b>Part B Prerequisite</b>	No

# TAZVERIK

## Products Affected

- TAZVERIK

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	Diagnosis
<b>Age Restrictions</b>	Epithelioid Sarcoma-16 years and older, Follicular Lymphoma-18 years and older
<b>Prescriber Restrictions</b>	N/A
<b>Coverage Duration</b>	1 year
<b>Other Criteria</b>	Epithelioid Sarcoma-approve if the patient has metastatic or locally advanced disease and the patient is not eligible for complete resection. Follicular Lymphoma-approve if the patient has relapsed or refractory disease and there are no appropriate alternative therapies or the patient has tried at least two prior systemic therapies.
<b>Indications</b>	All FDA-approved Indications.
<b>Off-Label Uses</b>	N/A
<b>Part B Prerequisite</b>	No

# TEPMETKO

## Products Affected

- TEPMETKO

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	Diagnosis
<b>Age Restrictions</b>	18 years and older
<b>Prescriber Restrictions</b>	N/A
<b>Coverage Duration</b>	1 year
<b>Other Criteria</b>	NSCLC-approve if the patient has advanced or metastatic disease and the tumor is positive for mesenchymal-epithelial transition (MET) exon 14 skipping mutations or patient has high-level MET amplification, as detected by an approved test.
<b>Indications</b>	All FDA-approved Indications, Some Medically-accepted Indications.
<b>Off-Label Uses</b>	Non-small cell lung cancer with high-level MET amplification.
<b>Part B Prerequisite</b>	No

# TERIPARATIDE

## Products Affected

- TERIPARATIDE SUBCUTANEOUS  
PEN INJECTOR 20 MCG/DOSE  
(620MCG/2.48ML)

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	Concomitant use with other medications for osteoporosis
<b>Required Medical Information</b>	N/A
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	N/A
<b>Coverage Duration</b>	High risk for fracture-2 yrs, Not high risk-approve a max of 2 yrs of therapy (total)/lifetime.
<b>Other Criteria</b>	<p>INITIAL THERAPY: Postmenopausal Osteoporosis (PMO) Treatment, Increase Bone Mass in Men (see Note 1 below) with Primary or Hypogonadal Osteoporosis, and Treatment of Glucocorticosteroid-Induced Osteoporosis (GIO): (one of A, B, C, D or E): A) tried one oral bisphosphonate or cannot take due to swallowing difficulties or inability to remain upright after administration, B) pre-existing gastrointestinal condition (e.g., esophageal lesions/ulcers, abnormalities of the esophagus that delay esophageal emptying [stricture, achalasia]), C) tried an IV bisphosphonate (PMO-ibandronate or zoledronic acid, all other diagnoses-zoledronic acid), D) severe renal impairment (creatinine clearance [CrCL] less than 35 mL/min) or chronic kidney disease (CKD), or E) patient had an osteoporotic fracture or fragility fracture at any time in the past.</p> <p>CONTINUATION THERAPY: ALL INDICATIONS: if the patient has taken teriparatide for two years, approve if the patient is at high risk for fracture. Examples of high risk for fracture include a previous osteoporotic fracture or fragility fracture, receipt of medications that increase the risk of osteoporosis, advanced age, and very low bone mineral density. Note 1: a</p>



PA Criteria	Criteria Details
	man is defined as an individual with the biological traits of a man, regardless of the individual's gender identity or gender expression.
<b>Indications</b>	All FDA-approved Indications.
<b>Off-Label Uses</b>	N/A
<b>Part B Prerequisite</b>	No

# TETRABENAZINE

## Products Affected

- tetrabenazine oral tablet 12.5 mg, 25 mg

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	N/A
<b>Age Restrictions</b>	18 years and older
<b>Prescriber Restrictions</b>	For treatment of chorea associated with Huntington's disease, Tourette syndrome or related tic disorders, hyperkinetic dystonia, or hemiballism, must be prescribed by or after consultation with a neurologist. For TD, must be prescribed by or after consultation with a neurologist or psychiatrist.
<b>Coverage Duration</b>	Authorization will be for 1 year.
<b>Other Criteria</b>	Chorea associated with Huntington's Disease-approve if the diagnosis of Huntington's Disease is confirmed by genetic testing.
<b>Indications</b>	All FDA-approved Indications, Some Medically-accepted Indications.
<b>Off-Label Uses</b>	Tardive dyskinesia (TD). Tourette syndrome and related tic disorders. Hyperkinetic dystonia. Hemiballism.
<b>Part B Prerequisite</b>	No

# THALOMID

## Products Affected

- THALOMID ORAL CAPSULE 100 MG, 150 MG, 200 MG, 50 MG

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	N/A
<b>Age Restrictions</b>	MM, myelofibrosis-18 years and older
<b>Prescriber Restrictions</b>	N/A
<b>Coverage Duration</b>	Authorization will be for 1 year
<b>Other Criteria</b>	Erythem Nodosum Leprosus-approve. Multiple Myeloma-approve. Discoid lupus erythematosus or cutaneous lupus erythematosus, approve if the patient has tried at least two other therapies (eg, corticosteroids [oral, topical, intralesional], hydroxychloroquine, tacrolimus [Protopic], methotrexate, dapson, acitretin [Soriatane]). Myelofibrosis, approve if according to the prescriber the patient has anemia and has serum erythropoietin levels greater than or equal to 500 mU/mL or if the patient has serum erythropoietin level less than 500 mU/mL and experienced no response or loss of response to erythropoietic stimulating agents. Prurigo nodularis, approve. Recurrent aphthous ulcers or aphthous stomatitis, approve if the patient has tried at least two other therapies (eg, topical or intralesional corticosteroids, systemic corticosteroids, topical anesthetics/analgesics [eg, benzocaine lozenges], antimicrobial mouthwashes [eg, tetracycline], acyclovir, colchicine). Kaposi's Sarcoma-approve if the patient has tried at least one regimen or therapy and has relapsed or refractory disease. Castleman's disease-approve if the patient has multicentric Castleman's disease, and is negative for the human immunodeficiency virus (HIV) and human herpesvirus-8 (HHV-8).
<b>Indications</b>	All FDA-approved Indications, Some Medically-accepted Indications.

PA Criteria	Criteria Details
<b>Off-Label Uses</b>	Discoid lupus erythematosus or cutaneous lupus erythematosus, Myelofibrosis, Prurigo nodularis, Recurrent aphthous ulcers or aphthous stomatitis, Kaposi's Sarcoma, Castleman's Disease.
<b>Part B Prerequisite</b>	No

# TIBSOVO

## Products Affected

- TIBSOVO

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	Diagnosis, IDH1 Status
<b>Age Restrictions</b>	All diagnoses (except chondrosarcoma)-18 years and older
<b>Prescriber Restrictions</b>	N/A
<b>Coverage Duration</b>	1 year
<b>Other Criteria</b>	AML- approve if the disease is isocitrate dehydrogenase-1 (IDH1) mutation positive, as detected by an approved test. Cholangiocarcinoma-approve if the disease is isocitrate dehydrogenase-1 (IDH1) mutation positive and has been previously treated with at least one chemotherapy regimen (Part B before Part D Step Therapy - applies only to beneficiaries enrolled in an MA-PD plan). Chondrosarcoma-approve if the disease is isocitrate dehydrogenase-1 (IDH1) mutation positive. Central nervous system cancer-approve if the patient has recurrent or progressive disease, AND patient has World Health Organization (WHO) grade 2 or 3 oligodendroglioma, OR Patient has WHO grade 2 astrocytoma. Myelodysplastic Syndrome-approve if patient has isocitrate dehydrogenase-1 (IDH1) mutation-positive disease AND relapsed or refractory disease.
<b>Indications</b>	All FDA-approved Indications, Some Medically-accepted Indications.
<b>Off-Label Uses</b>	Chondrosarcoma, Central nervous system cancer
<b>Part B Prerequisite</b>	Yes

## TOBRAMYCIN (NEBULIZATION)

### Products Affected

- *tobramycin in 0.225 % nacl*
- *tobramycin inhalation*

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	Diagnosis
<b>Age Restrictions</b>	Bronchiectasis, Non-cystic fibrosis-18 years and older
<b>Prescriber Restrictions</b>	CF-prescr/consult w/pulm/phys specializes in tx of CF.Bronchiectasis, non CF-prescr/consult w/pulm
<b>Coverage Duration</b>	1 year
<b>Other Criteria</b>	Part B versus Part D determination will be made at time of prior authorization review per CMS guidance. Cystic fibrosis/Bronchiectasis, non-cystic fibrosis-approve if the patient has pseudomonas aeruginosa in the culture of the airway.
<b>Indications</b>	All FDA-approved Indications, Some Medically-accepted Indications.
<b>Off-Label Uses</b>	Bronchiectasis, non-cystic fibrosis
<b>Part B Prerequisite</b>	No

# TOLVAPTAN

## Products Affected

- *tolvaptan*

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	Concurrent use with Jynarque.
<b>Required Medical Information</b>	Serum sodium less than 125 mEq/L at baseline or less marked hyponatremia, defined as serum sodium less than 135 mEq/L at baseline, that is symptomatic (eg, nausea, vomiting, headache, lethargy, confusion).
<b>Age Restrictions</b>	18 years and older
<b>Prescriber Restrictions</b>	N/A
<b>Coverage Duration</b>	Authorization will be for 30 days for initial therapy, 3 months for continuation of therapy
<b>Other Criteria</b>	Hyponatremia, initial therapy (including new starts, patients on therapy for less than 30 days, and patients restarting therapy) - Pt must meet ONE of the following: 1. serum sodium less than 125 mEq/L at baseline, OR 2. marked hyponatremia, defined as less than 135 mEq/L at baseline, that is symptomatic (eg, nausea, vomiting, headache, lethargy, confusion), OR 3. patient has already been started on tolvaptan and has received less than 30 days of therapy. Hyponatremia, continuation of therapy for patients established on therapy for at least 30 days - approve if the serum sodium level has increased from baseline (prior to initiating the requested drug) OR if the patient experienced improvement in at least one symptom, such as nausea, vomiting, headache, lethargy, or confusion.
<b>Indications</b>	All FDA-approved Indications.
<b>Off-Label Uses</b>	N/A
<b>Part B Prerequisite</b>	No

## TOPICAL AGENTS FOR ATOPIC DERMATITIS

### Products Affected

- *pimecrolimus*
- *tacrolimus topical*

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	N/A
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Authorization will be for 12 months.
Other Criteria	Authorize use in patients who have tried a prescription strength topical corticosteroid (brand or generic) for the current condition. Dermatologic condition on or around the eyes, eyelids, axilla, or genitalia, authorize use without a trial of a prescription strength topical corticosteroid.
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No



# TOPICAL RETINOID PRODUCTS

## Products Affected

- *tretinoin topical*

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	Coverage is not provided for cosmetic use.
<b>Required Medical Information</b>	N/A
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	N/A
<b>Coverage Duration</b>	Authorization will be for 12 months
<b>Other Criteria</b>	N/A
<b>Indications</b>	All Medically-accepted Indications.
<b>Off-Label Uses</b>	N/A
<b>Part B Prerequisite</b>	No

## TOPIRAMATE/ZONISAMIDE

### Products Affected

- EPRONTIA
- *topiramate oral capsule, sprinkle*
- *topiramate oral tablet*
- ZONISADE
- *zonisamide*

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	Coverage is not provided for weight loss or smoking cessation.
<b>Required Medical Information</b>	N/A
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	N/A
<b>Coverage Duration</b>	Authorization will be for 1 year.
<b>Other Criteria</b>	N/A
<b>Indications</b>	All Medically-accepted Indications.
<b>Off-Label Uses</b>	N/A
<b>Part B Prerequisite</b>	No

# TRANSDERMAL FENTANYL

## Products Affected

- *fentanyl transdermal patch 72 hour 100 mcg/hr, 12 mcg/hr, 25 mcg/hr, 50 mcg/hr, 75 mcg/hr*

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	Acute (i.e., non-chronic) pain.
<b>Required Medical Information</b>	Pain type (chronic vs acute), prior pain medications/therapies tried, concurrent pain medications/therapies
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	N/A
<b>Coverage Duration</b>	1 year
<b>Other Criteria</b>	For pain severe enough to require daily, around-the-clock, long-term opioid treatment, approve if all of the following criteria are met: 1) patient is not opioid naive, AND 2) non-opioid therapies have been tried and are being used in conjunction with opioid therapy according to the prescribing physician, AND 3) the prescribing physician has checked the patient's history of controlled substance prescriptions using state prescription drug monitoring program (PDMP), AND 4) the prescribing physician has discussed risks (eg, addiction, overdose) and realistic benefits of opioid therapy with the patient, AND 5) according to the prescriber physician there is a treatment plan (including goals for pain and function) in place and reassessments are scheduled at regular intervals. Patients with cancer, sickle cell disease, in hospice or who reside in a long term care facility are not required to meet above criteria. Clinical criteria incorporated into the quantity limit edits for all oral long-acting opioids (including transdermal fentanyl products) require confirmation that the indication is intractable pain (ie, FDA labeled use) prior to reviewing for quantity exception.
<b>Indications</b>	All FDA-approved Indications.

PA Criteria	Criteria Details
Off-Label Uses	N/A
Part B Prerequisite	No

## TRANSMUCOSAL FENTANYL DRUGS

### Products Affected

- *fentanyl citrate buccal lozenge on a handle*

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	N/A
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	N/A
<b>Coverage Duration</b>	Authorization will be for 12 months.
<b>Other Criteria</b>	For breakthrough pain in patients with cancer if patient is unable to swallow, has dysphagia, esophagitis, mucositis, or uncontrollable nausea/vomiting OR patient is unable to take 2 other short-acting narcotics (eg, oxycodone, morphine sulfate, hydromorphone, etc) secondary to allergy or severe adverse events AND patient is on or will be on a long-acting narcotic (eg, Duragesic), or the patient is on intravenous, subcutaneous, or spinal (intrathecal, epidural) narcotics (eg, morphine sulfate, hydromorphone, fentanyl citrate). Clinical criteria incorporated into the quantity limit edits for all transmucosal fentanyl drugs require confirmation that the indication is breakthrough cancer pain (ie, FDA labeled use) prior to reviewing for quantity exception.
<b>Indications</b>	All FDA-approved Indications.
<b>Off-Label Uses</b>	N/A
<b>Part B Prerequisite</b>	No

# TRELSTAR

## Products Affected

- TRELSTAR INTRAMUSCULAR SUSPENSION FOR RECONSTITUTION

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	Diagnosis
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	Prescribed by or in consultation with a oncologist or urologist
<b>Coverage Duration</b>	1 year
<b>Other Criteria</b>	Part B versus Part D determination will be made at time of prior authorization review per CMS guidance. Patients new to therapy, are required to try Eligard or Orgovyx prior to approval of Trelstar.
<b>Indications</b>	All FDA-approved Indications.
<b>Off-Label Uses</b>	N/A
<b>Part B Prerequisite</b>	No

# TREMFYA

## Products Affected

- TREMFYA SUBCUTANEOUS AUTO-INJECTOR
- TREMFYA SUBCUTANEOUS SYRINGE 100 MG/ML

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	Concurrent Use with other Biologics or Targeted Synthetic Disease-Modifying Antirheumatic Drugs (DMARDs)
<b>Required Medical Information</b>	Diagnosis
<b>Age Restrictions</b>	PP- 18 years of age and older (initial therapy)
<b>Prescriber Restrictions</b>	PP-Prescribed by or in consultation with a dermatologist (initial therapy only). PsA-prescribed by or in consultation with a dermatologist or rheumatologist (initial therapy).
<b>Coverage Duration</b>	Approve through end of plan year
<b>Other Criteria</b>	PP, initial therapy - approve if the pt meets one of the following criteria: 1) pt has tried at least one traditional systemic agent (eg, MTX, cyclosporine, acitretin, PUVA) for at least 3 months, unless intolerant (note: a biologic that is not a biosimilar of the requested product will also count) OR 2) pt has a contraindication to MTX as determined by the prescribing physician. PP/PsA continuation of therapy - approve if the pt is responding to therapy.
<b>Indications</b>	All FDA-approved Indications.
<b>Off-Label Uses</b>	N/A
<b>Part B Prerequisite</b>	No

# TRIENTINE

## Products Affected

- *trientine oral capsule 250 mg*

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	Diagnosis
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	Prescribed by or in consultation with a gastroenterologist, hepatologist, or liver transplant physician.
<b>Coverage Duration</b>	Authorization will be for 1 year
<b>Other Criteria</b>	For Wilson's Disease, approve if the patient meets A and B: A) Diagnosis of Wilson's disease is confirmed by ONE of the following (i or ii): i. Genetic testing results confirming biallelic pathogenic ATP7B mutations (in either symptomatic or asymptomatic individuals), OR ii. Confirmation of at least two of the following (a, b, c, or d): a. Presence of Kayser-Fleischer rings, OR b. Serum ceruloplasmin levels less than 20mg/dL, OR c. Liver biopsy findings consistent with Wilson's disease, OR d. 24-hour urinary copper greater than 40 micrograms/24 hours, AND B) Patient meets ONE of the following: 1) Patient has tried a penicillamine product and per the prescribing physician the patient is intolerant to penicillamine therapy, OR 2) Per the prescribing physician, the patient has clinical features indicating the potential for intolerance to penicillamine therapy (ie, history of any renal disease, congestive splenomegaly causing severe thrombocytopenia, autoimmune tendency), OR 3) Per the prescribing physician, the patient has a contraindication to penicillamine therapy, OR 4) The patient has neurologic manifestations of Wilson's disease, OR 5) The patient is pregnant, OR 6) the patient has been started on therapy with trientine.
<b>Indications</b>	All FDA-approved Indications.



PA Criteria	Criteria Details
Off-Label Uses	N/A
Part B Prerequisite	No

# TRIKAFTA

## Products Affected

- TRIKAFTA ORAL GRANULES IN PACKET, SEQUENTIAL
- TRIKAFTA ORAL TABLETS, SEQUENTIAL

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	Patients with unknown CFTR gene mutations. Combination therapy with Orkambi, Kalydeco or Symdeko.
<b>Required Medical Information</b>	Diagnosis, specific CFTR gene mutations, concurrent medications
<b>Age Restrictions</b>	2 years of age and older
<b>Prescriber Restrictions</b>	Prescribed by or in consultation with a pulmonologist or a physician who specializes in CF
<b>Coverage Duration</b>	1 year
<b>Other Criteria</b>	CF - Approve if the pt meets A, B and C: A) pt has at least one mutation in the CFTR gene that is considered to be pathogenic or likely pathogenic, and B) pt must have positive CF newborn screening test or family history of CF or clinical presentation consistent with signs and symptoms of CF and C) evidence of abnormal CFTR function as demonstrated by i, ii, or iii: (i) elevated sweat chloride test or (ii) two CFTR mutations or (iii) abnormal nasal potential difference.
<b>Indications</b>	All FDA-approved Indications.
<b>Off-Label Uses</b>	N/A
<b>Part B Prerequisite</b>	No

# TRUQAP

## Products Affected

- TRUQAP

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	Diagnosis
<b>Age Restrictions</b>	18 years and older
<b>Prescriber Restrictions</b>	N/A
<b>Coverage Duration</b>	1 year
<b>Other Criteria</b>	Breast Cancer-Approve if the patient meets the following (A, B, C, D and E): A) Patient has locally advanced or metastatic disease, AND B) Patient has hormone receptor positive (HR+) disease, AND C) Patient has human epidermal growth factor receptor 2 (HER2)-negative disease, AND D) Patient has at least one phosphatidylinositol 3-kinase (PIK3CA), serine/threonine protein kinase (AKT1), or phosphatase and tensin homolog (PTEN)-alteration, AND E) Patient meets one of the following (i or ii): i. Patient has had progression with at least one endocrine-based regimen in the metastatic setting (Note: Examples of endocrine therapy include anastrozole, exemestane, and letrozole.) and has had progression with at least one cyclin-dependent kinase (CDK) 4/6 inhibitor in the metastatic setting (Note: Examples of CDK4/6 inhibitor include: Ibrance (palbociclib tablets or capsules), Verzenio (abemaciclib tablets), Kisqali (ribociclib tablets), KISQALI Femara Co-Pack (ribociclib and letrozole tablets) OR ii. Patient has recurrence on or within 12 months of completing adjuvant endocrine therapy.
<b>Indications</b>	All FDA-approved Indications.
<b>Off-Label Uses</b>	N/A

PA Criteria	Criteria Details
Part B Prerequisite	No

# TUKYSA

## Products Affected

- TUKYSA ORAL TABLET 150 MG, 50 MG

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	Diagnosis, prior therapies
<b>Age Restrictions</b>	18 years and older
<b>Prescriber Restrictions</b>	N/A
<b>Coverage Duration</b>	1 year
<b>Other Criteria</b>	Breast Cancer-approve if the patient has recurrent or metastatic human epidermal growth factor receptor 2 (HER2)-positive disease, has received at least one prior anti-HER2-based regimen in the metastatic setting and Tukysa is used in combination with trastuzumab and capecitabine. Colon/Rectal Cancer-approve if the requested medication is used in combination with trastuzumab, patient has unresectable or metastatic disease, human epidermal growth factor receptor 2 (HER2)-amplified disease, AND Patient's tumor or metastases are wild-type RAS (KRAS wild-type and NRAS wild-type). Biliary tract cancer- approve if the patient meets all of (a, b, c, and d): a) unresectable or metastatic disease, b) HER2 positive disease, c) tried at least one systemic regimen, d) will use in combination with trastuzumab.
<b>Indications</b>	All FDA-approved Indications, Some Medically-accepted Indications.
<b>Off-Label Uses</b>	Biliary tract cancer
<b>Part B Prerequisite</b>	No

# TURALIO

## Products Affected

- TURALIO ORAL CAPSULE 125 MG

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	Diagnosis
<b>Age Restrictions</b>	18 years and older
<b>Prescriber Restrictions</b>	N/A
<b>Coverage Duration</b>	1 year
<b>Other Criteria</b>	Tenosynovial Giant Cell Tumor (Pigmented Villonodular Synovitis)-approve if, according to the prescriber, the tumor is not amenable to improvement with surgery. Histiocytic Neoplasms-approve if the patient has a colony stimulating factor 1 receptor (CSF1R) mutation AND has one of the following conditions (i, ii, or iii): i. Langerhans cell histiocytosis OR ii. Erdheim-Chester disease OR iii. Rosai-Dorfman disease.
<b>Indications</b>	All FDA-approved Indications, Some Medically-accepted Indications.
<b>Off-Label Uses</b>	Histiocytic Neoplasms
<b>Part B Prerequisite</b>	No

# UBRELVY

## Products Affected

- UBRELVY

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	Diagnosis
Age Restrictions	18 years and older
Prescriber Restrictions	N/A
Coverage Duration	1 year
Other Criteria	Migraine, Acute treatment-approve
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

## UPTRAVI

### Products Affected

- UPTRAVI ORAL TABLET
- UPTRAVI ORAL TABLETS,DOSE PACK

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	Concurrent Use with Other Inhaled or Parenteral Prostacyclin Agents Used for Pulmonary Hypertension.
<b>Required Medical Information</b>	Confirmation of right heart catheterization, medication history (as described in Other Criteria)
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	PAH must be prescribed by, or in consultation with, a cardiologist or a pulmonologist.
<b>Coverage Duration</b>	1 year
<b>Other Criteria</b>	Must have PAH (WHO Group 1) and had a right heart catheterization to confirm the diagnosis of PAH (WHO Group 1). Patient new to therapy must meet a) OR b): a) tried one or is currently taking one oral therapy for PAH for 30 days, unless patient has experienced treatment failure, intolerance, or oral therapy is contraindicated: PDE5 inhibitor (eg, sildenafil, Revatio), endothelin receptor antagonist (ERA) [eg, Tracleer, Letairis or Opsumit], or Adempas, OR b) receiving or has received in the past one prostacyclin therapy for PAH (eg, Orenitram, Ventavis, or epoprostenol injection).
<b>Indications</b>	All FDA-approved Indications.
<b>Off-Label Uses</b>	N/A
<b>Part B Prerequisite</b>	No



# VALCHLOR

## Products Affected

- VALCHLOR

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	Diagnosis
<b>Age Restrictions</b>	Cutaneous lymphoma-18 years and older
<b>Prescriber Restrictions</b>	N/A
<b>Coverage Duration</b>	1 year
<b>Other Criteria</b>	Cutaneous Lymphomas (Note-includes mycosis fungoides/Sezary syndrome, primary cutaneous B-cell lymphoma, primary cutaneous CD30+ T-cell lymphoproliferative disorders)-approve. Adult T-Cell Leukemia/Lymphoma-approve if the patient has chronic/smoldering subtype of adult T-cell leukemia/lymphoma. Langerhans cell histiocytosis-approve if the patient has unifocal Langerhans cell histiocytosis with isolated skin disease.
<b>Indications</b>	All FDA-approved Indications, Some Medically-accepted Indications.
<b>Off-Label Uses</b>	Adults with T-cell leukemia/lymphoma, Langerhans Cell Histiocytosis
<b>Part B Prerequisite</b>	No

# VALTOCO

## Products Affected

- VALTOCO

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	Diagnosis, other medications used at the same time
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	Prescribed by or in consultation with a neurologist
<b>Coverage Duration</b>	1 year
<b>Other Criteria</b>	Intermittent Episodes of Frequent Seizure Activity (i.e., seizure clusters, acute repetitive seizures)-approve if the patient is currently receiving maintenance antiseizure medication(s).
<b>Indications</b>	All FDA-approved Indications.
<b>Off-Label Uses</b>	N/A
<b>Part B Prerequisite</b>	No

# VANCOMYCIN

## Products Affected

- *vancomycin oral capsule 125 mg, 250 mg*

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	Diagnosis
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	2 weeks
Other Criteria	N/A
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

# VANFLYTA

## Products Affected

- VANFLYTA

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	Diagnosis
Age Restrictions	18 years and older
Prescriber Restrictions	N/A
Coverage Duration	1 year
Other Criteria	Acute Myeloid Leukemia: approve if the patient has FLT3-ITD mutation-positive disease as detected by an approved test and this medication is being used for induction, re-induction, consolidation, or maintenance treatment.
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

# VENCLEXTA

## Products Affected

- VENCLEXTA ORAL TABLET 10 MG, 100 MG, 50 MG
- VENCLEXTA STARTING PACK

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	Diagnosis, prior therapy
<b>Age Restrictions</b>	18 years and older
<b>Prescriber Restrictions</b>	N/A
<b>Coverage Duration</b>	1 year
<b>Other Criteria</b>	AML-approve if used in combination with azacitidine, decitabine, or cytarabine. CLL/SLL- approve. Mantle Cell Lymphoma- approve if the patient has tried at least one systemic regimen. Multiple Myeloma- approve if the patient has t (11,14) translocation AND has tried at least one systemic regimen for multiple myeloma AND Venclexta will be used in combination with dexamethasone. Systemic light chain amyloidosis-approve if the patient has t (11, 14) translocation and has tried at least one systemic regimen. Waldenstrom Macroglobulinemia/Lymphoplasmacytic Lymphoma-approve if the patient has tried at least one systemic regimen.
<b>Indications</b>	All FDA-approved Indications, Some Medically-accepted Indications.
<b>Off-Label Uses</b>	Mantle Cell Lymphoma, waldenstrom macroglobulinemia/lymphoplasmacytic lymphoma, multiple myeloma, systemic light chain amyloidosis
<b>Part B Prerequisite</b>	No

# VERZENIO

## Products Affected

- VERZENIO

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	Breast cancer: HR status, HER2 status, previous medications/therapies tried, concomitant therapy, menopausal status
<b>Age Restrictions</b>	18 years and older
<b>Prescriber Restrictions</b>	N/A
<b>Coverage Duration</b>	1 year
<b>Other Criteria</b>	<p>Breast Cancer, Early-Approve if pt meets (A,B,C and D): A)Pt has HR+disease, AND B)Pt has HER2-negative breast cancer, AND C)Pt has node-positive disease at high risk of recurrence AND D)Pt meets ONE of the following (i or ii): i.Verzenio will be used in combo w/anastrozole, exemestane, or letrozole AND pt meets one of the following (a,b, or c): a)Pt is a postmenopausal woman, OR b)Pt is a pre/perimenopausal woman and meets one of the following 1 or 2:1-Pt is receiving ovarian suppression/ablation with a gonadotropin-releasing hormone (GnRH) agonist, OR 2-Pt has had surgical bilateral oophorectomy or ovarian irradiation, OR c)Pt is a man and pt is receiving a GnRH analog, OR ii.Verzenio will be used in combo with tamoxifen AND pt meets one of the following (a or b): a)Pt is a postmenopausal woman or man OR b)Pt is a pre/perimenopausal woman and meets one of the following 1 or 2:1-Pt is receiving ovarian suppression/ablation with a GnRH agonist, OR 2-Patient has had surgical bilateral oophorectomy or ovarian irradiation. Breast Cancer-Recurrent or Metastatic in Women-Approve if pt meets (A, B, C and D): A) Pt has HR+ disease, AND B)Pt has HER2-negative breast cancer, AND C)Pt meets ONE of the following criteria (i or ii): i.Pt is a postmenopausal woman, OR ii.Pt is a pre/perimenopausal woman and meets one of the following (a or b): a)Pt is receiving ovarian</p>

PA Criteria	Criteria Details
	<p>suppression/ablation with a GnRH agonist, OR b)Pt has had surgical bilateral oophorectomy or ovarian irradiation, AND D)Pt meets ONE of the following criteria (i, ii, or iii): i.Verzenio will be used in combo with anastrozole, exemestane, or letrozole, OR ii.Verzenio will be used in combo with fulvestrant, OR iii.pt meets the following conditions (a, b, and c): a)Verzenio will be used as monotherapy, AND b)Pt's breast cancer has progressed on at least one prior endocrine therapy, AND c)Pt has tried chemotherapy for metastatic breast cancer.Breast Cancer-Recurrent or Metastatic in Men-Approve if pt meets the following criteria (A,B and C): A)Pt has HR+ disease, AND B)Pt has HER2-negative breast cancer, AND C)Pt meets ONE of the following criteria (i, ii, or iii): i.Pt meets BOTH of the following conditions (a and b): a)Pt is receiving a GnRH analog, AND b)Verzenio will be used in combo with anastrozole, exemestane, or letrozole, OR ii.Verzenio will be used in combo with fulvestrant, OR iii.Pt meets the following conditions (a, b, and c): a)Verzenio will be used as monotherapy, AND b)Pt's breast cancer has progressed on at least one prior endocrine therapy, AND c)Pt has tried chemotherapy for metastatic breast cancer. Endometrial cancer- approve if pt meets all of (A, B, And C): A) pt has recurrent or metastatic disease, and B) pt has estrogen receptor (ER)-positive tumors, and C) pt will be using in combination with letrozole.</p>
<b>Indications</b>	All FDA-approved Indications, Some Medically-accepted Indications.
<b>Off-Label Uses</b>	Endometrial cancer
<b>Part B Prerequisite</b>	No

# VIGABATRIN

## Products Affected

- *vigabatin*
- *vigadrone*
- *vigpoder*

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	Diagnosis, medication history (complex partial seizures)
<b>Age Restrictions</b>	Refractory complex partial seizures - patients 2 years of age or older. Infantile spasms - patients less than or equal to 2 years of age
<b>Prescriber Restrictions</b>	Prescribed by, or in consultation with, a neurologist
<b>Coverage Duration</b>	Infantile spasms- 6 months. Treatment-Refractory Partial Seizures- initial 3 months, cont 1 year
<b>Other Criteria</b>	Infantile spasms-requested medication is being used as monotherapy. Treatment refractory complex partial seizures intial-patient has tried and/or is concomitantly receiving at least two other antiepileptic drugs. Treatment refractory complex partial seizures continuation- the patient is responding to therapy (e.g., reduced seizure severity, frequency, and/or duration).
<b>Indications</b>	All FDA-approved Indications.
<b>Off-Label Uses</b>	N/A
<b>Part B Prerequisite</b>	No



# VITRAKVI

## Products Affected

- VITRAKVI ORAL CAPSULE 100 MG, 25 MG
- VITRAKVI ORAL SOLUTION

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	Diagnosis, NTRK gene fusion status
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	1 year
Other Criteria	Solid tumors - approve if the tumor is positive for neurotrophic receptor tyrosine kinase (NTRK) gene fusion AND the tumor is metastatic or surgical resection of tumor will likely result in severe morbidity.
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

# VIZIMPRO

## Products Affected

- VIZIMPRO

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	Diagnosis, EGFR status, exon deletions or substitutions
<b>Age Restrictions</b>	18 years and older
<b>Prescriber Restrictions</b>	N/A
<b>Coverage Duration</b>	1 year
<b>Other Criteria</b>	NSCLC-approve if the patient has advanced or metastatic disease, has sensitizing EGFR mutation-positive NSCLC as detected by an approved test. Note: Examples of sensitizing EGFR mutation-positive NSCLC include the following mutations: exon 19 deletions, exon 21 (L858R) substitution mutations, L861Q, G719X and S7681.
<b>Indications</b>	All FDA-approved Indications.
<b>Off-Label Uses</b>	N/A
<b>Part B Prerequisite</b>	No

# VONJO

## Products Affected

- VONJO

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	Diagnosis
<b>Age Restrictions</b>	18 years and older
<b>Prescriber Restrictions</b>	N/A
<b>Coverage Duration</b>	1 year
<b>Other Criteria</b>	Myelofibrosis (MF), including primary MF, post-polycythemia Vera MF, and Post-Essential Thrombocythemia MF-approve if the patient meets either (A, B, or C): (A) the patient has a platelet count of less than 50 X 10 <sup>9</sup> /L (less than 50,000/mcL) and meets one of the following criteria (a or b):a) Patient has intermediate-risk or high-risk disease and is not a candidate for transplant, or b) Patient has lower-risk disease OR (B) Patient has a platelet count of greater than or equal to 50 X 10 <sup>9</sup> /L (greater than or equal to 50,000/mcL) and meets all of the following criteria (a, b and c): a) Patient has high-risk disease, AND b) Patient is not a candidate for transplant, AND c) Patient has tried Jakafi (ruxolitinib tablets) or Inrebic (fedratinib capsules) OR (C) patient has myelofibrosis-associated anemia with symptomatic splenomegaly and/or constitutional symptoms.
<b>Indications</b>	All FDA-approved Indications.
<b>Off-Label Uses</b>	N/A
<b>Part B Prerequisite</b>	No

## VORICONAZOLE (ORAL)

### Products Affected

- voriconazole*

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	Diagnosis
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	N/A
<b>Coverage Duration</b>	Aspergillus-Prophy, systemic w/risk neutropenia-Prophy, systemic w/HIV-Prophy/Tx-6 mo, others-3 mo
<b>Other Criteria</b>	N/A
<b>Indications</b>	All FDA-approved Indications, Some Medically-accepted Indications.
<b>Off-Label Uses</b>	Aspergillus Infections - prophylaxis, oropharyngeal candidiasis (fluconazole-refractory) - treatment, candidia endophthalmitis - treatment, blastomycosis - treatment, fungal infections (systemic) in patients at risk of neutropenia - prophylaxis, fungal infections (systemic) in patients with human immunodeficiency virus (HIV) - prophylaxis or treatment.
<b>Part B Prerequisite</b>	No

# VOSEVI

## Products Affected

- VOSEVI

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	Diagnosis
<b>Age Restrictions</b>	18 years or older
<b>Prescriber Restrictions</b>	Prescribed by or in consultation with a gastroenterologist, hepatologist, infectious diseases physician, or a liver transplant physician
<b>Coverage Duration</b>	Will be c/w AASLD guidance and inclusive of treatment already received for the requested drug
<b>Other Criteria</b>	Criteria will be applied consistent with current AASLD/IDSA guidance.
<b>Indications</b>	All FDA-approved Indications, Some Medically-accepted Indications.
<b>Off-Label Uses</b>	Indications consistent with current AASLD/IDSA guidance
<b>Part B Prerequisite</b>	No

# VOTRIENT

## Products Affected

- *pazopanib*

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	N/A
<b>Age Restrictions</b>	18 years and older
<b>Prescriber Restrictions</b>	N/A
<b>Coverage Duration</b>	1 year
<b>Other Criteria</b>	<p>Soft tissue sarcoma other than GIST-approve if the patient has advanced or metastatic disease and has ONE of the following: alveolar soft part sarcoma, angiosarcoma, desmoid tumors (aggressive fibromatosis, dermatofibrosarcoma protuberans with fibrosarcomatous transformation, non-adipocytic sarcoma or pleomorphic rhabdomyosarcoma.</p> <p>Differentiated (ie, papillary, follicular, Hurthle) thyroid carcinoma, approve if the patient is refractory to radioactive iodine therapy. Uterine sarcoma, approve if the patient has recurrent or metastatic disease. Renal Cell Carcinoma, Clear Cell or non-Clear Cell histology-approved if the patient has relapsed or advanced disease or VonHippel-Lindau disease. Ovarian Cancer (ie, Epithelial Ovarian, Fallopian Tube, or Primary Peritoneal Cancer) - approve if the patient has persistent or recurrent disease. GIST - approve if the patient has succinate dehydrogenase (SDH)-deficient GIST OR the patient has tried TWO of the following: Gleevec, Ayvakit, Sutent, Sprycel, Qinlock or Stivarga. Medullary Thyroid Carcinoma, approve if the patient has tried at least one systemic therapy. Bone cancer-approve if the patient has chondrosarcoma and has metastatic widespread disease.</p>
<b>Indications</b>	All FDA-approved Indications, Some Medically-accepted Indications.

PA Criteria	Criteria Details
<b>Off-Label Uses</b>	Differentiated (ie, papillary, follicular, oncocytic carcinoma) thyroid carcinoma. Uterine sarcoma, Epithelial Ovarian, Fallopian Tube, or Primary Peritoneal Cancer, Gastrointestinal Stromal Tumor (GIST), Medullary thyroid carcinoma, bone cancer.
<b>Part B Prerequisite</b>	No

# VOWST

## Products Affected

- VOWST

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	Diagnosis
Age Restrictions	18 years and older
Prescriber Restrictions	N/A
Coverage Duration	30 days
Other Criteria	Prevention of recurrence of clostridioides difficile infection (CDI)-approve if the patient has completed a bowel prep, will not eat or drink for at least 8 hours prior to the first dose and will complete their antibacterial treatment for recurrent CDI 2-4 days before initiating treatment with Vowst and Vowst will not be used for the TREATMENT of CDI.
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Part B Prerequisite	No



# VUMERITY

## Products Affected

- VUMERITY

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	Concurrent use with other disease-modifying agents used for multiple sclerosis (MS)
<b>Required Medical Information</b>	Relapsing form of Multiple Sclerosis (MS), to include clinically-isolated syndrome, relapsing-remitting disease, and active secondary progressive disease
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	Prescribed by or in consultation with a neurologist or a physician who specializes in the treatment of MS.
<b>Coverage Duration</b>	Authorization will be for 1 year.
<b>Other Criteria</b>	N/A
<b>Indications</b>	All FDA-approved Indications.
<b>Off-Label Uses</b>	N/A
<b>Part B Prerequisite</b>	No

# WELIREG

## Products Affected

- WELIREG

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	Diagnosis
<b>Age Restrictions</b>	18 years and older
<b>Prescriber Restrictions</b>	N/A
<b>Coverage Duration</b>	1 year
<b>Other Criteria</b>	Renal Cell Carcinoma- approve if pt has advanced disease AND has tried at least one programmed death receptor-1 (PD-1) or programmed death-ligand 1 (PD-L1) inhibitor AND has tried at least one a vascular endothelial growth factor tyrosine kinase inhibitor (VEGF-TKI). [Note: Examples of PD-1 inhibitor or PD-L1 inhibitor include: Keytruda (pembrolizumab intravenous infusion), Opdivo (nivolumab intravenous infusion), and Bavencio (avelumab intravenous infusion). Examples of VEGF-TKI include Cabometyx (cabozantinib tablets), Lenvima (lenvatinib capsules), Inlyta (axitinib tablets), Fotivda (tivozanib capsules), pazopanib, sunitinib, and sorafenib.] Van Hippel-Lindau Disease-approve if the patient meets the following (A, B, and C): A) Patient has a von Hippel-Lindau (VHL) germline alteration as detected by genetic testing, B) Does not require immediate surgery and C) Patient requires therapy for ONE of the following conditions (i, ii, iii, or iv): i. Central nervous system hemangioblastomas, OR ii. Pancreatic neuroendocrine tumors, OR iii. Renal cell carcinoma, OR iv. Retinal hemangioblastoma.
<b>Indications</b>	All FDA-approved Indications.
<b>Off-Label Uses</b>	N/A

PA Criteria	Criteria Details
Part B Prerequisite	No

# XALKORI

## Products Affected

- XALKORI ORAL CAPSULE
- XALKORI ORAL PELLETT 150 MG, 20 MG, 50 MG

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	Diagnosis
<b>Age Restrictions</b>	Anaplastic large cell lymphoma/IMT-patients greater than or equal to 1 year of age. All other diagnoses-18 years and older
<b>Prescriber Restrictions</b>	N/A
<b>Coverage Duration</b>	1 year
<b>Other Criteria</b>	Metastatic non-small cell lung cancer-approve if the patient has anaplastic lymphoma kinase (ALK)-positive disease, as detected by an approved test and patients new to therapy must have a trial of Alecensa prior to approval of Xalkori. Metastatic non-small cell lung cancer, approve if the patient has ROS1 rearrangement positive disease, as detected by an approved test. Anaplastic Large Cell Lymphoma-approve if the patient has anaplastic lymphoma kinase (ALK)-positive disease AND (i or ii): (i) the medication is used for palliative-intent therapy, or (ii) pt has relapsed or refractory disease. Histiocytic neoplasm-approve if the patient has ALK rearrangement/fusion-positive disease and meets one of the following criteria (i, ii, or iii): (i. Patient has Langerhans cell histiocytosis, OR ii. Patient has Erdheim-Chester disease OR iii. Patient has Rosai-Dorfman disease. NSCLC with MET mutation-approve if the patient has high level MET amplification or MET exon 14 skipping mutation. Inflammatory Myofibroblastic Tumor-approve if the patient has ALK positive disease and the patient has advanced, recurrent or metastatic disease or the tumor is inoperable. Melanoma, cutaneous-approve if the patient has ALK fusion disease or ROS1 fusion disease.

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Indications</b>	All FDA-approved Indications, Some Medically-accepted Indications.
<b>Off-Label Uses</b>	NSCLC with high level MET amplification or MET Exon 14 skipping mutation, Histiocytic neoplasms, melanoma, cutaneous.
<b>Part B Prerequisite</b>	No

# XDEMVY

## Products Affected

- XDEMVY

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	Diagnosis
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	6 months
Other Criteria	N/A
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

# XELJANZ

## Products Affected

- XELJANZ ORAL SOLUTION
- XELJANZ ORAL TABLET
- XELJANZ XR

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	Concurrent use with a biologic or with a Targeted Synthetic DMARD for an inflammatory condition (eg, tocilizumab, anakinra, abatacept, rituximab, certolizumab pegol, etanercept, adalimumab, infliximab, golimumab). Concurrent use with potent immunosuppressants that are not methotrexate (MTX) [eg, azathioprine, tacrolimus, cyclosporine, mycophenolate mofetil].
<b>Required Medical Information</b>	Diagnosis, concurrent medications, previous drugs tried.
<b>Age Restrictions</b>	AS/PsA/RA/UC-18 years and older (initial therapy)
<b>Prescriber Restrictions</b>	RA, JIA/JRA/AS-prescribed by or in consultation with a rheumatologist. PsA-prescribed by or in consultation with a rheumatologist or a dermatologist. UC-prescribed by or in consultation with a gastroenterologist.
<b>Coverage Duration</b>	Approve through end of plan year
<b>Other Criteria</b>	RA initial-approve Xeljanz/XR tablets if the patient has had a 3 month trial of at least one tumor necrosis factor inhibitor or was unable to tolerate a 3 month trial. PsA initial, approve Xeljanz/XR tablets if the patient has had a 3 month trial of at least one tumor necrosis factor inhibitor or was unable to tolerate a 3 month trial and the requested medication will be used in combination with methotrexate or another conventional synthetic disease modifying antirheumatic drug (DMARD), unless contraindicated. UC- Approve Xeljanz/XR tablets if the patient has had a 3 month trial of at least ONE tumor necrosis factor inhibitor for ulcerative colitis or was unable to tolerate a 3-month trial. Juvenile Idiopathic Arthritis (JIA) [or Juvenile Rheumatoid Arthritis] (regardless of type of onset) [Note: This includes patients with juvenile spondyloarthritis/active sacroiliac arthritis]-initial-approve Xeljanz immediate release tablets or solution if the patient meets the following: patient has had a 3 month trial of at least one tumor

PA Criteria	Criteria Details
	necrosis factor inhibitor or was unable to tolerate a 3 month trial. AS-approve Xeljanz/XR tablets if the patient has had a 3 month trial of at least one tumor necrosis factor inhibitor or was unable to tolerate a 3 month trial. Continuation Therapy - Patient must have responded, as determined by the prescriber.
<b>Indications</b>	All FDA-approved Indications.
<b>Off-Label Uses</b>	N/A
<b>Part B Prerequisite</b>	No



# XERMELO

## Products Affected

- XERMELO

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	Diagnosis, previous therapy, concomitant therapy
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	N/A
<b>Coverage Duration</b>	1 year
<b>Other Criteria</b>	Initial therapy - approve if the patient meets ALL of the following criteria: 1) patient has been on long-acting somatostatin analog (SSA) therapy (eg, Somatuline Depot [lanreotide for injection]) AND 2) while on long-acting SSA therapy (prior to starting Xermelo), the patient continues to have at least four bowel movements per day, AND 3) Xermelo will be used concomitantly with a long-acting SSA therapy. Continuation therapy - approve if the patient is continuing to take Xermelo concomitantly with a long-acting SSA therapy for carcinoid syndrome diarrhea.
<b>Indications</b>	All FDA-approved Indications.
<b>Off-Label Uses</b>	N/A
<b>Part B Prerequisite</b>	No

# XIFAXAN

## Products Affected

- XIFAXAN ORAL TABLET 200 MG, 550 MG

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	Diagnosis
<b>Age Restrictions</b>	Traveler's diarrhea - 12 years of age or older. Hepatic encephalopathy, irritable bowel syndrome with diarrhea - 18 years of age or older.
<b>Prescriber Restrictions</b>	N/A
<b>Coverage Duration</b>	Hep Enceph-6mo, IBS/diarrhea-14days, Traveler's Diarr-3days, Sm intest bacterial overgrowth-14days
<b>Other Criteria</b>	Hepatic Encephalopathy-approve Xifaxan 550 mg tablets if the patient has previously had overt hepatic encephalopathy and the requested medication will be used concomitantly with lactulose, unless the patient has a contraindication or significant intolerance to treatment with lactulose. Irritable bowel syndrome with diarrhea-approve Xifaxan 550 mg tablets. Travelers Diarrhea-approve Xifaxan 200 mg tablets if the patient is afebrile and does not have blood in the stool. Small intestine bacterial overgrowth-approve Xifaxan 200mg or 550 mg tablets if the diagnosis has been confirmed by a glucose hydrogen breath test, lactulose hydrogen breath test, or small bowel aspiration and culture.
<b>Indications</b>	All FDA-approved Indications, Some Medically-accepted Indications.
<b>Off-Label Uses</b>	Small intestine bacterial overgrowth
<b>Part B Prerequisite</b>	No

# XOLAIR

## Products Affected

- XOLAIR SUBCUTANEOUS AUTO-INJECTOR 150 MG/ML, 300 MG/2 ML, 75 MG/0.5 ML
- XOLAIR SUBCUTANEOUS RECON SOLN
- XOLAIR SUBCUTANEOUS SYRINGE 150 MG/ML, 300 MG/2 ML, 75 MG/0.5 ML

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	Concurrent use with another monoclonal antibody therapy.
<b>Required Medical Information</b>	Diagnosis
<b>Age Restrictions</b>	Moderate to severe persistent asthma-6 years and older. CIU-12 years and older. Polyps-18 years and older. Food Allergy-1 yr and older
<b>Prescriber Restrictions</b>	Moderate to severe persistent asthma if prescribed by, or in consultation with an allergist, immunologist, or pulmonologist. CIU if prescribed by or in consultation with an allergist, immunologist, or dermatologist. Polyps-prescribed by or in consult with an allergist, immunologist, or otolaryngologist. Food allergy- allergist or immunologist
<b>Coverage Duration</b>	asthma/CIU-Initial tx 4 months, Polyps-initial-6 months, continued tx 12 months, Food allergy-1 yr
<b>Other Criteria</b>	MODERATE TO SEVERE PERSISTENT ASTHMA (A, B, C and D): A) baseline IgE greater than or equal to 30 IU/mL, and B) baseline positive skin test or in vitro test for 1 or more perennial or seasonal aeroallergens C) received at least 3 months of combination therapy with an inhaled corticosteroid (ICS) and additional asthma controller/maintenance medication (e.g., LABA, LAMA, leukotriene receptor antagonist, monoclonal antibody) [see Exception 1 below] and D) asthma is uncontrolled or was uncontrolled prior to receiving Xolair or another monoclonal antibody and meets one of the following (a, b, c, d, or e): a) experienced two or more asthma exacerbations requiring systemic CSs in the past year, b) experienced one or more asthma exacerbation requiring hospitalization/urgent care visit/emergency department visit in the past year, c) forced expiratory volume in 1 second (FEV1) less than 80%

PA Criteria	Criteria Details
	<p>predicted, d) FEV1/forced vital capacity (FVC) less than 0.80, or e) asthma worsens upon tapering of oral CS. CHRONIC RHINOSINUSITIS WITH NASAL POLYPS (CRwNP) [all of A, B, C, D, and E]: A) diagnosis by direct exam, endoscopy, or sinus CT scan, B) baseline (prior to Xolair or another monoclonal antibody that may lower IgE) IgE at least 30 IU/ml, C) at least two of the following symptoms for 6 months: nasal congestion, obstruction, discharge, reduction/loss of smell, D) tried intranasal CS and will continue in combination with Xolair, and E) one of the following (a, b, or c): a) had systemic CS at least 5 days in past 2 years, b) contraindication to systemic CS, or c) had nasal polyp surgery. CHRONIC IDIOPATHIC URTICARIA (CIU): urticaria more than 6 weeks prior to treatment with Xolair with symptoms present more than 3 days per week despite daily non-sedating H1-antihistamine therapy. IgE-MEDIATED FOOD ALLERGY (all of A, B, C and D): A) baseline IgE greater than or equal to 30 IU/mL, B) positive skin prick test and positive in vitro test for IgE to one or more foods, C) history of allergic reaction that met all of the following (a, b, and c): a) signs and symptoms of a significant systemic allergic reaction, b) reaction occurred within a short period of time following a known ingestion of the food, and c) prescriber deemed this reaction significant enough to require a prescription for an epinephrine auto-injector, and D) patient has been prescribed an epinephrine auto-injector. CONTINUATION THERAPY: ASTHMA: patient responded to therapy and continues to receive an ICS. CRwNP: patient responded after 6 months of therapy and continues intranasal CS. CIU: patient responded to therapy. Exception 1: an exception to the requirement of a trial of one additional asthma controller/maintenance medication can be made if the patient has already received anti-IL-4/13 therapy (Dupixent) used concomitantly with an ICS.</p>
<b>Indications</b>	All FDA-approved Indications.
<b>Off-Label Uses</b>	N/A
<b>Part B Prerequisite</b>	No

# XOSPATA

## Products Affected

- XOSPATA

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	Diagnosis, FLT3-mutation status
<b>Age Restrictions</b>	18 years and older
<b>Prescriber Restrictions</b>	N/A
<b>Coverage Duration</b>	1 year
<b>Other Criteria</b>	AML - approve if the patient has relapsed or refractory disease AND the disease is FLT3-mutation positive as detected by an approved test. Lymphoid, Myeloid Neoplasms-approve if the patient has eosinophilia and the disease is FLT3-mutation positive as detected by an approved test.
<b>Indications</b>	All FDA-approved Indications, Some Medically-accepted Indications.
<b>Off-Label Uses</b>	Lymphoid, Myeloid Neoplasms
<b>Part B Prerequisite</b>	No

# XPOVIO

## Products Affected

- XPOVIO

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	Diagnosis, prior therapies
<b>Age Restrictions</b>	18 years and older
<b>Prescriber Restrictions</b>	N/A
<b>Coverage Duration</b>	1 year
<b>Other Criteria</b>	Multiple Myeloma-Approve if the patient meets the following (A and B): A) The medication will be taken in combination with dexamethasone AND B) Patient meets one of the following (i, ii, or iii): i. Patient has tried at least four prior regimens for multiple myeloma OR ii. Patient meets both of the following (a and b): a) Patient has tried at least one prior regimen for multiple myeloma AND b) The medication will be taken in combination with bortezomib OR iii. Patient meets both of the following (a and b): a) Patient has tried at least one prior regimen for multiple myeloma AND b) The medication will be taken in combination with Darzalex (daratumumab infusion), Darzlaex Faspro (daratumumab and hyaluronidase-fihj injection), or Pomalyst (pomalidomide capsules). Note: Examples include bortezomib/Revlimid (lenalidomide capsules)/dexamethasone, Kyprolis (carfilzomib infusion)/Revlimid/dexamethasone, Darzalex (daratumumab injection)/bortezomib or Kyprolis/dexamethasone, or other regimens containing a proteasome inhibitor, immunomodulatory drug, and/or anti-CD38 monoclonal antibody. Diffuse large B-cell lymphoma Note:this includes patients with histologic transformation of indolent lymphomas to diffuse large B-cell lymphoma)-approve if the patient has been treated with at least two prior systemic therapies.
<b>Indications</b>	All FDA-approved Indications, Some Medically-accepted Indications.

PA Criteria	Criteria Details
Off-Label Uses	Treatment of multiple myeloma in combination with daratumumb or pomalidomide
Part B Prerequisite	No

# XTANDI

## Products Affected

- XTANDI ORAL CAPSULE
- XTANDI ORAL TABLET 40 MG, 80 MG

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	Diagnosis for which Xtandi is being used.
<b>Age Restrictions</b>	18 years and older
<b>Prescriber Restrictions</b>	N/A
<b>Coverage Duration</b>	Authorization will be for 1 year
<b>Other Criteria</b>	Prostate cancer-castration-resistant [Metastatic or Non-metastatic] and Prostate cancer-metastatic, castration sensitive-approve if Xtandi will be used concurrently with a gonadotropin-releasing hormone (GnRH) analog [for example: leuprolide acetate, Lupron Depot (leuprolide acetate intramuscular injection), Trelstar (triptorelin pamoate intramuscular injection), Zoladex (goserelin acetate subcutaneous implant), Vantas (histrelin acetate subcutaneous implant), Firmagon (degarelix subcutaneous injection), Orgovyx (relugolix tablets)] or if the patient has had a bilateral orchiectomy. Prostate cancer- Non-Metastatic, Castration-Sensitive - approve if pt has biochemical recurrence and is at high risk for metastasis. [Note: High-risk biochemical recurrence is defined as prostate-specific antigen (PSA) doubling time less than or equal to 9 months.]
<b>Indications</b>	All FDA-approved Indications.
<b>Off-Label Uses</b>	N/A
<b>Part B Prerequisite</b>	No



# XYREM

## Products Affected

- SODIUM OXYBATE (PREFERRED NDCS STARTING WITH 00054)

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	Concomitant use with Xywav, Wakix or Sunosi
<b>Required Medical Information</b>	Medication history (as described in Other Criteria field)
<b>Age Restrictions</b>	7 years and older
<b>Prescriber Restrictions</b>	Prescribed by a sleep specialist physician or a Neurologist
<b>Coverage Duration</b>	12 months.
<b>Other Criteria</b>	For Excessive daytime sleepiness (EDS) in patients with narcolepsy, 18 years and older - approve if the patient has tried one CNS stimulant (e.g., methylphenidate, dextroamphetamine), modafinil, or armodafinil and narcolepsy has been confirmed with polysomnography and a multiple sleep latency test (MSLT). For EDS in patients with narcolepsy, less than 18 years old-approve if the patient has tried one CNS stimulant (e.g., methylphenidate, dextramphetamine) or modafinil and narcolepsy has been confirmed with polysomnography and a multiple sleep latency test (MSLT). Cataplexy treatment in patients with narcolepsy-approve if narcolepsy has been confirmed with polysomnography and a multiple sleep latency test (MSLT).
<b>Indications</b>	All FDA-approved Indications.
<b>Off-Label Uses</b>	N/A
<b>Part B Prerequisite</b>	No

# ZEJULA

## Products Affected

- ZEJULA ORAL TABLET

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	N/A
<b>Age Restrictions</b>	18 years and older
<b>Prescriber Restrictions</b>	N/A
<b>Coverage Duration</b>	1 year
<b>Other Criteria</b>	Ovarian, fallopian tube, or primary peritoneal cancer, maintenance therapy - approve if the patient is in complete or partial response after platinum-based chemotherapy regimen and if the patient is new to therapy they must have a trial of Lynparza prior to approval of Zejula. Patients who have had a complete or partial response to first-line platinum based chemotherapy and do not have BRCA altered disease are not required to try Lynparza. Uterine leiomyosarcoma-approve if the patient has BRCA2-altered disease and has tried one systemic regimen. In addition, patients new to therapy must have a trial of Lynparza prior to approval of Zejula. Ovarian, fallopian tube or primary peritoneal cancer in the treatment setting-approve.
<b>Indications</b>	All FDA-approved Indications, Some Medically-accepted Indications.
<b>Off-Label Uses</b>	Uterine Leiomyosarcoma, Ovarian, fallopian tube or primary peritoneal cancer-treatment
<b>Part B Prerequisite</b>	No

# ZELBORAF

## Products Affected

- ZELBORAF

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	BRAFV600 mutation status required.
<b>Age Restrictions</b>	All diagnoses (except CNS cancer)-18 years and older
<b>Prescriber Restrictions</b>	N/A
<b>Coverage Duration</b>	Authorization will be for 1 year
<b>Other Criteria</b>	Melanoma, patient new to therapy must have BRAFV600 mutation for approval AND have unresectable, advanced or metastatic melanoma. HCL - must have tried at least one other systemic therapy for hairy cell leukemia OR is unable to tolerate purine analogs and Zelboraf will be used in combination with Gazyva (obinutuzumab intravenous infusion) as initial therapy. Thyroid Cancer-patient has disease that is refractory to radioactive iodine therapy. Erdheim-Chester disease, in patients with BRAF V600 mutation-approve. Central Nervous System Cancer-approve if the patient has BRAF V600 mutation-positive disease AND medication is being used for one of the following situations (i, ii, or iii): i. Adjuvant treatment of one of the following conditions (a, b, or c): a) Pilocytic astrocytoma OR b) Pleomorphic xanthoastrocytoma OR c) Ganglioglioma OR ii. Recurrent or progressive disease for one of the following conditions (a or b): a) glioma OR b) Glioblastoma OR iii. Melanoma with brain metastases AND the medication with be taken in combination with Cotellic (cobimetinib tablets). Histiocytic Neoplasm-approve if the patient has Langerhans cell histiocytosis and one of the following (i, ii, or iii): i. Multisystem disease OR ii. Pulmonary disease OR iii. Central nervous system lesions AND the patient has BRAF V600-mutation positive disease.
<b>Indications</b>	All FDA-approved Indications, Some Medically-accepted Indications.

PA Criteria	Criteria Details
<b>Off-Label Uses</b>	Patients with Hairy Cell Leukemia, Non-Small Cell Lung Cancer (NSCLC) with BRAF V600E Mutation, Differentiated thyroid carcinoma (i.e., papillary, follicular, or oncocytic carcinoma) with BRAF-positive disease, Central Nervous System Cancer, Histiocytic Neoplasm
<b>Part B Prerequisite</b>	No

# ZEPOSIA

## Products Affected

- ZEPOSIA
- ZEPOSIA STARTER KIT (28-DAY)
- ZEPOSIA STARTER PACK (7-DAY)

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	MS-Concurrent use with other disease-modifying agents used for multiple sclerosis. UC- Concurrent Use with a Biologic or with a Targeted Synthetic Disease-modifying Antirheumatic Drug (DMARD) for Ulcerative Colitis
<b>Required Medical Information</b>	Diagnosis
<b>Age Restrictions</b>	UC-18 years and older
<b>Prescriber Restrictions</b>	MS-Prescribed by or in consultation with a neurologist or a physician who specializes in the treatment of multiple sclerosis. UC-Prescribed by or in consultation with a gastroenterologist
<b>Coverage Duration</b>	1 year
<b>Other Criteria</b>	MS-approve. Ulcerative Colitis, initial-approve if the patient has tried TWO of the following: a preferred adalimumab product, Stelara, a preferred infliximab product, Rinvoq. Note-a trial of Simponi SC, Entyvio IV/SC, Omvoh IV/SC, a non-preferred adalimumab product or a non-preferred infliximab would also count). Cont tx-approve if the patient has been established on Zeposia. Please Note: preferred adalimumab products include Humira (NDCs starting with -00074), Cyltezo, Yuflyma. Preferred infliximab products include Remicade, Zymfentra.
<b>Indications</b>	All FDA-approved Indications.
<b>Off-Label Uses</b>	N/A
<b>Part B Prerequisite</b>	No

# ZOLINZA

## Products Affected

- ZOLINZA

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	Diagnosis
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	1 year
Other Criteria	Cutaneous T-Cell Lymphoma including Mycosis Fungoides/Sezary Syndrome-approve.
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

# ZTALMY

## Products Affected

- ZTALMY

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	Diagnosis
Age Restrictions	2 years and older
Prescriber Restrictions	Prescribed by or in consultation with a neurologist
Coverage Duration	1 year
Other Criteria	Seizures associated with cyclin-dependent kinase-like 5 (CDKL5) deficiency disorder-approve if the patient has a molecularly confirmed pathogenic or likely pathogenic mutation in the CDKL5 gene and patient has tried or is concomitantly receiving two other antiepileptic drugs.
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

# ZURZUVAE

## Products Affected

- ZURZUVAE ORAL CAPSULE 20 MG, 25 MG, 30 MG

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	Previous treatment with Zurzuvae during the current episode of postpartum depression
<b>Required Medical Information</b>	Diagnosis
<b>Age Restrictions</b>	18 years and older
<b>Prescriber Restrictions</b>	Prescribed by or in consultation with a psychiatrist or an obstetrician-gynecologist
<b>Coverage Duration</b>	14 days
<b>Other Criteria</b>	Postpartum depression-approve if the patient meets the following (A, B and C): A.Patient meets BOTH of the following (i and ii): i. Patient has been diagnosed with severe depression, AND ii. Symptom onset began during the third trimester of pregnancy or up to 4 weeks post-delivery, AND B. Patient is less than or equal to 12 months postpartum, AND C. Patient is not currently pregnant.
<b>Indications</b>	All FDA-approved Indications.
<b>Off-Label Uses</b>	N/A
<b>Part B Prerequisite</b>	No



# ZYDELIG

## Products Affected

- ZYDELIG

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	N/A
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Authorization will be for 1 year
Other Criteria	CLL/SLL-approve if the patient has tried at least one systemic regimen.
Indications	All FDA-approved Indications, Some Medically-accepted Indications.
Off-Label Uses	small lymphocytic lymphoma
Part B Prerequisite	No

# ZYKADIA

## Products Affected

- ZYKADIA

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	Diagnosis
<b>Age Restrictions</b>	18 years and older
<b>Prescriber Restrictions</b>	N/A
<b>Coverage Duration</b>	Authorization will be for 1 year
<b>Other Criteria</b>	Erdheim-Chester Disease-approve if the patient has anaplastic lymphoma kinase (ALK) rearrangement/fusion-positive disease. NSCLC, ALK positive-approve if the patient has advanced or metastatic disease that is ALK positive as detected by an approved test and for patients new to therapy must have a trial of Alecensa prior to approval of Zykadia. NSCLC, ROS1 Rearrangement-approve if the patient has advanced or metastatic disease.
<b>Indications</b>	All FDA-approved Indications, Some Medically-accepted Indications.
<b>Off-Label Uses</b>	Inflammatory Myofibroblastic Tumor (IMT) with ALK Translocation. Patients with NSCLC with ROS1 Rearrangement. Erdheim-Chester disease.
<b>Part B Prerequisite</b>	No

# ZYMFENTRA

## Products Affected

- ZYMFENTRA

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	Concurrent use with a biologic or with a targeted synthetic disease-modifying antirheumatic drug (DMARD).
<b>Required Medical Information</b>	Diagnosis
<b>Age Restrictions</b>	18 years and older (initial therapy)
<b>Prescriber Restrictions</b>	Prescribed by or in consultation with a gastroenterologist (initial therapy)
<b>Coverage Duration</b>	Approve through end of plan year
<b>Other Criteria</b>	<p>Crohn's Disease, initial therapy-Approve if the patient meets the following (i. and ii.): i.The patient is currently receiving infliximab intravenous maintenance therapy or will receive induction dosing with an infliximab intravenous product within 3 months of initiating therapy with Zymfentra, AND ii. Patient meets ONE of the following (a, b, c, or d): a) Patient has tried or is currently taking systemic corticosteroids, or corticosteroids are contraindicated in this patient, Note: Examples of corticosteroids are prednisone and methylprednisolone. OR b) Patient has tried one conventional systemic therapy for Crohn's disease, Note: Examples of conventional systemic therapy for Crohn's disease include azathioprine, 6-mercaptopurine, or methotrexate. An exception to the requirement for a trial of or contraindication to steroids or a trial of one other conventional systemic agent can be made if the patient has already tried at least one biologic other than the requested medication. A biosimilar of the requested biologic does not count. OR c) Patient has enterocutaneous (perianal or abdominal) or rectovaginal fistulas, OR d) Patient had ileocolonic resection (to reduce the chance of Crohn's disease recurrence). Crohn's Disease, continuation-approve if the patient has had a response to therapy.</p> <p>Ulcerative Colitis, initial therapy-Approve if the patient meets ALL of the following (i, ii, iii, and iv): i.The patient is currently receiving infliximab</p>

PA Criteria	Criteria Details
	<p>intravenous maintenance therapy or will receive induction dosing with an infliximab intravenous product within 3 months of initiating therapy with Zymfentra, AND ii. Patient meets ONE of the following (a or b): a) Patient had a trial of one systemic agent or was intolerant to one of these agents for ulcerative colitis, Note: Examples include 6-mercaptopurine, azathioprine, cyclosporine, tacrolimus, or a corticosteroid such as prednisone or methylprednisolone. A trial of a mesalamine product does not count as a systemic therapy for ulcerative colitis. A previous trial of one biologic other than the requested medication also counts as a trial of one systemic agent for ulcerative colitis. A biosimilar of the requested biologic does not count. OR b) Patient meets BOTH of the following [(1) and (2)]: (1) Patient has pouchitis AND (2) Patient has tried therapy with an antibiotic, probiotic, corticosteroid enema, or Rowasa (mesalamine enema). Note: Examples of antibiotics include metronidazole and ciprofloxacin. Examples of corticosteroid enemas include hydrocortisone enema (Cortenema, generics). Ulcerative Colitis, continuation-approve if the patient has had a response to therapy.</p>
<b>Indications</b>	All FDA-approved Indications.
<b>Off-Label Uses</b>	N/A
<b>Part B Prerequisite</b>	No

## ZYTIGA

### Products Affected

- *abiraterone oral tablet 250 mg, 500 mg*

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	N/A
Age Restrictions	18 years and older
Prescriber Restrictions	N/A
Coverage Duration	Authorization will be for 1 year
Other Criteria	<p>Prostate Cancer-Metastatic, Castration-Resistant (mCRPC)-Approve if abiraterone is being used in combination with prednisone or dexamethasone and the medication is used concurrently used with a gonadotropin-releasing hormone (GnRH) agonist, or the medication is concurrently used with Firmagon or the patient has had a bilateral orchiectomy. Prostate cancer-metastatic, castration-sensitive (mCSPC)-approve if the medication is used in combination with prednisone and the medication is concurrently used with a gonadotropin-releasing hormone agonist or concurrently used with Firmagon or the patient has had a bilateral orchiectomy. Prostate Cancer - Regional Risk Group - Approve if the patient meets all of the following criteria (A, B, and C): A) abiraterone is used in combination with prednisone AND B) Patient has regional lymph node metastases and no distant metastases AND C) Patient meets one of the following criteria (i, ii or iii): i. abiraterone with prednisone is used in combination with gonadotropin-releasing hormone (GnRH) agonist OR ii. Patient has had a bilateral orchiectomy OR iii. the medication is used in combination with Firmagon. Prostate cancer-very-high-risk-group-approve if according to the prescriber the patient is in the very-high-risk group, the medication will be used in combination with prednisone, the medication will be used in combination with external beam radiation therapy and the</p>

<b>PA Criteria</b>	<b>Criteria Details</b>
	<p>patient meets one of the following criteria (i, ii or iii): i. abiraterone is used in combination with gonadotropin-releasing hormone (GnRH) agonist OR ii. Patient has had a bilateral orchiectomy OR iii. the medication is used in combination with Firmagon. Prostate cancer-radical prostatectomy-approve if the medication is used in combination with prednisone, the patient has prostate specific antigen (PSA) persistence or recurrence following radical prostatectomy, patient has pelvic recurrence, the medication will be used concurrently with GnRH agonist, Firmagon or the patient has had a bilateral orchiectomy.</p>
<b>Indications</b>	All FDA-approved Indications, Some Medically-accepted Indications.
<b>Off-Label Uses</b>	Prostate Cancer-Regional Risk Group, Prostate cancer-very-high-risk group, Prostate cancer-radical prostatectomy
<b>Part B Prerequisite</b>	No

## PART B VERSUS PART D

### Products Affected

- ABELCET
- *acetylcysteine*
- *acyclovir sodium intravenous solution*
- *albuterol sulfate inhalation solution for nebulization 0.63 mg/3 ml, 1.25 mg/3 ml, 2.5 mg /3 ml (0.083 %), 2.5 mg/0.5 ml*
- *amphotericin b*
- *aprepitant*
- *arformoterol*
- *azathioprine oral tablet 50 mg*
- *budesonide inhalation suspension for nebulization 0.25 mg/2 ml, 0.5 mg/2 ml, 1 mg/2 ml*
- CLINIMIX 5%/D15W SULFITE FREE
- CLINIMIX 4.25%/D10W SULF FREE
- CLINIMIX 4.25%/D5W SULFIT FREE
- CLINIMIX 5% -D20W(SULFITE-FREE)
- *cromolyn inhalation*
- *cyclophosphamide oral capsule*
- CYCLOPHOSPHAMIDE ORAL TABLET
- *cyclosporine modified*
- *cyclosporine oral capsule*
- *dronabinol*
- ENGERIX-B (PF)
- ENGERIX-B PEDIATRIC (PF)
- ENVARSUS XR
- *everolimus (immunosuppressive)*
- *formoterol fumarate*
- *gengraf*
- *granisetron hcl oral*
- HEPLISAV-B (PF)
- *intralipid intravenous emulsion 20 %*
- *ipratropium bromide inhalation*
- *ipratropium-albuterol*
- JYLAMVO
- JYNNEOS (PF)
- *methotrexate sodium*
- *methotrexate sodium (pf) injection solution*
- *methylprednisolone oral tablet*
- *mycophenolate mofetil*
- *mycophenolate sodium*
- MYHIBBIN
- *ondansetron hcl oral solution*
- *ondansetron hcl oral tablet 4 mg, 8 mg*
- *ondansetron oral tablet, disintegrating 4 mg, 8 mg*
- *pentamidine inhalation*
- PLENAMINE
- PREHEVBRIO (PF)
- *premasol 10 %*
- PROGRAF ORAL GRANULES IN PACKET
- PULMOZYME
- RECOMBIVAX HB (PF)
- *sirolimus*
- *tacrolimus oral capsule*
- *travasol 10 %*
- TROPHAMINE 10 %
- VARUBI
- XGEVA

### Details

This drug may be covered under Medicare Part B or D depending upon the circumstances. Information may need to be submitted describing the use and setting of the drug to make the determination.

## Index

### A

ABELCET.....	343
abiraterone oral tablet 250 mg, 500 mg .	341, 342
acetylcysteine.....	343
ACTEMRA ACTPEN .....	1
ACTEMRA SUBCUTANEOUS .....	1
ACTIMMUNE.....	2
acyclovir sodium intravenous solution ...	343
acyclovir topical ointment.....	3
ADBRY SUBCUTANEOUS SYRINGE ...	6
ADEMPAS .....	7
AIMOVIG AUTOINJECTOR.....	8
AKEEGA .....	9
albuterol sulfate inhalation solution for nebulization 0.63 mg/3 ml, 1.25 mg/3 ml, 2.5 mg /3 ml (0.083 %), 2.5 mg/0.5 ml	343
alcohol pads .....	60
ALECENSA.....	10, 11
alosetron.....	12
ALUNBRIG ORAL TABLET 180 MG, 30 MG, 90 MG.....	14
ALUNBRIG ORAL TABLETS,DOSE PACK.....	14
alyq.....	209
ambrisentan.....	30
amikacin injection solution 500 mg/2 ml.	15, 16
amphotericin b .....	343
ampicillin sodium injection recon soln 1 gram, 10 gram, 125 mg.....	15, 16
ampicillin-sulbactam injection.....	15, 16
aprepitant.....	343
ARCALYST .....	18
arformoterol .....	343
ARIKAYCE.....	19
armodafinil.....	165
AUGTYRO.....	21
AVONEX INTRAMUSCULAR PEN INJECTOR KIT .....	22
AVONEX INTRAMUSCULAR SYRINGE KIT .....	22

AYVAKIT .....	23
azathioprine oral tablet 50 mg.....	343
azithromycin intravenous.....	15, 16
aztreonam.....	15, 16
<b>B</b>	
BALVERSA .....	24
BELBUCA.....	149, 150
BELSOMRA.....	68
BENLYSTA SUBCUTANEOUS.....	25
benztropine oral .....	102
BESREMI .....	26
BETASERON SUBCUTANEOUS KIT ..	27
bexarotene .....	28, 29
BICILLIN L-A.....	15, 16
bosentan .....	30
BOSULIF ORAL CAPSULE 100 MG, 50 MG .....	31
BOSULIF ORAL TABLET 100 MG, 400 MG, 500 MG.....	31
BRAFTOVI.....	32
BRUKINSA .....	33
budesonide inhalation suspension for nebulization 0.25 mg/2 ml, 0.5 mg/2 ml, 1 mg/2 ml.....	343
buprenorphine transdermal patch....	149, 150
BYDUREON BCISE.....	95
BYETTA SUBCUTANEOUS PEN INJECTOR 10 MCG/DOSE(250 MCG/ML) 2.4 ML, 5 MCG/DOSE (250 MCG/ML) 1.2 ML.....	95
<b>C</b>	
CABLIVI INJECTION KIT .....	35
CABOMETYX .....	36
CALQUENCE .....	37, 38
CALQUENCE (ACALABRUTINIB MAL) .....	37, 38
CAPRELSA ORAL TABLET 100 MG, 300 MG.....	39
carglumic acid.....	40
CAYSTON.....	41
cefoxitin .....	15, 16
ceftazidime.....	15, 16



cefuroxime sodium injection recon soln 750 mg .....	15, 16	cyclosporine oral capsule .....	343
cefuroxime sodium intravenous recon soln 1.5 gram .....	15, 16	CYLTEZO(CF) PEN .....	4, 5
CHEMET .....	42	CYLTEZO(CF) PEN CROHN'S-UC-HS4, 5	
CIBINQO .....	43	CYLTEZO(CF) PEN PSORIASIS-UV ..	4, 5
CIMZIA POWDER FOR RECONST .....	44, 45	CYLTEZO(CF) SUBCUTANEOUS	
CIMZIA SUBCUTANEOUS SYRINGE		SYRINGE KIT 10 MG/0.2 ML, 20	
KIT 400 MG/2 ML (200 MG/ML X 2) .....	44, 45	MG/0.4 ML, 40 MG/0.4 ML, 40 MG/0.8	
cinacalcet .....	46	ML .....	4, 5
CINRYZE .....	34	CYSTAGON .....	55
ciprofloxacin in 5 % dextrose intravenous piggyback 200 mg/100 ml .....	15, 16	CYSTARAN .....	54
clindamycin in 5 % dextrose .....	15, 16	<b>D</b>	
clindamycin phosphate injection .....	15, 16	dalfampridine .....	56
CLINIMIX 5%/D15W SULFITE FREE .....	343	DAURISMO ORAL TABLET 100 MG, 25	
CLINIMIX 4.25%/D10W SULF FREE ..	343	MG .....	57
CLINIMIX 4.25%/D5W SULFIT FREE .....	343	deferasirox .....	58
CLINIMIX 5%-D20W(SULFITE-FREE)		deferiprone .....	59
.....	343	DIACOMIT .....	64
clobazam oral suspension .....	47	diazepam intensol .....	101
clobazam oral tablet .....	47	diazepam oral solution 5 mg/5 ml (1 mg/ml)	
clorazepate dipotassium oral tablet 15 mg, 3.75 mg, 7.5 mg .....	101	.....	101
colistin (colistimethate na) .....	15, 16	diazepam oral tablet .....	101
COMETRIQ ORAL CAPSULE 100		diclofenac sodium topical gel 3 % .....	244
MG/DAY(80 MG X1-20 MG X1), 140		dimethyl fumarate oral capsule, delayed	
MG/DAY(80 MG X1-20 MG X3), 60		release(dr/ec) 120 mg, 120 mg (14)- 240	
MG/DAY (20 MG X 3/DAY) .....	48	mg (46), 240 mg .....	65
COPIKTRA .....	49	DOPTELET (10 TAB PACK) .....	66
COSENTYX (2 SYRINGES) .....	50, 51	DOPTELET (15 TAB PACK) .....	66
COSENTYX PEN (2 PENS) .....	50, 51	DOPTELET (30 TAB PACK) .....	66
COSENTYX SUBCUTANEOUS		dotti .....	106
SYRINGE 75 MG/0.5 ML .....	50, 51	doxy-100 .....	15, 16
COSENTYX UNOREADY PEN .....	50, 51	dronabinol .....	343
COTELLIC .....	52	droxidopa .....	67
CRESEMBA ORAL .....	53	DUPIXENT PEN SUBCUTANEOUS PEN	
cromolyn inhalation .....	343	INJECTOR 200 MG/1.14 ML, 300 MG/2	
cyclobenzaprine oral tablet 10 mg, 5 mg .....	103	ML .....	69
cyclophosphamide oral capsule .....	343	DUPIXENT SYRINGE SUBCUTANEOUS	
CYCLOPHOSPHAMIDE ORAL TABLET		SYRINGE 100 MG/0.67 ML, 200	
.....	343	MG/1.14 ML, 300 MG/2 ML .....	69
cyclosporine modified .....	343	<b>E</b>	
		ELIGARD .....	96
		ELIGARD (3 MONTH) .....	96
		ELIGARD (4 MONTH) .....	96
		ELIGARD (6 MONTH) .....	96
		EMGALITY PEN .....	70

EMGALITY SYRINGE  
 SUBCUTANEOUS SYRINGE 120  
 MG/ML ..... 70  
 ENBREL MINI..... 71, 72  
 ENBREL SUBCUTANEOUS SOLUTION  
 ..... 71, 72  
 ENBREL SUBCUTANEOUS SYRINGE71,  
 72  
 ENBREL SURECLICK..... 71, 72  
 ENGERIX-B (PF)..... 343  
 ENGERIX-B PEDIATRIC (PF)..... 343  
 ENVARSUS XR..... 343  
 EPIDIOLEX..... 74  
 EPRONTIA..... 282  
 ERIVEDGE..... 77  
 ERLEADA ORAL TABLET 240 MG, 60  
 MG ..... 78  
 erlotinib oral tablet 100 mg, 150 mg, 25 mg  
 ..... 79  
 ertapenem ..... 15, 16  
 estradiol oral..... 106  
 estradiol transdermal patch semiweekly . 106  
 estradiol transdermal patch weekly..... 106  
 estradiol-norethindrone acet..... 106  
 everolimus (antineoplastic) oral tablet..... 80  
 everolimus (antineoplastic) oral tablet for  
 suspension 2 mg, 3 mg, 5 mg..... 80  
 everolimus (immunosuppressive) ..... 343

**F**  
 FASENRA PEN..... 81  
 FASENRA SUBCUTANEOUS SYRINGE  
 10 MG/0.5 ML, 30 MG/ML ..... 81  
 fentanyl citrate buccal lozenge on a handle  
 ..... 285  
 fentanyl transdermal patch 72 hour 100  
 mcg/hr, 12 mcg/hr, 25 mcg/hr, 50 mcg/hr,  
 75 mcg/hr ..... 283, 284  
 fingolimod..... 82  
 FINTEPLA..... 83  
 FIRMAGON KIT W DILUENT SYRINGE  
 ..... 84  
 fluconazole in nacl (iso-osm) intravenous  
 piggyback 200 mg/100 ml, 400 mg/200  
 ml ..... 17

formoterol fumarate ..... 343  
 FOTIVDA ..... 85  
 FRUZAQLA ORAL CAPSULE 1 MG, 5  
 MG ..... 86, 87  
 FULPHILA ..... 88, 89  
 fyavolv ..... 106

**G**  
 gabapentin oral tablet extended release 24 hr  
 300 mg, 600 mg ..... 97  
 GATTEX 30-VIAL..... 90  
 GAUZE PAD TOPICAL BANDAGE 2 X 2  
 ..... 61  
 GAVRETO ..... 91  
 gefitinib ..... 92  
 gengraf ..... 343  
 gentamicin in nacl (iso-osm) intravenous  
 piggyback 100 mg/100 ml, 60 mg/50 ml,  
 80 mg/100 ml, 80 mg/50 ml..... 15, 16  
 gentamicin injection solution 40 mg/ml .. 15,  
 16  
 GILOTRIF ..... 93  
 glatiramer subcutaneous syringe 20 mg/ml,  
 40 mg/ml ..... 94  
 glatopa subcutaneous syringe 20 mg/ml, 40  
 mg/ml ..... 94  
 glutamine (sickle cell)..... 73  
 granisetron hcl oral ..... 343

**H**  
 HEPLISAV-B (PF) ..... 343  
 HUMIRA (PREFERRED NDCS  
 STARTING WITH 00074)  
 SUBCUTANEOUS SYRINGE KIT 40  
 MG/0.8 ML ..... 4, 5  
 HUMIRA PEN (PREFERRED NDCS  
 STARTING WITH 00074)..... 4, 5  
 HUMIRA(CF) (PREFERRED NDCS  
 STARTING WITH 00074)  
 SUBCUTANEOUS SYRINGE KIT 10  
 MG/0.1 ML, 20 MG/0.2 ML, 40 MG/0.4  
 ML..... 4, 5  
 HUMIRA(CF) PEN (PREFERRED NDCS  
 NDCS STARTING WITH 00074)  
 SUBCUTANEOUS PEN INJECTOR  
 KIT 40 MG/0.4 ML, 80 MG/0.8 ML.. 4, 5

HUMIRA(CF) PEN CROHNS-UC-HS (PREFERRED NDCS NDCS STARTING WITH 00074).....	4, 5	ivermectin oral .....	123
HUMIRA(CF) PEN PEDIATRIC UC (ONLY NDCS STARTING WITH 00074) .....	4, 5	IWILFIN .....	125
HUMIRA(CF) PEN PSOR-UV-ADOL HS (PREFERRED NDCS NDCS STARTING WITH 00074).....	4, 5	<b>J</b>	
hydromorphone oral tablet extended release 24 hr .....	149, 150	JAKAFI.....	126, 127
hydroxyzine hcl oral tablet.....	104	JAYPIRCA ORAL TABLET 100 MG, 50 MG .....	128, 129
<b>I</b>		jinteli .....	106
IBRANCE .....	107, 108	JYLAMVO .....	343
icatibant.....	109	JYNNEOS (PF).....	343
ICLUSIG.....	110	<b>K</b>	
IDHIFA .....	111	KALYDECO.....	130
imatinib oral tablet 100 mg, 400 mg	112, 113	KERENDIA .....	131, 132
IMBRUVICA ORAL CAPSULE 140 MG, 70 MG .....	114, 115	KESIMPTA PEN .....	133
IMBRUVICA ORAL SUSPENSION....	114, 115	KISQALI FEMARA CO-PACK ORAL TABLET 200 MG/DAY(200 MG X 1)- 2.5 MG, 400 MG/DAY(200 MG X 2)-2.5 MG, 600 MG/DAY(200 MG X 3)-2.5 MG .....	134
IMBRUVICA ORAL TABLET 140 MG, 280 MG, 420 MG.....	114, 115	KISQALI ORAL TABLET 200 MG/DAY (200 MG X 1), 400 MG/DAY (200 MG X 2), 600 MG/DAY (200 MG X 3).....	134
imipenem-cilastatin.....	15, 16	KOSELUGO .....	136, 137
INBRIJA INHALATION CAPSULE, W/INHALATION DEVICE .....	116	KRAZATI.....	138
INGREZZA.....	117	<b>L</b>	
INGREZZA INITIATION PK(TARDIV) .....	117	lapatinib.....	139, 140
INGREZZA SPRINKLE .....	117	LEDIPASVIR-SOFOSBUVIR .....	141
INLYTA ORAL TABLET 1 MG, 5 MG	119	lenalidomide.....	142, 143
INPEFA.....	120	LENVIMA ORAL CAPSULE 10 MG/DAY (10 MG X 1), 12 MG/DAY (4 MG X 3), 14 MG/DAY(10 MG X 1-4 MG X 1), 18 MG/DAY (10 MG X 1-4 MG X2), 20 MG/DAY (10 MG X 2), 24 MG/DAY(10 MG X 2-4 MG X 1), 4 MG, 8 MG/DAY (4 MG X 2).....	144, 145
INQOVI .....	121	leuprolide subcutaneous kit.....	96
INREBIC.....	122	levofloxacin in d5w intravenous piggyback 500 mg/100 ml, 750 mg/150 ml.....	15, 16
INSULIN SYRINGE-NEEDLE U-100 SYRINGE 0.3 ML 29 GAUGE, 1 ML 29 GAUGE X 1/2.....	63	LIBERVANT .....	146
INSULIN SYRINGES (NON-PREFERRED BRANDS) SYRINGE 1 ML 29 GAUGE X 1/2.....	62	lidocaine topical adhesive patch,medicated 5 % .....	147
intralipid intravenous emulsion 20 % .....	343	lidocan iii .....	147
ipratropium bromide inhalation .....	343	linezolid in dextrose 5%.....	15, 16
ipratropium-albuterol .....	343	LIVTENCITY .....	148
		LONSURF .....	151

lorazepam intensol ..... 101  
 lorazepam oral tablet 0.5 mg, 1 mg, 2 mg  
 ..... 101  
 LORBRENA ORAL TABLET 100 MG, 25  
 MG ..... 152  
 LUMAKRAS ORAL TABLET 120 MG,  
 320 MG ..... 153  
 LUPRON DEPOT..... 154, 155  
 lyllana..... 106  
 LYNPARZA ..... 156, 157  
 LYTGABI ORAL TABLET 12 MG/DAY  
 (4 MG X 3), 16 MG/DAY (4 MG X 4),  
 20 MG/DAY (4 MG X 5) ..... 158

**M**  
 MAVYRET ORAL PELLETS IN PACKET  
 ..... 159  
 MAVYRET ORAL TABLET..... 159  
 megestrol oral suspension 400 mg/10 ml (40  
 mg/ml), 625 mg/5 ml (125 mg/ml) ..... 160  
 megestrol oral tablet..... 160  
 MEKINIST ORAL RECON SOLN 161, 162  
 MEKINIST ORAL TABLET 0.5 MG, 2  
 MG ..... 161, 162  
 MEKTOVI ..... 163  
 memantine oral capsule, sprinkle, er 24hr 164  
 memantine oral solution..... 164  
 memantine oral tablet..... 164  
 meropenem intravenous recon soln 1 gram,  
 500 mg ..... 15, 16  
 methadone oral solution 10 mg/5 ml, 5 mg/5  
 ml ..... 149, 150  
 methadone oral tablet 10 mg, 5 mg. 149, 150  
 methotrexate sodium ..... 343  
 methotrexate sodium (pf) injection solution  
 ..... 343  
 methylprednisolone oral tablet..... 343  
 metronidazole in nacl (iso-os)..... 15, 16  
 metyrosine..... 208  
 mifepristone oral tablet 300 mg ..... 135  
 mimvey ..... 106  
 modafinil oral tablet 100 mg, 200 mg..... 165  
 morphine oral tablet extended release.... 149,  
 150  
 MOUNJARO ..... 95

moxifloxacin-sod.chloride(iso)..... 15, 16  
 mycophenolate mofetil..... 343  
 mycophenolate sodium ..... 343  
 MYFEMBREE..... 166  
 MYHIBBIN ..... 343

**N**  
 nafcillin injection ..... 15, 16  
 NAMZARIC ..... 164  
 NAYZILAM ..... 167  
 NERLYNX ..... 168  
 NEXLETOL..... 169, 170  
 NEXLIZET ..... 171, 172  
 nilutamide ..... 173  
 NINLARO..... 174  
 nitisinone..... 175  
 NIVESTYM..... 176  
 norethindrone ac-eth estradiol oral tablet  
 0.5-2.5 mg-mcg, 1-5 mg-mcg ..... 106  
 NUBEQA..... 179  
 NUCALA SUBCUTANEOUS AUTO-  
 INJECTOR..... 180, 181  
 NUCALA SUBCUTANEOUS RECON  
 SOLN ..... 180, 181  
 NUCALA SUBCUTANEOUS SYRINGE  
 100 MG/ML, 40 MG/0.4 ML ..... 180, 181  
 NUEDEXTA..... 182  
 NUPLAZID..... 183  
 NURTEC ODT ..... 184  
 NYVEPRIA ..... 185, 186

**O**  
 OCALIVA..... 187, 188  
 octreotide acetate injection solution..... 189  
 ODOMZO ..... 190  
 OFEV ..... 191  
 OGSIVEO ORAL TABLET 100 MG, 150  
 MG, 50 MG..... 192  
 OJEMDA ORAL SUSPENSION FOR  
 RECONSTITUTION ..... 193  
 OJEMDA ORAL TABLET 400 MG/WEEK  
 (100 MG X 4), 500 MG/WEEK (100 MG  
 X 5), 600 MG/WEEK (100 MG X 6) . 193  
 OJJAARA ..... 194  
 OMNITROPE ..... 98, 99, 100  
 ondansetron hcl oral solution ..... 343

ondansetron hcl oral tablet 4 mg, 8 mg... 343	penicillin g potassium injection recon soln 20 million unit..... 15, 16
ondansetron oral tablet,disintegrating 4 mg, 8 mg ..... 343	penicillin g sodium..... 15, 16
ONUREG..... 195	pentamidine inhalation..... 343
OPSUMIT..... 196	phenobarbital..... 105
OPSYNVI..... 197	pimecrolimus..... 280
ORENCIA CLICKJECT..... 198	PIQRAY ORAL TABLET 200 MG/DAY (200 MG X 1), 250 MG/DAY (200 MG X1-50 MG X1), 300 MG/DAY (150 MG X 2) ..... 210
ORENCIA SUBCUTANEOUS SYRINGE 125 MG/ML, 50 MG/0.4 ML, 87.5 MG/0.7 ML ..... 198	pirfenidone oral capsule..... 211
ORGOVYX..... 199	pirfenidone oral tablet 267 mg, 801 mg.. 211
ORKAMBI ORAL GRANULES IN PACKET ..... 200	PLEGRIDY SUBCUTANEOUS PEN INJECTOR 125 MCG/0.5 ML ..... 212
ORKAMBI ORAL TABLET ..... 200	PLEGRIDY SUBCUTANEOUS SYRINGE 125 MCG/0.5 ML ..... 212
ORSERDU ORAL TABLET 345 MG, 86 MG ..... 201	PLENAMINE ..... 343
OTEZLA..... 202	POMALYST..... 213
OTEZLA STARTER ORAL TABLETS,DOSE PACK 10 MG (4)- 20 MG (51), 10 MG (4)-20 MG (4)-30 MG (47)..... 202	posaconazole oral tablet,delayed release (dr/ec)..... 214
oxacillin..... 15, 16	PREHEVBRIO (PF) ..... 343
oxacillin in dextrose(iso-osm)..... 15, 16	premasol 10 %..... 343
OXERVATE..... 203	PREVYMIS ORAL ..... 215
OXYCONTIN ORAL TABLET,ORAL ONLY,EXT.REL.12 HR 10 MG, 15 MG, 20 MG, 30 MG, 40 MG, 60 MG, 80 MG ..... 149, 150	PRIVIGEN..... 124
OZEMPIC SUBCUTANEOUS PEN INJECTOR 0.25 MG OR 0.5 MG (2 MG/3 ML), 1 MG/DOSE (4 MG/3 ML), 2 MG/DOSE (8 MG/3 ML) ..... 95	PROCRIT INJECTION SOLUTION 10,000 UNIT/ML, 2,000 UNIT/ML, 20,000 UNIT/ML, 3,000 UNIT/ML, 4,000 UNIT/ML, 40,000 UNIT/ML ..... 75, 76
<b>P</b>	PROGRAF ORAL GRANULES IN PACKET ..... 343
PANRETIN..... 204	PROLASTIN-C INTRAVENOUS SOLUTION..... 13
pazopanib ..... 310, 311	PROLIA ..... 216, 217
PEMAZYRE..... 205	PROMACTA ..... 218
PEN NEEDLES (NON-PREFERRED BRANDS) NEEDLE 29 GAUGE X 1/2 ..... 62	promethazine oral..... 104
penicillamine oral tablet..... 206	PULMOZYME ..... 343
PENICILLIN G POT IN DEXTROSE INTRAVENOUS PIGGYBACK 2 MILLION UNIT/50 ML, 3 MILLION UNIT/50 ML..... 15, 16	pyrimethamine ..... 219
	<b>Q</b>
	QINLOCK..... 220
	QULIPTA ..... 221
	<b>R</b>
	RADICAVA ORS STARTER KIT SUSP ..... 222
	RECOMBIVAX HB (PF)..... 343



RELEUKO SUBCUTANEOUS.....	223	SODIUM OXYBATE (PREFERRED	
REPATHA .....	224	NDCS STARTING WITH 00054) .....	329
REPATHA PUSHTRONEX.....	224	sodium phenylbutyrate.....	207
REPATHA SURECLICK.....	224	SOFOSBUVIR-VELPATASVIR.....	243
RETACRIT .....	75, 76	SOMAVERT.....	245
RETEVMO ORAL CAPSULE 40 MG, 80		sorafenib.....	246, 247
MG .....	225	SOTYKTU .....	248
RETEVMO ORAL TABLET 120 MG, 160		SPRYCEL ORAL TABLET 100 MG, 140	
MG, 40 MG, 80 MG .....	225	MG, 20 MG, 50 MG, 70 MG, 80 MG	249
REVLIMID .....	142, 143	STELARA SUBCUTANEOUS	
REZDIFFRA.....	226	SOLUTION.....	250, 251
REZLIDHIA .....	227	STELARA SUBCUTANEOUS SYRINGE	
REZUROCK.....	228	45 MG/0.5 ML, 90 MG/ML .....	250, 251
riluzole .....	229	STIVARGA .....	252, 253
RINVOQ ORAL TABLET EXTENDED		STREPTOMYCIN.....	15, 16
RELEASE 24 HR 15 MG, 30 MG, 45		SUCRAID .....	254, 255
MG .....	230	sunitinib malate.....	256, 257
roflumilast.....	231	SYMDEKO.....	258
ROZLYTREK ORAL CAPSULE 100 MG,		SYMLINPEN 120.....	259
200 MG .....	232	SYMLINPEN 60.....	259
ROZLYTREK ORAL PELLETS IN		SYMPAZAN.....	47
PACKET .....	232	<b>T</b>	
RUBRACA .....	233, 234	TABRECTA .....	260
rufinamide.....	235	tacrolimus oral capsule .....	343
RYBELSUS .....	95	tacrolimus topical.....	280
RYDAPT.....	236	tadalafil (pulm. hypertension).....	209
<b>S</b>		tadalafil oral tablet 2.5 mg, 5 mg .....	261
sajazir .....	109	TAFINLAR ORAL CAPSULE .....	263, 264
sapropterin.....	237	TAFINLAR ORAL TABLET FOR	
SCEMBLIX ORAL TABLET 100 MG, 20		SUSPENSION .....	263, 264
MG, 40 MG.....	238	TAGRISSO .....	265
SIGNIFOR .....	239	TALZENNA .....	266
sildenafil (pulmonary arterial hypertension)		TASIGNA ORAL CAPSULE 150 MG, 200	
oral tablet .....	209	MG, 50 MG.....	267, 268
sirolimus.....	343	tazarotene topical cream 0.1 % .....	269
SIRTURO .....	240	tazarotene topical gel .....	269
SKYRIZI SUBCUTANEOUS PEN		tazicef injection.....	15, 16
INJECTOR.....	241, 242	TAZVERIK.....	270
SKYRIZI SUBCUTANEOUS SYRINGE		TEFLARO.....	15, 16
150 MG/ML .....	241, 242	TEPMETKO .....	271
SKYRIZI SUBCUTANEOUS WEARABLE		teriflunomide.....	20
INJECTOR 180 MG/1.2 ML (150		TERIPARATIDE SUBCUTANEOUS PEN	
MG/ML), 360 MG/2.4 ML (150 MG/ML)		INJECTOR 20 MCG/DOSE	
.....	241, 242	(620MCG/2.48ML).....	272, 273

testosterone cypionate.....	118	TYENNE AUTOINJECTOR.....	1
testosterone enanthate .....	118	TYENNE SUBCUTANEOUS.....	1
testosterone transdermal gel in metered-dose pump 12.5 mg/ 1.25 gram (1 %), 20.25 mg/1.25 gram (1.62 %).....	177, 178	<b>U</b>	
testosterone transdermal gel in packet 1 % (25 mg/2.5gram), 1 % (50 mg/5 gram), 1.62 % (20.25 mg/1.25 gram), 1.62 % (40.5 mg/2.5 gram) .....	177, 178	UBRELVY.....	295
testosterone transdermal solution in metered pump w/app.....	177, 178	UPTRAVI ORAL TABLET.....	296
tetrabenazine oral tablet 12.5 mg, 25 mg	274	UPTRAVI ORAL TABLETS,DOSE PACK .....	296
THALOMID ORAL CAPSULE 100 MG, 150 MG, 200 MG, 50 MG .....	275, 276	<b>V</b>	
TIBSOVO .....	277	VALCHLOR.....	297
tigecycline .....	15, 16	VALTOCO .....	298
tobramycin in 0.225 % nacl .....	278	vancomycin intravenous recon soln 1,000 mg, 10 gram, 500 mg, 750 mg.....	15, 16
tobramycin inhalation .....	278	vancomycin oral capsule 125 mg, 250 mg .....	299
tobramycin sulfate injection solution..	15, 16	VANFLYTA.....	300
tolvaptan.....	279	VARUBI .....	343
topiramate oral capsule, sprinkle .....	282	VENCLEXTA ORAL TABLET 10 MG, 100 MG, 50 MG.....	301
topiramate oral tablet .....	282	VENCLEXTA STARTING PACK .....	301
torpenz.....	80	VERZENIO.....	302, 303
travasol 10 %.....	343	vigabatrin .....	304
TRELSTAR INTRAMUSCULAR SUSPENSION FOR RECONSTITUTION .....	286	vigadrone.....	304
TREMFYA SUBCUTANEOUS AUTO-INJECTOR.....	287	vigoder .....	304
TREMFYA SUBCUTANEOUS SYRINGE 100 MG/ML .....	287	VITRAKVI ORAL CAPSULE 100 MG, 25 MG.....	305
tretinoin topical .....	281	VITRAKVI ORAL SOLUTION .....	305
tridacaine ii.....	147	VIZIMPRO .....	306
trientine oral capsule 250 mg.....	288, 289	VONJO .....	307
TRIKAFTA ORAL GRANULES IN PACKET, SEQUENTIAL .....	290	voriconazole.....	17, 308
TRIKAFTA ORAL TABLETS, SEQUENTIAL.....	290	VOSEVI.....	309
TROPHAMINE 10 % .....	343	VOWST .....	312
TRULICITY .....	95	VUMERITY .....	313
TRUQAP.....	291, 292	VYNDAMAX.....	262
TUKYSA ORAL TABLET 150 MG, 50 MG .....	293	<b>W</b>	
TURALIO ORAL CAPSULE 125 MG..	294	WELIREG.....	314, 315
		<b>X</b>	
		XALKORI ORAL CAPSULE.....	316, 317
		XALKORI ORAL PELLETT 150 MG, 20 MG, 50 MG.....	316, 317
		XDEMVI .....	318
		XELJANZ ORAL SOLUTION .....	319, 320
		XELJANZ ORAL TABLET.....	319, 320
		XELJANZ XR .....	319, 320
		XERMELO .....	321



XGEVA..... 343  
 XIFAXAN ORAL TABLET 200 MG, 550  
 MG ..... 322  
 XOLAIR SUBCUTANEOUS AUTO-  
 INJECTOR 150 MG/ML, 300 MG/2 ML,  
 75 MG/0.5 ML ..... 323, 324  
 XOLAIR SUBCUTANEOUS RECON  
 SOLN ..... 323, 324  
 XOLAIR SUBCUTANEOUS SYRINGE  
 150 MG/ML, 300 MG/2 ML, 75 MG/0.5  
 ML..... 323, 324  
 XOSPATA ..... 325  
 XPOVIO ..... 326, 327  
 XTANDI ORAL CAPSULE..... 328  
 XTANDI ORAL TABLET 40 MG, 80 MG  
 ..... 328  
**Y**  
 YUFLYMA(CF) AI CROHN'S-UC-HS. 4, 5  
 YUFLYMA(CF) AUTOINJECTOR  
 SUBCUTANEOUS AUTO-INJECTOR,  
 KIT 40 MG/0.4 ML, 80 MG/0.8 ML.. 4, 5

YUFLYMA(CF) SUBCUTANEOUS  
 SYRINGE KIT 20 MG/0.2 ML, 40  
 MG/0.4 ML ..... 4, 5  
**Z**  
 ZEJULA ORAL TABLET..... 330  
 ZELBORAF ..... 331, 332  
 ZEPOSIA ..... 333  
 ZEPOSIA STARTER KIT (28-DAY).... 333  
 ZEPOSIA STARTER PACK (7-DAY).. 333  
 ZOLINZA ..... 334  
 ZONISADE..... 282  
 zonisamide ..... 282  
 ZTALMY ..... 335  
 ZURZUVAE ORAL CAPSULE 20 MG, 25  
 MG, 30 MG..... 336  
 ZYDELIG ..... 337  
 ZYKADIA ..... 338  
 ZYMFENTRA ..... 339, 340

**CareSource® MyCare Ohio (Medicare-Medicaid Plan) is a health plan that contracts with both Medicare and Ohio Medicaid to provide benefits of both programs to enrollees.**



H8452\_OH-MYC-M-3096364-V.1

ODM Approved: 8/15/2024