

# MEDICAL POLICY STATEMENT Wisconsin Marketplace

Policy Name & Number	Date Effective			
Breast Reduction Surgery-WI MP-MM-1653	12/01/2024			
Policy Type				
MEDICAL				

Medical Policy Statements are derived from literature based on and supported by clinical guidelines, nationally recognized utilization and technology assessment guidelines, other medical management industry standards, and published MCO clinical policy guidelines. Medically necessary services include, but are not limited to, those health care services or supplies that are proper and necessary for the diagnosis or treatment of disease, illness, or injury and without which the patient can be expected to suffer prolonged, increased, or new morbidity, impairment of function, dysfunction of a body organ or part, or significant pain and discomfort. These services meet the standards of good medical practice in the local area, are the lowest cost alternative, and are not provided mainly for the convenience of the member or provider. Medically necessary services also include those services defined in any Evidence of Coverage or Certificate of Coverage documents, Medical Policy Statements, Provider Manuals, Member Handbooks, and/or other plan policies and procedures.

Medical Policy Statements do not ensure an authorization or payment of services. Please refer to the plan contract (often referred to as the Evidence of Coverage or Certificate of Coverage) for the service(s) referenced in the Medical Policy Statement. Except as otherwise required by law, if there is a conflict between the Administrative Policy Statement and the plan contract, then the plan contract will be the controlling document used to make the determination.

According to the rules of Mental Health Parity Addiction Equity Act (MHPAEA), coverage for the diagnosis and treatment of a behavioral health disorder will not be subject to any limitations that are less favorable than the limitations that apply to medical conditions as covered under this policy.

#### **Table of Contents**

٩.	Subject	. 2
	Background	
	Definitions	
	Policy	
≣.	Conditions of Coverage	6
=.	Related Policies/Rules	6
Э.	Review/Revision History	6
4	References	6

Breast Reduction Surgery-WI MP-MM-1653 Effective Date: 12/01/2024



#### A. Subject

# **Breast Reduction Surgery**

## B. Background

Breast reduction surgery is performed for females or males for different indications. Women diagnosed with macromastia (excessively large breasts) seeking breast reduction typically present with complaints of a feeling of heaviness, chronic pain, and tension in the neck, shoulders, and upper back. Macromastia commonly causes permanent grooving and ulceration of the shoulder following years of wearing support bras to try to minimize symptoms. The physical and psychological symptoms of macromastia can significantly and negatively impact an individual's life and should be taken into consideration when evaluating surgical intervention.

Reduction mammoplasty is a surgical procedure that reduces the weight and volume of the breast. As much as two to five pounds of excess breast tissue is routinely removed during a reduction mammoplasty. Indications for surgery include chronic pain and skin symptoms, neuropathy, breast discomfort, physical impairment, and psychological symptoms that can be associated with poor self-esteem and loss of desire to engage in activities.

Gynecomastia is a benign proliferation of glandular tissue of the breast in males. This condition may be caused by androgen deficiency, medications, chronic medical conditions, tumors, or endocrine disorders. Depending on the cause of the tissue proliferation, surgical removal may be considered cosmetic or medically necessary. In order to evaluate the excess breast tissue, mammography and grading using the breast imaging reporting and data system (BI-RADS) may be appropriate. BI-RADS standardizes risk assessment and quality control for mammography by using categorizations and reporting/documentation standards. The 6 classifications include:

- BI-RADS 0 Incomplete evaluation with further imaging required including additional mammographic views (spot compression, magnification, or ultrasound).
- BI-RADS 1 Negative examination with no masses, suspicious calcifications, or areas of architectural distortion.
- BI-RADS 2 Consistent with benign findings (eg, secretory calcifications, simple cysts, fat-containing lesions, calcified fibroadenomas, implants, intramammary lymph nodes.
- BI-RADS 3 Probably benign (risk of malignancy below 2%) with a shortened follow-up interval to determine stability. Must include a non-palpable, circumscribed mass on a baseline mammogram; a focal asymmetry, which becomes less dense on spot compression images, or a solitary group of punctate calcifications.
- BI-RADS 4 A suspicious abnormality, which is subdivided further into subcategories:
  - (a) low probability of malignancy
  - (b) intermediate chance of malignancy (10-50%)
  - (c) high probability of malignancy (50-95%)

The MEDICAL Policy Statement detailed above has received due consideration as defined in the MEDICAL Policy Statement Policy and is approved.



- BI-RADS 5 Highly suggestive of malignancy (greater than 95%).
- BI-RADS 6 Pathology proven malignancy.

#### C. Definitions

- **Body Surface Area (BSA)** A metric used for physiologic measurements, pharmacologic dosing, and therapeutic calculations, including the Schnur Sliding Scale for breast reduction surgery.
- **Cosmetic Procedures** Procedures performed for aesthetic purposes that do not improve or restore physiologic function.
- Functional/Physical or Physiological Impairment Impairment causes deviation
  from the normal function of a tissue or organ. This results in a significantly limited,
  impaired or delayed capacity to move and coordinate actions or perform physical
  activities and is exhibited by difficulties in physical and motor tasks, independent
  movement, or performing basic life functions.
- **Gynecomastia** Enlargement of the male breast secondary to a proliferation of ductal, stromal, and/or fatty tissue.
- Intertriginous Rash Dermatitis occurring between juxtaposed folds of skin, caused by retention of moisture and warmth and providing an environment favoring overgrowth of normal skin micro-organisms.
- Kyphosis Over-curvature of the thoracic vertebrae (upper back) associated with degenerative diseases, such as arthritis, developmental problems, or with osteoporotic compression fractures of vertebral bodies.
- **Macromastia (Breast Hypertrophy)** An increase in the volume and weight of breast tissue relative to the general body habitus.
- **Mammography** An imaging technique that uses low-energy x-rays to examine breast tissue for diagnosis and screening.
- Symptomatic Breast Hypertrophy A syndrome of persistent neck and shoulder pain, shoulder grooving from brassiere straps, chronic intertriginous rash of the inframammary fold and/or frequent episodes of headache, backache, and upper extremity neuropathies caused by an increase in the volume and weight of breast tissue beyond normal proportions.
- **Schnur Sliding Scale** Used in calculating the amount of breast tissue to be removed in reduction mammoplasty (Appendix A).

#### D. Policy

- I. Common Ground Healthcare Cooperative considers breast reduction surgery for macromastia medically necessary when ALL the following criteria are met:
  - A. Member is 18 years or older or under 18 years with documented evidence that breasts have finished growing for a minimum of one year. Parental/guardian consent is required for members under age 18.
  - B. Breast size interferes with activities of daily living, as indicated by 1 or more of the following:
    - 1. arm numbness consistent with brachial plexus compression syndrome
    - 2. cervical pain



- 3. chronic breast pain
- 4. headaches
- 5. nipple position greater than 21 cm below suprasternal notch
- 6. persistent redness and erythema (intertrigo) below breasts
- 7. restriction of physical activity
- 8. severe bra strap grooving or ulceration of shoulder
- 9. shoulder pain
- 10. thoracic kyphosis
- 11. upper or lower back pain
- C. Preoperative evaluation by surgeon concludes that amount of breast tissue to be removed (by mass or volume) will provide a reasonable expectation of symptomatic relief.
- D. No evidence of breast cancer
  - 1. As evidenced by results of a physical exam completed by a physician within the last year if under 40 years of age.
  - 2. Women 40 to 54 years of age or older must have documentation of a mammogram negative for cancer performed within the year prior to the date of the planned breast reduction surgery.
  - 3. Women 55 years of age and older may switch to mammograms every 2 years.
- II. Breast reduction surgery following mastectomy to achieve symmetry is covered as part of the Women's Health and Cancer Rights Act (WHCRA). Please refer to the Common Ground Healthcare Cooperative Medical policy, *Breast Reconstruction Surgery*, for additional information.
- III. For males, cosmetic removal of excess breast tissue is not a covered benefit. Removal of breast tissue is considered medically necessary when a breast mass is expected or felt. This must be accompanied by documentation of a mammogram with a BI-RADS of 4 or higher in each breast that is surgically reduced.

## IV. Schnur Sliding Scale

The Schnur Sliding Scale is an evaluation tool used to determine the appropriate volume of tissue to be removed relative to a member's total body surface area (BSA). This estimation can be instrumental in determining whether breast reduction surgery is being planned for cosmetic reasons or as a medically necessary procedure.

- A. The weight of tissue to be removed from each breast must be above the 22<sup>nd</sup> percentile on the Schnur Sliding Scale (Appendix A below) based on the member's BSA.
- B. The BSA in meters squared (m²) is calculated using the Mosteller formula (square root of the result of height (inches) multiplied by weight (lbs) and divided by 3131).



Appendix A: Schnur Sliding Scale

Body Surface Area and Minimum Requirement for Breast Tissue Removal		
Body Surface Area (m²)	Grams per Breast of Minimum Breast Tissue to be Removed	
1.350-1.374	199	
1.375-1.399	208	
1.400-1.424	218	
1.425-1.449	227	
1.450-1.474	238	
1.475-1.499	249	
1.500-1.524	260	
1.525-1.549	272	
1.550-1.574	284	
1.575-1.599	297	
1.600-1.624	310	
1.625-1.649	324	
1.650-1.674	338	
1.675-1.699	354	
1.700-1.724	370	
1.725-1.749	386	
1.750-1.774	404	
1.775-1.799	422	
1.800-1.824	441	
1.825-1.849	461	
1.850-1.874	482	
1.875-1.899	504	
1.900-1.924	527	
1.925-1.949	550	
1.950-1.974	575	
1.975-1.999	601	
2.000-2.024	628	
2.025-2.049	657	
2.050-2.074	687	
2.075-2.099	717	



2.100-2.124	750
2.125-2.149	784
2.150-2.174	819
2.175-2.199	856
2.200-2.224	895
2.225-2.249	935
2.250-2.274	978
2.275-2.299	1022
2.300-2.324	1068
2.325-2.349	1117
2.350-2.374	1167
2.375-2.399	1219
2.400-2.424	1275
2.425-2.449	1333
2.450-2.474	1393
2.475-2.499	1455
2.500-2.524	1522
2.525-2.549	1590
2.550 or greater	1662
I and the second	

# E. Conditions of Coverage N/A

# F. Related Policies/Rules Breast Reconstruction Surgery

# G. Review/Revision History

	DATE	ACTION
Date Issued	08/14/2024	New policy. Approved at Committee.
Date Revised		
Date Effective	12/01/2024	
Date Archived		


#### H. References

- 1. Billa E, Kanakis GA, Goulis DG. Imaging in gynecomastia. *Andrology*. 2021;9(5):1444-1456. doi:10.1111/andr.13051
- 2. Biro FM, Chan YM. Normal puberty. UpToDate. Updated October 25, 2023. Accessed May 30, 2024. www.uptodate.com

The MEDICAL Policy Statement detailed above has received due consideration as defined in the MEDICAL Policy Statement Policy and is approved.



- 3. Briefing paper: plastic surgery for teenagers. American Society of Plastic Surgeons. Accessed May 30, 2024. www.plasticsurgery.org
- 4. Common Ground HealthCare Cooperative Certification of Coverage. Common Ground; 2025. Accessed May 30, 2024. www.caresource.com
- 5. Guliyeva G, Cheung JY, Avila FR, et al. Effect of reduction mammoplasty on pulmonary function tests: a systematic review. *Ann Plast Surg.* 2021;87(6):694-698. doi:10.1097/SAP.0000000000002834
- 6. Hansen J, Chang S. Overview of breast reduction. UpToDate. Updated April 19, 2023. Accessed May 30, 2024. www.uptodate.com
- 7. Holzmer SW, Lewis PG, Landau MJ, et al. Surgical management of gynecomastia: a comprehensive review of the literature. *Plast Reconstr Surg Glob Open*. 2020;8(10):e3161. doi:10.1097/GOX.000000000003161
- 8. Kanakis GA, Norkap L, Bang AK, et al. EAA clinical practice guidelines—gynecomastia evaluation and management. *Andrology*. 2019;7(6):778-793. doi:10.1111/andr.12636
- 9. Kimia R, Magee L, Caplan HS, et al. Trends in insurance coverage for adolescent reduction mammoplasty. *Am J Surg.* 2022;224(4):1068-1073. doi:10.1016/j.amjsurg.2022.07.030
- 10. Klement KA, Hijjawi BJ, Neuner J, et al. Discussion of preoperative mammography in women undergoing reduction mammoplasty. *Breast J*. 2019;25(3):439-44. doi:10.1111/tbj.13237
- Knox JA, Nelson DA, Latham KP, et al. Objective effects of breast reduction surgery on physical fitness. *Ann Plast Surg.* 2018;80(1):14-17. doi:10.1097/SAP.000000000001167
- 12. Lewin R, Liden M, Lundberg J, et al. Prospective evaluation of health after breast reduction surgery using the Breast-Q, Short-Form 36, Breast-Related Symptoms Questionnaire, and Modifed Breast Evaluation Form. *Ann Plast Surg.* 2019;83(2):143-151. doi:10.1097/SAP.000000000001849
- 13. Magny SJ, Shikhman R, Keppke AL. *Breast Imaging Reporting and Data System*. StatPearls Publishing; 2024. Accessed April 15, 2024. www.ncbi.nlm.nih.gov
- Morrison KA, Vernon R, Choi M, et al. Quantifying surgical complications for reduction mammaplasty in adolescents. *Plast Reconstr Surg.* 2023;151(3):376e-383e. doi:10.1097/PRS.0000000000009905
- 15. Nuzzi LC, Firriolo JM, Pike CM, et al. The effect of reduction mammoplasty on quality of life in adolescents with macromastia. *Pediatrics*. 2017;140(5):e20171103. doi:10.1542/peds.2017-1103
- Nuzzi LC, Pramanick T, Walsh LR, et al. Optimal timing for reduction mammoplasty in adolescents. *Plast Reconstr Surg.* 2020;146(6):1213-1220. doi:10.1097/PRS.0000000000007325
- 17. Patel K, Corcoran J. Breast reduction surgery in adolescents. *Pediatr Ann.* 2023;52(1):e31-e35. doi:10.3928/19382359-20221114-06
- 18. Perdikis G, Dillingham C, Boukovalas S, et al. American Society of Plastic Surgeons evidence-based clinical practice guideline revision: reduction mammaplasty. *Plast Reconstr Surg.* 2022;149(3):392e-409e. doi:10.1097/PRS.0000000000008860





- Perez-Panzano E, Gascon-Catalan A, Sousa-Dominguez R, et al. Reduction mammoplasty improves levels of anxiety, depression and body image satisfaction in patients with symptomatic macromastia in the short and long term. *J Psychosom Obstet Gynaecol.* 2017;38(4):268-275. doi:10.1080/0167482X.2016.1270936
- 20. Reduction mammoplasty: A-0274. MCG. 28th ed. Updated March 14, 2024. Accessed May 30, 2024. www.careweb.careguidelines.com
- 21. Reeves RA, Kaufman T. *Mammography*. StatPearls Publishing; 2023. Updated July 24, 2023. Accessed May 30, 2024. www.ncbi.nlm.nih.gov
- 22. Sears ED, Lu YT, Swiatek PR, et al. Use of preoperative mammography during evaluation for nononcologic breast reduction surgery. *JAMA Surg.* 2019;154(4):356-358. doi:10.1001/jamasurg.2018.4875
- 23. Xia TY, Scomacao I, Duraes E, et al. Aesthetic, quality-of-life, and clinical outcomes after inferior pedicle oncoplastic reduction mammoplasty. *Aesthetic Plast Surg.* 2023;47(3):905-911. doi:10.1007/s00266-023-03257-7
- 24. Zeiderman MR, Kelishadi SS, Tutela JP, et al. Reduction mammoplasty: intraoperative weight versus pathology weight and its implications. *Eplasty*. 2017;17:e32. Accessed June 3, 2024. www.pubmed.ncbi.nlm.nih.gov